Since the work of the heart bears a relation to the body weight, an attempt should be made to bring the bulk of the body, when it is excessive, within normal limits. Careful dieting to obtain this result should be carried out and the results noted. An abnormally high blood pressure can sometimes be modified with definite relief to the overloaded myocardium.

Much can be done in the secondary anginas when by meticulous cross-examination the cause of the lowered threshold to pain has been discovered. All possible sources of persistent low-grade infections should be investigated and radically treated—dental sepsis, sinus disease, abnormal intestinal conditions, pyelitis, chronic appendicitis, etc.

An attempt must be made to eliminate worry and make sure of sleep. For this a mixture of ammonium bromide and phenazonum is especially helpful. In the readjustment of the patient's life, factors giving rise to mental strain and irritations should be obviated as far as possible.

As my last word, may I lay stress again on the very great importance of an early diagnosis when the symptoms, though but slight and elusive, are both ominous and sinister. An accurate diagnosis *then*, followed by wise treatment, offers some chance of arresting, or at any rate of retarding, the development of the pathological changes responsible for angina.

THE ACTUAL PLACE AND FUNCTION OF THE TUBERCULOSIS DISPENSARY IN THE TUBERCULOSIS SCHEME.*

BY

SIR ROBERT PHILIP, M.D., LL.D.,

PROFESSOR OF TUBERCULOSIS IN THE UNIVERSITY OF EDINBURGH.

WHEN the council honoured me with the request to open this discussion I accepted with alacrity. The fascination of tuberculosis has held me a bondsman throughout my professional life. The range of interests is always widening. In offering a few thoughts for your consideration it will be my endeavour to do so briefly and simply, in the hope that it may stimulate members of the Conference to give us freely of their experience.

Forty Years Ago.

It is difficult to-day for the man in middle life—and still more for the younger man—to realize the attitude of mind towards tuberculosis which prevailed forty years ago and previously among both doctors and laymen. What was *then* accepted as care and treatment is *now* hardly imaginable.

The victim of tuberculosis—so-called consumption—was for the most part recognized too late. He was generally expected to die. The treatment meted out to him was empirical, unphysiological, and usually futile. The vain attempts at protection tended to hasten the end. With the death of the patient the interest that had been kindled by his illness was extinguished.

Speaking of the expectancy of life of the consumptive patient, Dr. C. J. B. Williams, in his treatise on *Pulmonary Consumption* (1871), says, with reference to the first ten years of his practice, 1830-40: "My general recollection of the histories of the developed disease at that time is that of distressing tragedies, in which no means used seemed to have any power to arrest the malady; and life was rarely prolonged beyond the limit of two years, assigned by Laënnec and Louis as the average duration of the life of a consumptive."

Conception of Tuberculosis Dispensary (1887).

Let me try to recall some of the circumstances and events which led to the remarkable alteration of outlook and action which now prevails.

It is some forty years since, following on the discovery of the essential cause of tuberculosis, its infective nature came to be admitted. While engaged in laboratory

work on tubercle, and attached to a general medical dispensary, it was borne in on me that acceptance of the infective nature of the disease carried with it far-reaching consequences. It seemed clear that, the infective nature of tuberculosis being granted, the principles which had been effectively applied in relation to other infective conditions were, with appropriate modifications, applicable to tuberculosis likewise.

Apart from a few special hospitals devoted to diseases of the chest, the chief avenue to care and treatment open to the poorer citizen suffering from tuberculosis forty years ago was by way of the out-patient department of a general hospital or similar medical institution. What happened daily in out-patient departments amounted to this—that the patient, suspected to be suffering from what was then spoken of as consumption, was examined for the presence of certain recognized signs of chest disease; when such were ascertained to be present, the sick man was accorded a prescription for his cough or other chief symptom, and commonly for cod-liver oil, and was invited to report himself from time to time for similar consideration and treatment. Occasionally he was passed on for admission to hospital, when such facilities were available. The number of special beds at that date was extremely scanty, and he was not very warmly welcomed in the general wards. The demand on beds was out of all proportion to the number of tuberculous applicants, with the result that even the small percentage which attained admission enjoyed the available benefit for a relatively short period. There was little serious effort to ensure anything more than amelioration or arrest of existing symptoms. The infective nature of the disease hardly came under review; nor were the facts of the illness linked in any close fashion with the conditions of the patient's home and household.

All this was radically wrong. It involved a disregard of the natural history of the disease, its infective origin, its widespread distribution, its lengthy course, its protean manifestations, its dependence on environmental conditions, its tractability when efficiently handled for a sufficient period, and many other facts which we now regard as commonplaces.

Reflection on such lines led me to press for a wider conception and fresh line of procedure. The proposal culminated in the establishment of the tuberculosis dispensary in 1887.

Purpose and Programme.

The essential purpose was the creation of a centre towards which patients suffering from this infective disease, and the friends of patients, and the public, could readily turn for every sort of direction and help, and from which there would radiate information of many kinds for the guidance of individual patients, households, and the community as a whole.

The original aims of the tuberculosis dispensary have been often detailed. They are so well known that on the present occasion there is little need to recapitulate them in detail. They constituted the first attempt to hunt down the infection, to discover the haunts and methods of the invader, and to raid him on his own territory. The principles and methods were incorporated in the well known circular of the Local Government Board of Scotland, dated March 10th, 1906, which reproduced *in extenso* a paper given by me to the International Congress at Paris in 1905. The principles and methods were frankly accepted by the Departmental Committee on Tuberculosis (1912), which adopted the programme of the Edinburgh Tuberculosis Dispensary as a model for the tuberculosis dispensaries of the country.

The Departmental Committee, in describing the functions of the tuberculosis dispensary, affirmed that it should serve as (1) receiving house and centre for diagnosis; (2) clearing house and centre for observation; (3) centre for curative treatment and supervision of ambulant cases; (4) centre for the methodical examination of contacts and the investigation of environmental conditions; (5) centre for continuous care of patients—so-called "after-care"; (6) information bureau and educational centre. The Departmental Committee placed the tuberculosis dispensary

* An address introductory to a discussion at the twelfth annual Conference of the National Association for the Prevention of Tuberculosis, Glasgow, July 1st, 1926.

as the first unit in the tuberculosis scheme and confirmed the view

"that the tuberculosis dispensary should be the common centre for the diagnosis and for the organization of treatment of tuberculosis in each area, at which the various bodies and persons connected with the campaign against tuberculosis should be brought together. The aim should be that no single case of tuberculosis should remain uncared for in the community. . . . The tuberculosis dispensary should be linked up to those institutions (hospitals, sanatoria, farm colonies, open-air schools) for which it will act as a clearing house."

Larger Results.

For the moment we are less concerned with the original purpose but rather with results which have emerged and developed from the continuous application of the methods proposed. Forty years constitute a period of sufficient length to enable us to appraise the significance of the activities.

At the outset you will allow me to say that, great as seemed the possibilities at the start, appreciation of their extent and value has grown with each succeeding year. A few of the more conspicuous developments may be cited.

1. *Widespread Distribution of Tuberculosis.*—The tuberculosis dispensary has demonstrated, as no other agency could, the extensive ramifications of tuberculosis throughout the area served by the dispensary. It has collected living proof of the universality of tuberculosis throughout the community. A comparison of maps showing the distribution of tuberculosis, taken at successive intervals throughout the area, demonstrates that the more an area is thoroughly investigated, the more is the population found to be permeated by the tuberculous taint. It has shown further that different portions of a given area differ in degree of saturation. It has thus thrown a flood of light on the influence of environment as a factor in the production and maintenance of tuberculosis.

2. *Spread through Households.*—By the systematic examination of households—the "march past" of the contacts—it has afforded incontrovertible evidence of the spread of the disease throughout a household from a primary source of infection.

3. *Infection in Childhood.*—It has emphasized the frequency with which tuberculosis is contracted in childhood. In household after household it has determined the existence of tuberculosis in the majority of the children involved. As many as fifteen cases have thus been detected by systematic search in one family circle alone.

4. *Heredity.*—Its investigations have dealt a serious blow at the doctrine of heredity—fondly cherished as that doctrine still is by some who are little acquainted with the natural history of the disease. The investigations offer strong support to the applicability of the views based on the observations of Bang and others in cattle, to the effect that the tuberculous mother very seldom produces offspring already tainted by tuberculosis, and that the offspring of such mother is saved from the risk of infection from the mother in proportion as it is protected by separation from her, while, conversely, the risk of early contraction of tuberculosis is accentuated in proportion as close contact with the infected mother—or with other infected member of the group—is maintained. Tuberculosis is thus seen to be a family or household disease, not because of hereditary transmission, but because of the abundant opportunities afforded for the transmission of infection from one infected individual to the other members. This risk exists in high degree in the earliest years of childhood.

5. *Recognition at Earliest Moment.*—The system of examining contacts has not only detected many cases that otherwise would have been missed. It has also afforded opportunity for the very early detection of infection as it occurs in successive additions to the household. It has shown how it is possible by careful observation to register in a growing family the actual advent of infection in successive children. In this way it has rendered possible the practice of anticipatory treatment—that is, of treatment, antigenic or other, which has in view the raising of the resistance of the infected individual with a view to preventing the worse consequences of such infection.

6. *Continuity of Observation.*—The dispensary has thrown a flood of light—by its continuity of treatment and care—on the natural history of tuberculosis. It has

enabled the observer to follow the child from the moment of recognized infection throughout a long period of years. In this way it has been possible in my own experience to observe the same infected individual throughout twenty to thirty years and longer.

7. *Natural History of the Disease.*—By this means it has been possible to construct a picture of the successive manifestations which most commonly repeat themselves in the evolution of tuberculosis. It has shown that in the vast majority of cases there is a gradual extension—just as in the experimental animal—from the point of primary infection by way of the lymphatic system, until the deeper viscera are successively involved. The dispensary has thus served to illuminate the relation of pulmonary tuberculosis and the more likely points of entrance of infection, and has relegated pulmonary disease to its proper place as constituting for the most part a relatively late visceral involvement.

8. *Continuity of Infection.*—Observations at the dispensary have shown, not only the continuity, but the varying manifestations of the tuberculous process in the same individual throughout a long series of years. It has revealed the ebb and flow in the progress of infection throughout a long life. It has shown how cases in the so-called third stage of pulmonary disease—with cavity formation which used to be regarded, and is still regarded by many, as the terminal picture—may yet live and go comparatively strong for thirty years or longer. It has illustrated how a cavity, especially when it occurs in young subjects, may, during the process of growth and development, gradually contract and finally disappear.

9. *Necessity for Long Supervision.*—By continued observation on ambulant patients and on those who have undergone sanatorium or other institutional treatment, the dispensary has proved how essential is the prolonged maintenance of such supervision, and how valueless may be the result of short periods of residential treatment—as, for example, the classic three months in a sanatorium, mere snippets in relation to the time limit necessary for the establishment of sound and effective arrest by the necessary histological changes.

10. *Economy in Treatment.*—Continuous, prolonged supervision at the dispensary has demonstrated how much can be done for tuberculous patients by ambulant treatment. From the records abundant proof may be had of how health may be maintained successfully without much expenditure. By means of weekly, fortnightly, or monthly guidance received at the dispensary, large numbers of patients may be kept on their feet and successfully treated without recourse to the sanatorium, or following on sanatorium treatment—and this at a minimum of cost.

11. *Housing: Reform in the Home.*—By a sound system of domiciliary visitation, the dispensary has succeeded in reforming homes and households, reducing the effects of unhealthy environment and recreating insanitary houses after the image of the sanatorium. In this direction the dispensary's educational value to a community through doctor and nurse can hardly be overestimated.

These examples of the far-reaching activities of the tuberculosis dispensary might be multiplied. They are cited merely as illustrations. Others will emerge in the course of the discussion from able men who, in town and county, have had many years' experience in developing and refining this significant instrument of preventive medicine.

What the Tuberculosis Dispensary should not be.

In drawing these desultory remarks to a close, a few words seem desirable on the other side of the picture—namely, what a tuberculosis dispensary should not be.

The establishment of the tuberculosis dispensary was a reaction from the comparatively ineffective out-patient department of the past. Splendid work has been effected by tuberculosis officers in different parts of the country; yet visits to not a few dispensaries, and extended inquiries in different directions, constrain me to the view that the ideal of the dispensary and its great possibilities are not realized everywhere.

It is unfortunately common to find that the tuberculosis dispensary concerns itself solely or chiefly with cases of

pulmonary tuberculosis, although these constitute for the most part a late manifestation of the infection. Even then the investigation of the patient lacks reality and thoroughness. The idea of using the pronounced case of infection as a clue to the earlier—the conception of hunting down the infection in the households of the area with a view to its eradication—is overlooked or ignored. The opportunity of the dispensary as a field of fresh observation in many directions is entirely lost. With so limited an outlook the work tends to become routine.

Such procedure cannot be expected to lead us very far towards the final goal. As I ventured to say on a previous occasion:

"The calm repose of the physician who, stethoscope in hand, decides for or against a cavity in the lung, and then contents himself with prescribing cod-liver oil, and perhaps a chronic pectoral mixture, must be disturbed. His cerebration and locomotion both require stimulation, so that he may be led to anticipate the disease by earlier diagnosis, and to determine the potentialities of contagion in the home. With the infection detected he must leave no stone unturned in the study of the patient's life-history and environment, which may throw light on causation or maintenance of disease. Chiefly must the physician concern himself with the health of the rest of the household. The whole family—or as many members as possible—must be scrutinized with scientific suspicion.

"The patient will not be sent home merely rich in weighty counsels. He will be followed to his home by doctor and nurse with trained eye and skilled finger, prepared to observe and sift every fact and to hold a critical 'march past' of the household. That is the way to raid the haunts of the tubercle bacillus effectively."

Searchlight Installation.

Ernest Renan compared the human conscience to a light-house with recurrent flashes of illumination which guide the uncertain mariner on the sea of life. I like to think of the tuberculosis dispensary as a searchlight installation in the vast tubercle-tainted plantations of our communal life.

In proportion as the installation is complete and the operators are understanding, results will surely follow. The instrument suffuses with light the area of operations so that no facet of the problem remains in shadow. It superilluminates every case of tuberculosis and ensures a just appreciation of its character and bearings. The penetrating ray enters the home and, linking effect with cause, brings the means of healing and the assurance of prevention to each of the dwellers.

And, throughout the area thus illuminated, watchful observers "battling with custom, prejudice, and disease," may yet snatch further triumphs and gather new facts and ideas of which, to-day, we hardly dream.

HYPOGLYCAEMIA DUE TO INSULIN IN CHILDREN.

BY

G. A. HARRISON, B.A., M.D.,*

BIOCHEMIST, HOSPITAL FOR SICK CHILDREN, GREAT ORMOND STREET, LONDON.

It is not widely known that the level of blood sugar at which symptoms of hypoglycaemia occur is decidedly lower in children under 12 years of age than in adults. There is some variation in the level at which symptoms arise following insulin administration in adult diabetics. This is observed not only when comparing one patient with another, but also in repeated observations on the same patient. In my experience the level of blood sugar when hypoglycaemic reactions occur in the adult usually lies between 60 and 80 mg. per 100 c.cm., though very occasionally values as high as 100 and even 110 or 130 mg., or as low as 50 mg., have been encountered at the time of the reaction. Anomalous results (outside the range of 60 to 80 mg.) have been experienced too often now to be explicable solely on the basis of analytical errors. It is possible that the rate of fall of the blood sugar is of more consequence than the actual level in determining the onset of symptoms, though I have not uncommonly met with rapid falls (for example, from 328 to 53 mg. per 100 c.cm.

in five hours) without resultant symptoms. Another possible explanation is that it is the concentration of sugar in the central nervous system that matters, and the sugar content of the peripheral blood may be either above or below this critical value in certain circumstances.

In eleven diabetic children, aged 1 to 11 years, treated with insulin, twenty-eight analyses out of some hundreds showed a blood sugar of less than 50 mg. per 100 c.cm. without any concomitant clinical symptoms of hypoglycaemia. The lowest value was 28 mg., and the average of the twenty-eight analyses 39 mg. In three additional records (29, 44, and 45 mg. respectively) there were slight symptoms only. On thirty occasions in the above cases values between 50 and 59 mg. were found in the absence of symptoms. Typical examples follow. All analyses were made by Maclean's method, and in duplicate.

Case 1.—Male, aged 1 year and 2 months. Blood sugar on admission 442 mg. per 100 c.cm. On the third day of insulin treatment the blood sugar was 32 mg. four hours after the morning injection of 10 units, but there were no signs of hypoglycaemic reaction. Before the injection the blood sugar was 244 mg. The diet at this time consisted of 24 oz. of milk daily. On six subsequent occasions the blood sugar was reduced to values between 28 and 49 mg. per 100 c.cm. some three to five hours after 6 units of insulin, without any signs of a reaction. On another occasion there were definite signs of hypoglycaemia. The baby was pale and slightly drowsy. He tended to fall forward, burying his head in the clothes. Blood analysis yielded the figure 44 mg. sugar per 100 c.cm. Five grams of carbohydrate (orange juice) restored the patient to normal.

The low levels of blood sugar are striking in this baby. As in adults, the level at which clinical signs occur is not a fixed point. On one day there were definite signs with a blood sugar of 44 mg., on another no signs of a reaction with a blood sugar of 32 mg.

Case 2.—Female, aged 2 years and 1 month. The daily diet contained 37 grams of carbohydrate, 31 grams of protein, and 50 grams of fat; 10 units of insulin were given daily. There were no signs of hypoglycaemic reaction on seven occasions on which the blood sugar varied from 32 to 49 mg. per 100 c.cm. There were slight but definite signs (pallor and extreme hunger) on one occasion when the blood sugar was 29 mg. per 100 c.cm.

Case 3.—Male, aged 3 years and 3 months. The blood sugar was 46 and 49 mg. per 100 c.cm. respectively on two occasions when there were no signs of a reaction.

Case 4.—Female, aged 4. Blood sugar 38 mg. per 100 c.cm. unaccompanied by clinical symptoms.

Case 5.—Female, aged 6. No symptoms when the blood sugar was 49 mg. per 100 c.cm.

Case 6.—Female, aged 9. Slight symptoms with blood sugar of 45 mg. per 100 c.cm. Symptoms relieved by 5 grams of carbohydrate (orange).

Case 7.—Male, aged 11. Fixed diet of 60 grams of carbohydrate, 70 grams of protein, and 140 grams of fat daily. He was given 8 units of insulin half an hour before breakfast, and 5 units a quarter of an hour before tea. The blood sugar results are shown in the following table. There were no signs or symptoms of hypoglycaemic reaction throughout. The boy walked to the laboratory from the ward, and "played about" in the intervals between the collections of the samples of blood.

Date.	Blood Sugar, mg. per 100 c.cm.					1½ Hours after Evening Dose.
	Hours after Morning Dose of Insulin.					
	1½	2½	3½	5	6	
1926.						
April 19th ...	63	60	55	137	123	—
April 26th ...	—	53	—	62	—	53
April 27th ...	—	43	—	61	—	55
April 28th ...	—	35	—	108	—	—
April 29th ...	Morning dose reduced to 5 units.					
May 3rd ...	—	61	—	108	—	53
May 4th ...	—	59	—	—	156	—

It will be seen that in children the blood sugar may fall some 20 to 40 mg. below the "adult hypoglycaemic level" and yet be unaccompanied by signs of a reaction. It is well recognized that the general level of blood sugar in children is lower than that of adults, and therein may lie the explanation of these observations.

There is another point of practical importance. The hypoglycaemic reaction is not nearly so well defined in children as in adults. In the younger children and infants,

* Working on diabetes with a grant from the Medical Research Council.