Utilization of Exogenous and Endogenous Ornithine by *Neurospora crassa*

R. H. DAVIS

Department of Botany, The University of Michigan, Ann Arbor, Michigan 48104

Received for publication 13 May 1968

Through the use of a mutant deficient in ornithine- δ -transaminase (OTA), it is shown that this enzyme normally has no obligate or even major biosynthetic role in *Neurospora*. The pathways of ornithine and proline synthesis proceed wholly independently of each other in OTA-less strains. It is probable that OTA functions as an enzyme of arginine catabolism. With mutants affected in OTA, ornithine transcarbamylase, and the synthesis of ornithine, it was demonstrated that exogenous and endogenous ornithine are utilized in different ways. Exogenous ornithine is destined mainly for catabolism, whereas endogenous ornithine is destined mainly for biosynthesis. It is suggested that this distinction depends upon differences in the intracellular location or origin of the two sources of ornithine.

Ornithine- δ -transaminase (OTA), despite its ability to interconvert the arginine precursor, ornithine, and the proline precursor, glutamic- γ -semialdehyde, is generally presumed to have no major role in the biosynthesis of arginine or proline in *Neurospora* (9, 17; Vogel and Vogel, Genetics 48:914, 1963). This view is reinforced by our having recently obtained "ota" mutants, which lack OTA activity, but which are nevertheless healthy and prototrophic (7). The enzyme probably functions, with arginase, as an enzyme of arginine catabolism. An OTA-deficient mutant was used in the work reported here to study the catabolism, via OTA, or the biosynthetic utilization, via ornithine transcarbamylase (OTC), of endogenous or exogenous ornithine. Evidence that endogenous and exogenous ornithine are utilized in different ways is presented.

The work reported depends upon the use of singly and multiply mutant strains. A simplified metabolic diagram, with the positions of the relevant mutations, including *ota*, is presented in Fig. 1. Previous work has shown that the *arg-8* and *arg-9* mutations, though "leaky," are blocked in the proline pathway prior to glutamic- γ -semialdehyde (13, 16, 17). They grow well on medium supplemented with proline alone, or with ornithine, citrulline, or arginine. The last three compounds are utilized via the arginase or OTA reactions, or by both (2). The *arg-5* mutant, one of several blocked in the acetylglutamate path of ornithine synthesis, grows on ornithine, citrulline, or arginine (13). The *arg-12* mutant

completely lacks OTC and grows on citrulline or arginine, but not on ornithine (8). The *arg-12^s* mutation is a unique allele of the *arg-12* locus (4, 5, 8). Despite the large deficiency of OTC (approximately 2% of wild-type activity) and the lowered affinity of the residual enzyme for ornithine (5), a strain carrying *arg-12^s* in a wild-type background is a vigorous prototroph.

The *arg-12^s* mutation has two important effects. First, it renders the *arg-8* and *arg-9* mutations independent of proline supplementation; the double mutants *arg-8*, *arg-12^s* and *arg-9*, *arg-12^s* are prototrophic (11). It is thought that ornithine not utilized as a result of the *arg-12^s* mutation (and perhaps overproduced as a regulatory response to this impairment of arginine synthesis) overflows into the proline pathway via OTA (4). The leakiness of the *arg-8* and *arg-9* single mutants may have a similar basis. An explicit test of this view is presented in this paper. The other effect of *arg-12^s* is on the utilization of ornithine from the medium. Although *arg-12^s* by itself imposes no requirement, it renders the double mutant *arg-5*, *arg-12^s* unable to use exogenous ornithine for growth (11). The behavior of *arg-5*, *arg-12^s* allowed us to select ornithine-utilizing derivatives, most of which lacked OTA (7). The latter result indicates that OTA is severely competitive with OTC in the use of exogenous ornithine. Since the *arg-12^s* single mutant is a healthy prototroph, the same degree of competition cannot prevail for endogenous

FIG. 1. Schematic diagram of arginine and proline metabolism in *Neurospora*, with the metabolic positions of certain genes. The *arg-8* and *arg-9* loci are nonallelic. The *arg-12* and *arg-12^s* mutations are allelic, but affect ornithine transcarbamylase (OTC) to different degrees. The *ota* mutation blocks ornithine transaminase (OTA).

sources of ornithine. This difference is analyzed below.

MATERIALS AND METHODS

Strains. In addition to the strains used in the work reported in the previous paper (7), the *arg-8* (allele 44207) and the *arg-12* (allele UM-3) mutations were employed. In this paper, the allele of the *ota* locus used is UM-728.

Growth. The medium used throughout was Vogel's medium N (15). It was supplemented with 1.5% sucrose as a carbon source unless otherwise noted. Stationary cultures of 10 ml were incubated at 30°C for 4 days, at which time dry weights were maximal. Logarithmic cultures were grown as described in the previous paper (7). Harvests of 10 to 40 ml, acetone-dried on 2.5-cm Whatman no. 540 filter circles, were used to monitor growth rate. The range of growth used was from 0.3 to 1.0 mg (dry weight) per ml (about 10 to 13 hr after inoculation). Material for mycelial ornithine determinations was collected on 2.5-cm membrane filters (0.45-μ pore size, GA-6; Gelman Instrument Co., Ann Arbor, Mich.), washed, and scraped cleanly from the filter with a spatula. The moist pad was placed immediately into 5 ml of cold, 5% HClO₄, and an extract was prepared.

Pool size determinations. Extracts made in HClO₄ were neutralized with 3 N KOH, and KClO₄ was allowed to precipitate in the cold overnight. After centrifugation, supernatant fractions were recovered, and the KClO₄ was washed twice with 1.0 ml of sodium citrate buffer (0.116 M Na₃ citrate, brought to pH 5.3 with 4 N HCl). The combined washes and supernatant fluid of each sample (about 6.5 ml) were adjusted to pH 5.3. These were applied to small columns of Dowex-50W (0.7 × 15 cm, 200 to 400 mesh X8) equilibrated with the buffer above. Each of the many columns used for the experiments was fitted with a 10-ml reservoir. Collection of 10-ml fractions began

as the sample was applied. The first fraction was brought to 10 ml by adding sufficient buffer to the column. Four additional fractions were taken. All proline and glutamic-γ-semialdehyde, if present, emerged in the first fraction, except for a trace occasionally found in the second. All ornithine was found in the third and fourth fractions. Arginine was retained on the column and could be displaced by 0.2 N NaOH. Ornithine was determined by the method of Chinard (3), and is expressed in terms of micromoles of ornithine per gram (dry weight) of mycelium. Ornithine determinations on samples of medium were also made after fractionation on columns.

Protein arginine. Moist harvests were made, as before, and extracted in 4 ml of cold 5% HClO₄; the precipitates, after washing, were autoclaved for 1 hr in 4 ml of 3 N HCl. This treatment released all arginine into a soluble form, measurable directly, and with the proper colorigenic value (Davis, unpublished data), by the method of van Pilsum et al. (14). The method is free from interference, but measures the small amount of arginine derived from arginyl-soluble ribonucleic acid, in addition to protein arginine.

RESULTS

The prototrophy of the *ota* mutants (7) indicates that OTA is not required in biosynthesis. The conclusion rests on the assumption that the deficiency is complete. The enzymatic assay fails to detect as much as 3% of induced levels of OTA, however, and the slow growth of transfers or of ascospores on medium containing ornithine as a sole nitrogen source (7) suggested that residual enzyme might be present. Alternatively, other enzymes might be responsible for this behavior.

A more specific physiological test of the *ota* block was performed. The *arg-8* mutant, blocked

in the proline pathway prior to glutamic- γ -semialdehyde, grew well on ornithine. Most *arg-8* strains grew poorly on minimal medium at first, but some, after a pronounced lag period, grew well. Both genetic background and environmental conditions affected this behavior, presumably by affecting the amount of endogenous ornithine available to the OTA reaction. When *ota* was inserted into a genome carrying *arg-8*, its ability to grow on minimal medium or on ornithine-supplemented medium was abolished (Table 1). The double mutant became an absolute and specific requirer of proline. These data show that the OTA block is complete in the *ota* mutant. They also show that *arg-8*, at its primary site of action, imposes a complete block in the proline pathway. An extension of these findings is that, although *arg-8*, *arg-12*^s grows well on minimal medium without a lag period (presumably through diversion of endogenous ornithine to proline synthesis), the *arg-8*, *arg-12*^s, *ota* triple mutant is completely dependent upon exogenous proline (Table 1).

Because OTA is a major catabolic route for ornithine, the *ota* mutation would be expected to increase the proportion of exogenous ornithine available for arginine synthesis. If, in addition, ornithine derived from the catabolism of arginine can enter the OTC reaction, *ota* should also increase the proportion of citrulline and arginine utilized for protein synthesis. Both expectations were borne out by comparison of *arg-5* and *arg-5*, *ota*, each grown to maximal dry weight in limiting, equimolar (0.5 mM) ornithine, citrulline, or arginine (Table 2). In the case of the double mutant, dry weights were greater on all three supplements, and there was little difference among dry weights on the three compounds. This shows that, with an intact arginine cycle and a block in ornithine catabolism, all three compounds are indeed used more efficiently. A control in regard to citrulline and arginine utiliza-

TABLE 2. Effect of *ota* on the utilization of arginine and its precursors by *arg-5* and *arg-12* strains

Strain	Dry wt (mg) per 10 ml of medium with			
	L-Ornithine (0.5 mM)	L-Citrulline (0.5 mM)	L-Arginine	
			0.5 mM	1.0 mM ^a
<i>arg-5</i>	10	19	20	28
<i>arg-5</i> , <i>ota</i>	26	25	28	42
<i>arg-12</i>	0	13	14	20
<i>arg-12</i> , <i>ota</i>	0	16	18	28

^a Used to demonstrate that 0.5 mM arginine is growth-limiting.

tion was provided by a similar comparison of *arg-12* and *arg-12*, *ota* (Table 2). The *arg-12* mutation caused a complete OTC deficiency, and therefore neither strain grew on ornithine. Since catabolically derived ornithine cannot be reutilized, however, *ota* would not be expected to enhance the growth on citrulline or arginine. Table 2 shows that there was an effect, though less than in the *arg-5* experiment. The small enhancement seen in *arg-12*, *ota* cannot be interpreted with certainty, but may reflect inhibition of the arginase reaction by accumulated ornithine. Because this has not specifically been tested, the enhancement of the growth of *arg-5*, *ota* over *arg-5* on citrulline and arginine does not definitively prove that an arginine cycle is actually operating here.

Effect of mutations on mycelial ornithine concentration. The role of OTA and OTC in the utilization of exogenous and endogenous ornithine in logarithmic cultures was studied. In these experiments, the mycelial ornithine extractable in cold HClO₄ was related to growth rate, genotype, and supplementation. In all experiments, mycelial ornithine, provisionally assumed to be intracellular, was more concentrated than ornithine in the medium. A concentration of 1 μ mole per g (dry weight) is equivalent to an intracellular concentration of about 0.4 mM (12). Table 3 summarizes determinations on a number of strains growing in minimal medium or ornithine-supplemented medium. The figures for mycelial ornithine in strains grown in minimal medium are quite characteristic, varying only 15 to 20% between experiments. Cultures grown on ornithine show characteristic and regular patterns of change.

By using wild type grown on minimal medium as a control, the effect of the *arg-12*^s and *ota* mutations could be determined. Individually, each mutation caused a slight but definite and repeatable elevation of ornithine in mycelia

TABLE 1. Dry weight (milligrams per 10 ml of medium) of strains carrying the *arg-8* mutation, after growth in various media

Genotype	Supplementation				
	None	L-Ornithine		L-Arginine (0.48 mM)	L-Proline (0.87 mM)
		0.3 mM	2.9 mM		
<i>arg-8</i> R20 ^a	0.4	11.3	30.5	7.6	27.3
<i>arg-8</i> <i>ota</i>	0	0	0	0	26.3
<i>arg-8</i> , <i>arg-12</i> ^s	28.7	—	—	—	32.0
<i>arg-8</i> , <i>arg-12</i> ^s , <i>ota</i> .	0	0	0	0	27.8

^a R20 is a particularly auxotrophic isolate of this genotype.

grown in minimal medium. More ornithine accumulated in *arg-12^s*, partially blocked in OTC, than in the OTA-less mutant, *ota*. The very small magnitude of the effect of *ota* suggests that in the wild type very little endogenous ornithine normally passes through the OTA reaction. This is consistent with the fact that *arg-8*, unimpaired in ornithine synthesis, nevertheless requires exogenous ornithine or proline for optimal growth.

In addition to its effect on mycelial ornithine content, *arg-12^s* caused endogenous ornithine to overflow into the proline pathway. This was inferred from the prototrophic phenotype of *arg-8*, *arg-12^s* (Table 1). The behavior of the double mutant was confirmed on logarithmic cultures, which grew with a wild-type doubling time. The double mutant had an elevated ornithine pool identical to the *arg-12^s* single mutant. (It is of interest that both strains have only slightly induced OTA activities on this medium. The activities were 0.008 unit per mg of protein for both *arg-12^s* and *arg-8*, *arg-12^s*, as compared with 0.006 for the wild type on minimal medium and 0.025 for the wild type on the strongest inducer, arginine.) The altered fate of endogenous ornithine in *arg-12^s* strains can also be inferred by analysis of the minimal-grown double mutant, *arg-12^s*, *ota*. By virtue of the OTA block, the ornithine diverted by *arg-12^s* accumulated to high levels in the mycelium (Table 3).

Wild type, the single mutants *arg-12^s* and *ota*, and the double mutant *arg-12^s*, *ota*, were also

grown in ornithine-containing media (Table 3). Transient or minor elevation of the ornithine pool was evident in strains having normal OTA activity. The decreasing ornithine contents of these strains in the course of growth reflects, in most cases, its exhaustion from the medium. Strains lacking OTA accumulate ornithine to much higher levels, showing that OTA plays a strong role (in wild type) in controlling the size of the ornithine pool. Ornithine accumulation in cells to levels above 100 μ moles per g is correlated with a definite, though variable, impairment of growth rate. An initial lag in growth also occurred under these conditions. The *ota* strains show a change in ornithine content prior to the rapid growth phase without appreciable consumption of ornithine from the medium. The change involves factors which have not yet been evaluated.

The arg-5, arg-12^s phenotype. The inability of *arg-5*, *arg-12^s* to grow on exogenous ornithine may merely reflect its inability to concentrate ornithine from the medium. If its failure to grow were associated with a cellular ornithine content at least as high as in a minimal-grown *arg-12^s* mycelium, however, it would show that exogenously and endogenously derived sources of ornithine are distributed differently in the cell. This would be true regardless of the small apparent induction of OTA activity in the presence of exogenous ornithine (7). The ornithine pool of *arg-5*, *arg-12^s* and of *arg-5*, *arg-12^s*, *ota* in logarithmic culture was therefore studied. The inability of *arg-5*, *arg-12^s* to grow on ornithine was very clear under these conditions, although it occasionally grew slightly after prolonged incubation.

Unlike *arg-5*, *arg-12^s*, the triple mutant *arg-5*, *arg-12^s*, *ota* grew on ornithine alone, and throughout it had an ornithine content considerably higher than that of the double mutant under comparable conditions (Table 3). This is consistent with the view that a high mycelial ornithine content is vital to the progress of the OTC reaction in *arg-5*, *arg-12^s*, *ota*. Whether a high ornithine content satisfies a concentration or a locational requirement, or both, cannot be determined with this information.

The difference between *arg-12^s*, which grows at a maximal rate in minimal medium, and *arg-5*, *arg-12^s*, which fails to utilize exogenous ornithine, was then investigated. Cultures of the two types did not differ appreciably in mycelial ornithine content (Table 3). In all experiments with *arg-5*, *arg-12^s*, limiting citrulline (0.115 mM) was used as a basic supplement so that ornithine utilization during growth could be studied. When ornithine was added to a citrulline-grown culture after

TABLE 3. *Mycelial ornithine contents and growth rates of various cultures*

Strain	Medium ^a	Doubling time	Ornithine ^b
		hr	μ moles/g
Wild type 74A . . .	MIN	2.5	30 $\pm$ 5
	ORN	2.5	80 to 59
<i>arg-12^s</i>	MIN	2.5	55 $\pm$ 3
	ORN	2.5	54 to 48
<i>ota</i>	MIN	2.5	40 $\pm$ 4
	ORN	3.5	800 to 240
<i>arg-12^s</i> , <i>ota</i>	MIN	4.0	100 $\pm$ 5
	ORN	3.0	310 to 260
<i>arg-8</i> , <i>arg-12^s</i>	MIN	2.5	52 $\pm$ 2
	CIT + ORN ^c	3.5	63 to 26
<i>arg-5</i> , <i>arg-12^s</i>	ORN	4.5	206 to 109

^a MIN = minimal; ORN = 0.595 mM L-ornithine; CIT = 0.115 mM L-citrulline.

^b Figures for minimal cultures given with standard deviation. Figures for ornithine-grown cultures are the first and last determinations in the course of logarithmic cultures. Ornithine was present in the medium in all cultures when the first sample was taken. At the last determination, ornithine had disappeared from the medium of cultures of strains not carrying the *ota* mutation and persisted in *ota* strains.

^c See text and Fig. 2 for related experiments.

growth had stopped, growth did not resume (Fig. 2A). If the medium was supplemented with 0.595 mM (Fig. 2A) or 2.97 mM (Fig. 2B) L-ornithine from the time of inoculation, all ornithine was consumed in the course of growth in the first case, and 24% was consumed by the stationary phase in the second. The plateau dry weights were not appreciably different in the two cases, but both were somewhat higher than in cultures provided only with citrulline. The considerable catabolism of added ornithine, when seen in light of the relatively small enhancement of growth, indicates that ornithine enters the cell, but is destined mainly for a catabolic fate in this strain. This conclusion is strengthened by other observations, reported below.

The enhancement of growth observed in cultures grown in citrulline plus ornithine, as noted above, is not related to the concentration of ornithine. This suggested that the effect of ornithine was not to provide arginine, but to improve the efficiency of citrulline utilization. To test

this hypothesis circumstantially, protein arginine from cultures grown in 0.115 mM citrulline or in 0.115 mM citrulline plus 2.97 mM ornithine was measured after growth had ceased. In the absence of ornithine, only 84% as much arginine was recovered as had been added to the medium as citrulline. In the presence of ornithine, 102% as much protein arginine (0.118 μ mole per ml of culture) was recovered as had been added as citrulline, a value which is constant in the final phase of growth (Fig. 2B). The data suggest that ornithine may inhibit the catabolism of arginine (derived from citrulline) in the arginase reaction. The experiment therefore conforms to other observations that exogenous ornithine cannot be transformed to arginine in *arg-5*, *arg-12^s*, and suggests that exogenous ornithine is destined exclusively for catabolism in this strain.

When growth of *arg-5*, *arg-12^s* ceases in the presence of ornithine, the ornithine content is 50 to 60 μ moles per g, a figure which falls slowly to about 25 μ moles per g (Table 3 and Fig. 2B).

FIG. 2. (A) Growth of *arg-5*, *arg-12^s* on various media. Abscissa: time after inoculation. Symbols: ●, 0.95 mM L-arginine; ○, 0.115 mM L-citrulline plus 0.595 mM L-ornithine from time of inoculation; △, 0.115 mM L-citrulline; ▲, 0.115 mM L-citrulline, with 0.595 mM L-ornithine added at arrow. (B) Growth of *arg-5*, *arg-12^s* on 0.115 mM L-citrulline plus 2.97 mM L-ornithine (bottom). The accumulation of protein arginine per milliliter of culture (middle) and the mycelial ornithine per gram of dry weight (top) are also shown.

The slow decrease in mycelial ornithine may reflect the inability of the cells to maintain uptake during the stationary phase, and in fact ornithine consumption from the medium ceases. The mycelial ornithine concentration at the end of the growth period, however, is indistinguishable from the corresponding figure for *arg-12^s* mycelia growing vigorously in minimal medium. This indicates that the origin of the ornithine, rather than its mycelial concentration alone, determines whether the OTC reaction will proceed in strains carrying *arg-12^s*.

DISCUSSION

The prototrophy of the *ota* strain, together with the interaction of the *ota* mutation with the proline mutation, *arg-8*, proves that OTA has no obligate biosynthetic role, and that *arg-8* blocks completely the normal route of proline synthesis. The use of the *ota* strain should make the selection of additional proline mutants very efficient. The other known mutant of the proline pathway, *arg-9*, is similar to *arg-8* in its nutrition and its interaction with *ota*, though its genetic lesion may not be complete. The *arg-9* mutation is closely linked to, but nonallelic with *ota* (Davis, *preliminary results*).

The OTA deficiency, by blocking catabolism, allows exogenous ornithine to be transformed more efficiently to arginine. The data suggest that OTA is mainly an enzyme of arginine catabolism, of significance in the recovery of δ -amino nitrogen from ornithine. In addition, since ornithine can be used (though poorly) as a carbon source by wild-type *Neurospora*, it is probable that glutamic- γ -semialdehyde can be transformed to glutamic acid.

The *arg-5*, *arg-12^s* strain cannot grow on exogenous ornithine. The fact that ornithine is consumed by growing mycelia of this genotype shows that entry into the cell is not blocked, although the data suggest that *arg-12^s* may diminish the maximal ornithine content of mycelia. The data indicate that access of exogenous ornithine to the intracellular site of the OTC reaction is poor. Only when the cellular ornithine concentration is abnormally high, as in the OTA-deficient *arg-5*, *arg-12^s*, *ota*, will growth on ornithine occur. Because Abelson and Vogel (1) detected preferential use of ornithine via OTA in the wild type, the poor access of exogenous ornithine to OTC in *arg-5*, *arg-12^s* cannot be due solely to the *arg-12^s* mutation. Rather, the phenomenon is a general one for *Neurospora*.

A counterpart to poor access of exogenous ornithine to OTC is the ability of *arg-12^s* to grow without any nutritional supplementation, that is, on endogenous ornithine. While much endoge-

nous ornithine is catabolized by this strain, it is possible that endogenous ornithine originates at a surface or in a compartment (6) such that it is also accessible to OTC. The idea of a spatial distinction is supported by the observation that *arg-12^s* mycelia contain no more ornithine than those of *arg-5*, *arg-12^s* which have ceased growth in the presence of exogenous ornithine. The effect of the "channeling" mechanism, if it also prevails in the wild type, is to distribute exogenous ornithine mainly to OTA and endogenous ornithine mainly to OTC. Until it is proved that exogenously derived ornithine in the double mutant is actually within the cell membrane, conclusions regarding spatial parameters are tentative. Further work on this aspect of the problem is continuing with the use of tracers, and with tests for different localizations of OTC and OTA. Up to now, both enzymes have been considered "soluble" in *Neurospora* (5, 18). A suggestion that OTC is in fact localized comes from a recent report by Kinsey and Wagner (10), who showed de novo arginine synthesis in an isolated mitochondrial fraction of *Neurospora*.

ACKNOWLEDGMENTS

I thank J. D. Zimmerman and M. Lawless for technical assistance, J. Mora for discussion, and L. G. Williams and M. L. Sargent for critical reading of the manuscript.

This investigation was supported by grants GB-2342 and GB-6414 from the National Science Foundation.

LITERATURE CITED

1. Abelson, P. H., and H. J. Vogel. 1955. Amino acid biosynthesis in *Torulopsis utilis* and *Neurospora crassa*. *J. Biol. Chem.* **213**:355-364.
2. Castañeda, M., J. Martuscelli, and J. Mora. 1967. The catabolism of L-arginine by *Neurospora crassa*. *Biochim. Biophys. Acta* **141**:276-286.
3. Chinard, F. P. 1952. Photometric estimation of proline and ornithine. *J. Biol. Chem.* **199**:91-95.
4. Davis, R. H. 1962. Consequences of a suppressor gene effective with pyrimidine and proline mutants of *Neurospora*. *Genetics* **47**:351-360.
5. Davis, R. H. 1967. A mutant form of ornithine transcarbamylase found in a strain of *Neurospora* carrying a pyrimidine-proline suppressor gene. *Arch. Biochem. Biophys.* **97**:185-191.
6. Davis, R. H. 1967. Channeling in *Neurospora* metabolism, p. 303-322. In H. J. Vogel, J. O. Lampen, and V. Bryson (ed.), *Organizational biosynthesis*. Academic Press, Inc., New York.
7. Davis, R. H., and J. Mora. 1968. Mutants of *Neurospora crassa* deficient in ornithine- δ -transaminase. *J. Bacteriol.* **96**:383-388.
8. Davis, R. H., and W. M. Thwaites. 1963. Structural gene for ornithine transcarbamylase in *Neurospora*. *Genetics* **48**:1551-1558.
9. DeDeken, R. H. 1963. Biosynthèse de l'arginine

- chez la levure. I. Le sort de la $N\alpha$ -acétylornithine. *Biochim. Biophys. Acta* **78**:606-616.
10. Kinsey, J. A., and R. P. Wagner. 1966. Amino acid production by a mitochondrial fraction of *Neurospora crassa*. *Proc. Natl. Acad. Sci. U.S.* **55**:404-409.
 11. Mitchell, M. B., and H. K. Mitchell. 1952. Observations on the behavior of suppressors in *Neurospora*. *Proc. Natl. Acad. Sci. U.S.* **38**: 205-214.
 12. Slayman, C. W., and E. L. Tatum. 1964. Potassium transport in *Neurospora* I. Intracellular sodium and potassium concentrations, and cation requirements for growth. *Biochim. Biophys. Acta* **88**:578-592.
 13. Srb, A. M., J. R. S. Fincham, and D. M. Bonner. 1950. Evidence from gene mutations in *Neurospora* for close metabolic relationships among ornithine, proline, and α -amino- δ -hydroxyvaleric acid. *Am. J. Botany* **37**:533-538.
 14. Van Pilsum, J. F., R. P. Martin, E. Kito, and J. Hess. 1956. Determination of creatine, creatinine, arginine, guanidoacetic acid, guanidine, and methyl-guanidine in biological fluids. *J. Biol. Chem.* **222**:225-236.
 15. Vogel, H. J. 1964. Distribution of lysine pathways among fungi: evolutionary implications. *Am. Naturalist* **98**:435-446.
 16. Vogel H. J., and D. M. Bonner. 1954. On the glutamate-proline-ornithine inter-relation in *Neurospora crassa*. *Proc. Natl. Acad. Sci. U.S.* **40**:688-694.
 17. Vogel, R. H., and M. J. Kopac. 1959. Glutamic- γ -semialdehyde in arginine and proline synthesis of *Neurospora*: a mutant-tracer analysis. *Biochim. Biophys. Acta* **36**:505-510.
 18. Vogel, R. H., and M. J. Kopac. 1960. Some properties of ornithine- δ -transaminase from *Neurospora*. *Biochim. Biophys. Acta* **37**:539-540.