described above. I submit the following proposals for further investigation:

1. The preparation of suitable media of human and bovine material containing unheated blood serum.
2. The evolving of a vigorous strain of tubercle bacilli which will grow on human and bovine media which are not sterilized by heat.
3. The elimination of autolysis in such media and investigation into the possible action of autolytic enzymes on the tubercle bacillus.
4. The preparation of a tuberculin by the method described above.
5. Preliminary investigations to be carried out with bovine tubercle bacilli on media of bovine material.

In conclusion I have to express my thanks to Professor Tulloch for his valuable advice and assistance, and to Dr. Cumming for the time and labour he has given to the preliminary investigation.

A CASE OF TRANSVERSE ECTOPIA OF THE TESTIS.

BY

A. GRAY BANKS, M.D. GLASG., F.R.C.S. ED.,
SURGEON, EAST SUFFOLK AND IPSWICH HOSPITAL.

The interest of the following case is increased by the note Sir Arthur Keith has been good enough to append.

A man, aged 28, married three years, no children, was admitted to the East Suffolk and Ipswich Hospital in December, 1925, with a painful swelling in the left groin. Eighteen years earlier an operation for left inguinal hernia had been performed in the same hospital, but details are unobtainable. He had been aware of the lump ever since the operation and had noticed it becoming gradually larger, more superficial, and moving downwards. Nine months ago he had an attack of acute pain after lifting a heavy weight, and the lump became tender and swollen, and had remained more or less tender since. Two further attacks of pain occurred, one three months, and the other three weeks ago. On each occasion micturition was frequent for about two days; between the attacks there were no urinary symptoms. Sexual power developed when he was about 14, and has been normal since.

He was a spare, vigorous, active-looking man of good intelligence; the secondary sexual characters were well developed. A linear scar of old standing was present over the left inguinal canal, and lying at the external ring just below the pubic spine was a firm ovoid swelling about the size of a walnut, slightly movable and tender on pressure. The scrotum was small and contained only one testis (left), from which an easily palpable cord could be traced upwards. The right side of the scrotum was empty, and no opening could be felt at the usual site of the external ring (the entire absence of any weakness here was confirmed by intra-abdominal palpation during the operation).

Operation.

An oblique incision was made over the inguinal canal and the swelling identified as a testis (although at first sight the appearance suggested an ovary). It had a well marked tunica vaginalis. From the upper border of the testis a fold of fascia and peritoneum, forming a definite mesorchium and having a cord-like structure with fimbriated end along its free edge, entered the peritoneal cavity through the ring and crossed obliquely over the bladder as a lax membrane to the opposite pelvic brim, where it was attached about the point of bifurcation of the iliac artery.

The left testis was drawn out of the scrotum and examined. It was rather undersized, devoid of a tunica vaginalis, and on incising the tunica albuginea a normal testicular appearance was seen; the vas deferens of this testis was adherent to, and partially embedded in, a firm structure about a fingerbreadth thick, which ran upwards from the testis, through the inguinal opening, and down extraperitoneally along the left side of the bladder towards the base. In the intra-abdominal part of its course both vasa deferentia were embedded in it, the left more intimately, so that they could be dissected out only with great difficulty. When opened it was found to have a cavity admitting a fair-sized probe, the wall about a quarter of an inch thick, rugose, and apparently lined with endothelium.

The distal part was excised, about one to one and a half inches being left to avoid injury to the associated left vas deferens. The ectopic testis was removed with the accompanying fold, the cord having been freed from the above mentioned structure as far as possible, and the oblique muscles and skin sutured. The wound healed by first intention and the patient was perfectly comfortable on discharge two weeks later. A cystoscopic examination was refused.

Serial sections of the ectopic testis, kindly made by Sir Arthur Keith, showed only testicular tubules—no mixture of sex. "No spermatozoa are to be found and none in process of formation. The interstitial tissue is normal in amount (A. K.)."

The excessive rarity of the condition is shown by the fact that apparently only four cases of transverse ectopia are recorded.

1. Lenhossék in 1845 first described the condition in a man of 35 with two testes in the left half of the scrotum, the right lying a little higher than the left, in front and medianward. The vasa deferentia were closely united but distinct, the tubes separating only near the prostate.

2. Max Jordan in 1885 reported a case in a child of 8. There were two testes in one patent vaginal process with one vas which divided higher up. The testes were in the left half of the scrotum and the united vasa had coalesced for the greater part of their length.

3. Berg in 1901, operating on a congenital left inguinal hernia in a boy of 13, found two testes—one normal in a well developed scrotum, and the other, almond size, at the left ring. Both cords were intimately connected with the hernial sac; the ectopic cord had a distinct mesentery and passed up across the space of Retzius to the right inguinal region, whence it descended to the pelvis. The right ring was palpated and found closed.

4. Halstead (1907) records a case with two small testes and a large left inguinal hernia; the epididymes were fused, but the vasa deferentia were distinct up to the region of the internal ring where they were united.

NOTE BY SIR A. KEITH.

The abnormal structure found passing upwards from the testis, and descending to the base of the bladder, can only be the result of fusion and persistence of the Müllerian ducts, forming a uterus masculinus of unusual size and persistence. If this were drawn down into the hernial sac, as is quite possible, it would bring with it the attached undescended right testis. The vasa deferentia should be embedded in the "structure," exactly as found at operation. This, I think, is the true interpretation, not a double or dichotomous condition of the left testis.

It is to be supposed, therefore, that in this patient there had been a persistence of the stage reached and usually passed through in the third month of development.

REFERENCES.

- ¹ Lenhossék: *Anatomischer Anzeiger*, 1886. ² Jordan: *Deut. med. Woch.*, 1885. ³ Berg: *Annals of Surgery*, August, 1904. ⁴ Halstead: *Surg., Gynecol. and Obstet.*, 1907.

NEPHRO-URETERAL ANASTOMOSIS AFTER COMPLETE AVULSION OF THE URETER.

BY

R. CAMPBELL BEGG, M.C., M.D., F.R.C.S. ED.,
M.Sc.N.Z.,

GENITO-URINARY SURGEON, WELLINGTON HOSPITAL, N.Z.

A MAN, aged 42, was referred to me on January 19th, 1926, complaining of constant pain in the right loin. It had existed with varying intensity for many years, but he had had nothing in the nature of an attack of renal colic. Investigation showed that there was a round calculus in the right renal pelvis, and that the urine was infected with *B. coli* and was delivering pus. The kidney function, however, by dye tests showed that there was only slight loss of function on the right side as compared with the left. In the latter, indigo carmine, after intravenous injection, appeared in strong concentration in five minutes, while the dye did not appear till six and a half minutes from the right side, but the concentration was good. The calculus appeared to be about 1 cm. in diameter, so pyelo-lithotomy was advised.

Operation.

On February 12th the kidney was exposed with some difficulty owing to its high position; a portion of the twelfth rib had to be resected. The organ seemed healthy except for one or two points of subcapsular scarring. The pelvis was mostly intrarenal and the calculus was lying just outside the hilum, while immediately below it a ureter of normal calibre passed downwards. The calculus was removed by a vertical incision. The walls of the pelvis seemed to be very thin and there was a ring of ulceration where the stone had been lying. An attempt to stitch the walls of the incision with the finest round needle and catgut resulted in every stitch pulling through. My assistant had to exercise a certain amount of tension on the kidney to keep the field in view, and suddenly the ureter came completely away from the pelvis at the line of the circular ulcer. The line of separation was very close to the kidney, and the little pelvic tissue available for anastomosis was so friable that an end-to-end junction was impossible; after some attempts at this there remained no extrarenal pelvis, but merely a few bits of ragged tissue around the orifice of the renal sinus. Nephrectomy was considered, but owing to the comparatively good renal function present seemed undesirable. The ureter was not dilated and had good muscular walls. A No. 6F catheter