the abdomen was noticed to be rather distended, and an area of dullness was found in the right flank continuous with the liver dullness. No tumour could be palpated. A diagnosis of haemorrhage into the peritoneum was made, possibly from some abdominal tumour. At 3.30 p.m. the child died.

Post-mortem Examination.—This was performed about 9 p.m. on the same day. The thoracic viscera appeared normal, but there was very little blood in the heart or great vessels. Through the right dome of the diaphragm a dark mass could be seen. On opening the abdomen about 6 oz. of free blood and blood clot were found, mostly in the right flank. The right lobe of the liver was dark from blood effused under the capsule, which was stripped up over an area extending above into the diaphragmatic surface of the liver, and in front to within about three-quarters of an inch of the falciform ligament. From a minute perforation on the anterior aspect of the capsule blood was still oozing. On cutting across this area a deep sulcus was seen on the anterior surface, running from above downwards and slightly inwards, about 2½ inches in length. To the right of this a dark area, roughly ovoid, sharply demarcated from the liver, was made out. This area was situated in the right lobe, and extended into the cystal surface in front, and slightly on to the abdominal aspect beneath. It measured about 1½ inches across and about 2½ inches from above downwards. The liver weighed 243.06 grams. The remaining organs appeared normal.

Microscopical Examination of the Liver.—There was congestion and marked fatty change. An area of haemorrhage showed intermingled strands of fibrous tissue and a boundary of compressed haemorrhagic and fibroid liver tissue. This boundary showed in parts a leucocytic infiltration, and throughout the haemorrhagic area the leucocytes appeared to be increased, and were for the most part mononuclear in type. There was some early organization of the blood clot which would correspond with the period—five days—between the birth and death of the infant. The appearances suggested that the haemorrhage occurred at birth from trauma.

There can be very little doubt that the fissure found in the liver was a laceration which probably occurred during birth, although the fact of labour having been easy and normal makes it rather difficult to understand how it occurred. Holt states that small haemorrhages are not uncommon upon the surface of any of the viscera covered by peritoneum, and that intraperitoneal haemorrhages are rare, but may be very extensive, amounting to six or eight ounces, as in the present instance. Small surface haemorrhages of the liver, he states, are not infrequent. Occasionally one of considerable size occurs, separating the peritoneal covering and forming a tumour, generally upon the superior surface. Such lacerations may be produced during labour, and a slow accumulation of blood may take place beneath the capsule, death resulting from rupture into the peritoneal cavity.

RECOVERY FROM TUBAL DEAFNESS CAUSED BY A SARCOMA.

BY

SAMUEL LODGE, O.B.E., M.D.,

AND

WILLIAM OLIVER LODGE, M.D., F.R.C.S.E.,

BRADFORD.

(From the Aural Department, Royal Halifax Infirmary.)

THE following case is one of removal from the soft palate of a sarcoma which resembled clinically the type of endothelioma found in this situation.¹ As a result of relief of pressure on the Eustachian tube the patient recovered from unilateral deafness of eight years' standing.

An active man engaged in responsible affairs became deaf in the right ear at the age of 35 and noticed a swelling at the back of the mouth three years later. At the age of 40 he could only hear ordinary conversation close to the right ear. The swelling grew larger internally and began to project outwards behind the angle of the jaw. The voice had a nasal intonation, but there was no interference with breathing or swallowing.

The smooth yielding tumour occupied the right side of the soft palate and nasopharynx, the tonsil being displaced medially, and was slightly movable on bimanual palpation. The drumhead of the right ear was indrawn and with the C 256 tuning-fork bone conduction was relatively increased. With the stem of the vibrating fork on the vertex the sound was heard loudly in the right ear; by air conduction C 1024 was heard well, C 128 not at all. The left ear was normal. Systemic examination, including a blood count, revealed nothing abnormal.

Operation.

General anaesthesia was induced with chloroform and continued through a laryngotomy tube. The tumour, which extended deeply into the pterygo-maxillary and submaxillary regions, and was

surrounded with a false capsule, was dissected away. The walls of the cavity were coagulated by diathermy. A week later a tube of radium emanation was placed in the depth of the cavity and another towards the root of the tongue; these were secured in position for twenty-four hours by packing.

On section the tumour proved to be a spindle-celled sarcoma; portions of muscular tissue removed were seen to be invaded.

After-Treatment.

As soon as the sloughs had separated the patient was referred to Dr. William Mitchell for x-ray treatment, and five exposures were given. Three months later the drum was less retracted, and on inspection with Hay's pharyngoscope the Eustachian orifice appeared normal. After one or two inflations with the catheter the hearing for the watch and voice was completely restored. Rinne's test was positive, though air conduction was still a few seconds less in excess of air conduction than on the left side. Four years after the operation there was no sign of recurrence, and the hearing was normal.

Appearance of the Tumour.

What was the pathological condition of the right ear from which the patient recovered so promptly when the patency of the Eustachian tube was restored? In three cases of obliteration of the Eustachian tube by malignant tumours of the nasopharynx Brock found serous exudation into the tympanic cavity and its adnexa, attributable to hydrops *ex vacuo* rather than to inflammation. Karl Beck of Heidelberg, experimenting with dogs, closed the tube with wooden plugs, or, as was found more effective, with the thermo-cautery. After killing the animals, microscopic examination revealed hypertrophy and cellular infiltration of the mucous membrane, some bony thickening, and a sterile exudate containing polymorphs and lymphocytes. According to Habermann² this condition differs from chronic catarrh of the middle ear, in which proliferative changes in the tympanic mucosa are identical with those of hypertrophic rhinitis. In our case, in which closure of the Eustachian tube was probably gradual and possibly intermittent, there were no subjective symptoms of fluid in the tympanic cavity, nor could any fluid exudate be seen through the drum.

Only by the finest discrimination between the forms of chronic deafness can we hope to judge beforehand whether operations on the nose and throat will restore hearing.

REFERENCES.

- ¹ G. B. New: *Collected Papers of the Mayo Clinic*, vol. xii, 1920, p. 841. ² J. Habermann: *Archiv f. Ohren-, Nasen-, und Kehlkopfheilkunde*, 109 Band, 2/3 Heft, Seite 105, April, 1922.

CONGENITAL OCCLUSION OF DUODENUM.

BY

ANDREW V. MAGEE, M.B., CH.B.GLAS.,

SENIOR RESIDENT ASSISTANT MEDICAL OFFICER, BOOTH HALL INFIRMARY FOR CHILDREN, MANCHESTER.

DR. H. C. CAMERON, in the Lumleian Lectures, delivered last year,¹ discussed in the first lecture vomiting in infancy due to obstruction from developmental defect in the upper alimentary tract. The following case is reported because, as Cameron says, "the diagnosis, supported by a characteristic radiographic appearance, is often not difficult, and because, sometimes at least, we may hope to save life by means of a timely operation following upon early diagnosis." The condition, however, appears to be very rare. Only one case² has been reported in this infirmary, with about 3,500 admissions annually, in nine years.

A male infant, aged 5 days, was admitted to Booth Hall Infirmary, Manchester, on January 15th, 1926. There was a history of vomiting everything taken by the mouth since birth. The patient was the weaker and smaller of uniovular twins and weighed 4½ lb. at birth. The pregnancy and labour were normal and the parents and their children were healthy.

On examination the heart and lungs were found normal. The tongue and pharynx were red and congested. The abdomen was not distended, but some fullness was seen and felt in the epigastric region. No intestinal peristaltic waves were seen; there was no jaundice. The anus and rectum were normal. X-ray examination of the stomach was not done.