It has been possible to keep in touch with practically all those treated up to date, as the majority of them are so afraid of becoming chronic invalids again that they readily carry out instructions and report themselves regularly.

I wish it to be clearly understood that what I have here written is but a brief summary of beneficial effects so far obtained in asthma by means of radiation applied in a manner not usually adopted. The number of cases so far treated (21) is too small and the time too short for any statement as to the permanency or true efficiency of the method to be made. It is hoped, however, that those who have access to a larger number of patients will give it a trial, so that a true valuation may be reached.

So far radiation applied as described has proved a valuable palliative agent; time will indicate its true curative value. Radiation attacks this distressing complaint from an entirely new angle, and while at present it is not based on any theory more light may be thrown on the cause if we eventually find out what influence small radiation doses over a large body area have on general metabolism.

Radiation, of course, has been tried before in asthma, and the results reported have on the whole been disappointing. Ramirez and Cole come to the conclusion that radiation offers no particular advantage over other methods, but that definite effects noted in some of their cases would indicate that a modification of technique might yield a more encouraging result. I believe the fault in the methods so far tried lies in the limited radiation field—that is, the lungs alone are looked upon as the site of the trouble. This, combined with overdosage, accounts for the poor results.

This preliminary note may encourage others to give the method a trial, but careful attention to essential details is necessary.

LITERATURE.

- Ramirez and Cole: Roentgen-Ray Therapy in Bronchial Asthma, *Amer. Journ. of Roentgenology*, October, 1925.
 Immelmann: Report to Balneological Congress, Berlin, 1907.
 Pohlmann: *Munch. med. Woch.*, 1925.
 Grodel: *Die Röntgen Behandlung Inner. Krankheiten*, 1924.
 Gerber: X-Ray Treatment of Bronchial Asthma, *Journ. Amer. Med. Assoc.*, October 3rd, 1925.

HAEMATOMA IN THE ABDOMINAL WALL SIMULATING PELVIC NEW GROWTH.*

BY

W. E. FOTHERGILL, M.D.,

PROFESSOR OF CLINICAL OBSTETRICS AND GYNAECOLOGY, VICTORIA
UNIVERSITY, AND SURGEON TO ST. MARY'S HOSPITALS FOR
WOMEN, MANCHESTER.

HAEMATOMATA in the lower part of the abdominal wall sometimes present such close resemblances to tumours of pelvic origin that very nice diagnostic problems may arise. Cuthbertson¹ has recently collected forty-one cases in women of which no fewer than twelve were diagnosed as ovarian cysts and ectopic gestation. In only three of the forty-one cases was the haematoma recognized before operation, while in two others it was suspected. Thus the recognition of this condition is clearly a matter of some difficulty, and one which has points of special interest for those who deal with the diseases of women. This is illustrated by a recent case.

A lady, aged 57, was quite well until she had a day of severe abdominal pain. She sent for Dr. Bennett of Hyde, who found a considerable swelling rising above the pubes on the right side. The pain subsided under treatment; but, two days later, diffuse black bruises appeared upon the surface of the lower abdomen. When I saw her during the following week there was a discoloured area, measuring about eight inches by four, containing many patches of subcutaneous haemorrhage. There was a rounded swelling above the pubes, mainly to the right of the mid-line. It resembled a four and a half months pregnant uterus in shape and size, but was more like a fibroid in consistency. The cervix was pushed forward by a cystic swelling, which could be felt in the pouch of Douglas. Thus the physical signs suggested the presence of a fibroid above and an ovarian cyst below. But when the patient was asked to sit up, and so brought the recti into contraction, the swelling above the pubes could still be felt, and became quite fixed. This showed that it was really in the abdominal wall. The sudden onset of pain accompanied by the appearance of the lump and followed by the extravasation of

blood under the skin indicated that rupture of some fibres of the right rectus had occurred. The diagnosis, therefore, was haematoma of the abdominal wall, together with ovarian cyst in the pouch of Douglas.

The abdomen was opened through blood-stained tissues to the left of the swelling, the linea alba being displaced towards that side. A simple cyst of the left ovary was then removed. The uterus and right tube and ovary were normal. The appendix was removed, as it was involved with a portion of omentum, which was adherent near the internal abdominal ring. The haematoma lay within the right rectus, with a thick layer of muscle over it. The mass of blood clot projected into the abdominal cavity, with a thin layer of muscle and the peritoneum under it. The clot was turned out through a cut in the edge of the muscle. The wound was then closed in the usual way, a strip of gauze being left in the cavity of the haematoma for one day. Healing occurred quickly.

I once saw a case in which someone had diagnosed ovarian tumour in the absence of the patient's usual medical adviser, Dr. Robertson, then of Rawtenstall.

The patient was a stout, healthy woman, past the menopause, and she told us that the pain and swelling dated from a fall over a clothes-basket, which hurt her very much, but did not prevent her from finishing her washing. The right lower abdomen was quite dull and solid, as if the right iliac fossa was filled by a mass which rose above the umbilicus and crossed the middle line. The mass could still be felt, and became fixed when the patient sat up, and this sign, with the history of the fall, gave the diagnosis of haematoma. Expectant treatment was employed.

Later there was some constitutional disturbance, and the mass became softer. When I saw the patient again the physical signs were those of a large cyst adherent to the abdominal wall, and it was clear that the blood-clot had become infected and liquefied. An incision through the right rectus liberated a great quantity of chocolate-coloured pus mixed with blood-clot which had been distending a cavity walled in front by the rectus and separated from the abdominal cavity by peritoneum below and by peritoneum and the sheath of the rectus above.

I subsequently saw a similar case, and, in the light of previous experience, recognized it as an infected and liquefied haematoma, and arranged for the patient's admission to hospital. When she came in the lapse of time had introduced a new and disconcerting feature which rather shook the diagnosis, for the swelling, previously cystic, was now tympanitic. I remember the horror of some of those present when I incised this resonant area, and their relief when a blast of foul gas was followed by a gush of chocolate-pudding pus.

I once watched an operation after a very sporting diagnosis had been made—namely, myoma of the round ligament. The outcome was that the surgeon struck pus before he had cut through the abdominal wall. No doubt when he examined the patient the mass felt solid and closely resembled a fibroid. Infection and liquefaction must have occurred in the time that elapsed between the making of the diagnosis and the operation.

Infection does not always occur. Dr. J. W. Stenhouse had a patient, a healthy elderly lady, with a firm fixed swelling above Poupart's ligament which appeared suddenly with pain. We concluded that this was a haematoma, and it disappeared entirely in the course of a few weeks. When I was house-surgeon to the late Sir A. R. Simpson two students brought their old landlady to see him in the Buchanan ward of the Edinburgh Royal Infirmary. She thought she had strained herself while leaning out to clean a window. There was a lateral solid mass which seemed to rise out of the left iliac fossa and to be attached to the iliac bone. Simpson did not consider it suitable for operative treatment. He told us to strap her well up with strips of belladonna plaster. This gave her much comfort, and after a time she came back saying the plaster was too loose. It was renewed, and this happened two or three times. In the end the lump disappeared. These two cases stand out in my memory as the only disappearances of considerable lumps other than inflammatory deposits which I have personally observed. I believe they were both haematomas, though this was not proved by incision in either case.

As to the general aspects of the condition but little has been written; it is hardly mentioned in some textbooks and is treated briefly in others. Numerous cases have been recorded in the medical journals, most of which would seem to be the results of strains and accidents in men of active habits—cavalry soldiers, acrobats, and others whose occupation or sport demands violent effort. In women, on the other hand, the lesion occurs, apart from obvious

* Read before the North of England Obstetrical and Gynaecological Society, April 16th, 1926.

trauma, in those of middle or advanced age. It is sometimes associated with coughing or vomiting. In the cases I have mentioned the patients were all past the menopause. There was a history of a fall in one case and of a strain in another. Out of Cuthbertson's¹ forty-one cases, thirteen were over 50 years of age; seven were between 40 and 50; six were under 40. Twelve of the patients were pregnant or had recently been delivered.

Haematomata which simulate pelvic growths are below the umbilicus, and are thus due to the rupture of muscular branches of the deep epigastric artery. Being impressed by the marked lateral extension of the swelling in these cases I consulted Professor Stopford on the point. He says that if the haemorrhage occurs below the fold of Douglas, owing to the formation of the sheath of the rectus at that level, there is nothing to prevent the lateral extravasation of blood deep to the aponeurosis of the oblique muscles. This explains the relatively limited character of haematomata higher up in the rectus muscles. The extravasation of blood cannot cross the linea alba, but the swelling can extend beyond the middle line by pushing the linea alba over toward the sound side of the body.

The most interesting thing about these cases is the diagnosis. The patient complains of pain and the medical man finds the swelling. The trouble is that he seldom knows how long the swelling has been present; for patients frequently have masses of various kinds in the pelvis or abdomen whose presence is not suspected until they complain of pain and submit to examination. The main point is the recognition that these swellings are part and parcel of the abdominal wall. This is generally made by noting that they can still be felt when the recti are in action, and that they become fixed as the muscles contract. Another point is that, when a mass in the abdominal wall is not too large, a resonant note may be obtained right through it by deep percussion. But these signs are not infallible, for a mass within the abdomen that has become adherent to the wall gives the same signs as one which is a part of the wall. Even physical signs can tell lies, and must be checked.

This point decided, the next question is, What other swellings may be found in the wall of the abdomen? My first employment after qualification was to do duty for the house-surgeon of a hospital in a country town. He called my attention to a girl with a so-called ovarian cyst who was being prepared for operation next morning. The girl told us that the swelling was there sometimes, but not always. It was cystic, in the mid-line, and reached just to the umbilicus. It was still felt, and was fixed when the recti were in contraction. Being very young then, and full of book learning, I suggested that it might be a cyst of the urachus communicating with the bladder. Strange to say, it was, and we massaged the contents into the bladder and drew it off with a catheter.

A girl, aged 13, was brought to the Northern Hospital, Manchester, with a swelling above the pubes, which was part of the abdominal wall. The mother had been told that it was a pregnant uterus, and, when assured that this was not the case, she took the child away. But, later, the patient reached the Children's Hospital at Pendlebury, where the late Mr. Joseph Collier opened an abscess and found in the centre of it a rusty needle.

The commonest swellings in the abdominal wall are new growths—namely, those masses of spindle-celled tissue called desmoids. They are generally single, but I have seen three in the recti of one patient. Slow-growing and painless, they are easily recognized. They were frequently excised formerly, but they yield to treatment by irradiation in most cases.

Another new growth in the abdominal wall was seen in a patient from whom I had removed an ovarian cystadenoma six years previously. She came back saying she had another tumour upon which her corsets pressed uncomfortably when she sat down. It was a solid encapsuled mass under the scar of the former operation, and, on section, its structure was that of the solid portions of a cystadenoma.

Thus, before completing the diagnosis of a haematoma, it is necessary to exclude new growths, inflammatory masses, and developmental errors; for they can all produce swellings

in the abdominal wall which simulate tumours of pelvic origin. As to treatment, a small haematoma in this position may doubtless be left alone. But should we wait for the larger ones to become infected before opening and emptying them? I think not. If the mass of blood-clot is of considerable size the patient will be sound and well again much sooner under active surgical than under an expectant treatment.

REFERENCE.

¹ *Journ. Amer. Med. Assoc.*, December 19th, 1925.

RECTAL ANAESTHESIA.

BY

ROBERT B. COLEMAN, M.B., Ch.B.,
MEDICAL OFFICER, C.M.S. HOSPITAL, OLD CAIRO.

FROM the course of the correspondence on rectal anaesthesia appearing in the *BRITISH MEDICAL JOURNAL* during the last few months it appears that few surgeons practise it, and from communications received by me it would seem that some failures have been encountered. Such failures must, I think, be due to faulty method of administration, and it may probably be useful to describe that used at the C.M.S. Hospital at Cairo, where it has invariably proved perfectly safe. It has been the routine method since I first saw it used by Dr. Stiven at the Government Hospital, Port Said.

We confine it almost entirely to operations on the head, neck, chest, or upper extremity. Spinal anaesthesia by stovaine, which we find so eminently satisfactory, with an experience of nearly 8,000 cases, as described in the *JOURNAL* of March 21st, 1925, is our routine procedure for all operations up to the costal margin. Naturally, however, the ether method can be used for most operations.

The mixture used is three parts of ether to one part of olive oil, and the amount given is regulated by the weight of the patient, the allowance being one ounce of the mixture for every 20 lb. body weight. In actual practice we find this estimate is rather on the conservative side, and in cases requiring more perfect anaesthesia we often order a small quantity in excess of this, up to half an ounce for an average adult. We consider that about eight ounces is a maximum dose; this means six ounces of ether.

The anaesthetic is given to the patient in the bed, but some care is necessary in choosing a suitable room. The ideal condition is to have a small private ward, the windows of which can be darkened by shutters or blinds. It should be in a quiet place, and if possible not too far from the theatre, and preferably on the same floor, to avoid disturbing the patient by lift or stairs. If given in a general ward this should be darkened and quietened if possible, and screens kept round the bed. The stretcher and the macintoshes, etc., are put under the patient previously, so that he may be moved to trolley or carried without risk of waking.

An enema and rectal washout having been given some hours previously morphine with atropine is injected hypodermically an hour and three-quarters before the time fixed for the operation; fifteen minutes later administration of the mixture is commenced.

The mixture, slightly warmed, is run into the rectum high up by a catheter attached to a tube and glass funnel, which should not be raised to any appreciable height. All the apparatus should be well warmed before use to avoid cooling of the mixture. It is important that this should be given slowly, and it is reckoned that the administration should take twenty to thirty minutes. This is by no means always an easy matter, especially as sometimes the patient attempts to strain and return it, but with patience and tact it can generally be accomplished. It is found to facilitate the giving of the whole dose if it is preceded and followed by just a little olive oil. The administration can safely be entrusted to a competent sister, nurse, or orderly. At the completion the catheter is clipped, and great care is taken to withdraw it gradually, so as not to excite straining. A damp cloth is put over the patient's face to prevent excessive evaporation, and he is allowed to lie quietly till required for the theatre. With the above