FIG. 1.—CASE 2.

FIG. 2.—CASE 3. Lesser curve ulcer with spasm opposite. Duodenal cap normal.

FIG. 3.—CASE 4.

FIG. 4.—CASE 5

FIG. 5.—CASE 6.

FIG. 6.—CASE 7.

that the immediate cause of his trouble was mechanical I had first to consider how this could be relieved. Something more than aspiration through a needle was required, but drainage, from the risk of secondary infection, was out of the question. Under local anaesthesia, and without even moving the patient in bed, I made a small opening in the eighth space, into which I fitted a smooth silver obturator as a plug. Through this opening, which remained perfectly clean, we removed every morning from one to two pints of fluid. At the end of a fortnight his condition had so far improved that I was able to resect a rib and to explore the chest. On opening the pleural cavity twelve pints of milky fluid escaped. Both surfaces of the pleura were covered with tubercles. The wound was closed in the hope that adhesions might form. Six weeks later, fluid having been repeatedly aspirated in the meantime, the whole of the left dorsal region was blocked with novocain, and the posterior portions of the ribs from the fifth to the eighth were resected. As the ninth and tenth had previously been resected a satisfactory collapse of the lower part of the chest was thus obtained. The result was very remarkable. He made an uninterrupted recovery, and a year later he was in perfect health, doing full work on a farm. He has remained perfectly well, there is no sign of fluid in his chest, and the external deformity is trivial.

CONCLUSION.

The thorax thus presents to the surgeon peculiar difficulties and great opportunities. The difficulties are largely mechanical, and they can be overcome by mechanical means. Given the necessary appliances, and the still more necessary experience, there is no region which can be explored with more safety or with greater thoroughness. Many of the conditions have already been brought under our control, and the conquest of many others is only a matter of time. But this of all the branches of our art is the one in which co-operation between physician and surgeon is the most essential, and I therefore welcome this opportunity of bringing it before a body of practitioners who are meeting with these cases every day. I have only touched on a few features in a very large field, but I hope that I may have enabled you to realize the great opportunities, the brilliant hopes, and above all the safety, of modern thoracic surgery.

An Address ON GASTRIC DIAGNOSIS.* BY

L. R. BROSTER, O.B.E., D.M., M.Ch., F.R.C.S.,
ASSISTANT SURGEON, CHARING CROSS HOSPITAL; SURGEON, QUEEN'S
HOSPITAL FOR CHILDREN.

(With Special Plate.)

THIS paper contains an analysis of the diagnosis of four related gastric conditions of similar symptomatology, based on operation findings. From the patient's point of view the stomach is the most plaintive organ of the body, and we all know that our methods of diagnosis and treatment of it are not yet so standardized as to meet the requirements of each individual case.

Having been personally interested in the correlation of the x-ray diagnosis and the operative findings, I wish to direct attention chiefly to this aspect, and, with the aid of a series of plates supplied by Dr. S. Cochrane Shanks, to distinguish the finer points of diagnosis.

I will, so far as possible, limit myself to the analysis of some 140 cases treated at the Charing Cross Hospital; these include 51 cases of gastric ulcer, 33 cases of duodenal ulcer, 27 cases of carcinoma of the stomach, and 26 cases mimicking these lesions, but due to other causes.

PROPHYLAXIS.

We have not been able as yet to determine the exact causative factor in gastric and duodenal ulcer. There is no doubt that predisposing causes exist. The patient who early seeks medical advice can expect amelioration of his symptoms from judicious treatment, while in the later stages he can be saved much pain and bouts of semi-invalidism, and fear less the complications which threaten his life.

The practitioner whose aim it is to prevent the occurrence of these lesions has to devise a daily routine for those who seek his guidance. Such questions as diet, the treatment by drugs, the restriction of smoking, proper rest, the elimination of oral sepsis, and of foci of infection must be attended to.

When the disease is established examination by means of the x rays and the fractional test meal should be employed. It is not until such treatment has been tried and has failed to prevent recurrence of the symptoms that the need for surgery arises. It then becomes a question of co-operation between the physician and surgeon, and the best results are obtained when this partnership is continued after the operation. The patient is prone to think that an operation means an end to his disabilities. It is true that most patients are cured, but some are only improved; a few with gross lesions cannot expect to become entirely normal again, while in a small percentage of cases jejunal ulcers will develop. It is to this small minority that continued after-care is so important.

DIAGNOSIS

With regard to the efficacy of the diagnostic means at our disposal, opinions differ. M'ynihan and the Mayos base most reliance on the x-ray findings, Walton stresses the clinical history, while others place their faith in the fractional test meal. The necessity for such indirect methods indicates a very varied symptomatology, and a lack of definite physical signs.

This being so, I am convinced that when taking the patient's history it is necessary to eliminate, as far as possible, matters of fancy, and to consider only the bald facts. For this purpose I use a chart, which is comprehensive and standardizes the results. I prefer to take clinical histories myself, and am often struck by how materially they can differ from those taken by my assistants. In reviewing the x-ray and operative findings they are found to agree in the majority of cases, and in this series the margin of error worked out as follows:

Gastric Ulcers.

Of 49 cases—		
History and x rays disagreed in 13—		
History correct, x rays wrong or doubtful in ...	8	
History wrong, x rays correct in ...	4	
History doubtful, x rays wrong in ...	1	

Duodenal Ulcers.

Of 28 cases—		
History correct, x rays wrong or doubtful in ...	8	
History wrong, x rays correct in ...	4	
History and x rays wrong in ...	3	

Carcinoma.

Of 22 cases—		
The x-ray findings were wrong or doubtful in ...	5	

Whatever the merits or demerits of these various forms of investigation may be, the main point is that we each evolve our own particular lines of thought and action, and are best fitted to carry these into effect. Personally I consider a good x-ray examination a necessity. Apart from the diagnostic point of view, it is an invaluable guide to the surgeon in planning his operation, and revealing such complications as a small coexisting ulcer. Should there be any doubt, or should the x-ray examination and the clinical history disagree, then positive evidence from the test meal becomes a deciding factor.

SYMPTOMATOLOGY.

1. General.

Perhaps the most striking features of gastric and duodenal ulceration are its selectivity for the male sex, and the definite periodicity of its symptoms. Bouts of pain lasting generally from two to three weeks are followed by intervals of complete freedom ranging from a few months to a year or more. In duodenal ulcer these remissions are most constant; the patient has one or two attacks a year, and knows sometimes exactly when the next one is due.

In the early stages the intervals are not so marked as when the ulcer is established. In the later stages any change in this periodicity becomes significant; the painful attacks become prolonged, while the periods of freedom

* Delivered to the Southampton Division of the British Medical Association.

become shorter. Such a change is invariably associated with some complication, such as pyloric stenosis, hour-glass contraction of the stomach, or adhesions to the pancreas. When the change is sudden, and the attack lasts much longer and the pain is constant, the onset of carcinomatous change in the ulcer must be suspected.

What happens to the ulcer in these symptomless intervals? Operation at these times reveals the ulcer still active or in a smouldering condition. Occasionally the scar of an old healed ulcer is found, and it is fair to assume that healing has occurred when a painless interval has lasted an inordinately long time.

2. Age and Duration.

Although ulcers may appear at unexpected ages, the average age of incidence lies between the third and fourth decades. The average for duodenal ulcers is a little earlier than for gastric ulcers, and carcinoma follows after the forties. In this series the age incidence and duration of the symptoms work out as follows:

	Age.	Duration.
Gastric	44	6.5 years.
Duodenal	41	6.6
Carcinoma	53	5.0
Non-ulcers	38	5.7

In women the onset is slightly earlier than in males, and the ratio of females to males is: gastric 7:51; carcinoma 9:27; duodenal 7:33; non-ulcers 7:26.

3. Pain.

Although pain is popularly regarded as a *sine qua non*, it is important to remember that certain ulcers are unassociated with pain, and manifest their presence by a sudden perforation or haemorrhage. I have recently met with two perforations of this type, where a careful interrogation failed to elicit any previous history of dyspepsia. One patient was a workman, who twisted himself on a scaffolding, and walked into hospital with slight rigidity of the right upper rectus and a pulse of 72. The other patient experienced slight pain for the first time, which he relieved with oil, and perforation occurred the next morning. Apart from slight rigidity and a pulse of 80, he had no symptoms of dyspepsia.

The intermittent character of the pain has already been referred to. It seldom occurs within half an hour after a meal, but may appear at any time up to the next meal. As a general rule, in uncomplicated cases the time of onset is delayed the farther away the ulcer is from the cardia, and a two-hour limit is roughly the dividing line between the pyloric and duodenal ulcers. Each has a food fluctuation. In gastric ulcer it is food, comfort, pain, comfort; whereas in duodenal ulcer it is food, comfort, pain. The relief afforded to a duodenal ulcer by taking more food is often referred to as the "hunger pain."

The pain varies in intensity—sometimes dull and aching, at other times sharp, boring, or burning. It is determined partly by the acidity of the gastric contents, by the nature of the food, and by spasm. The time of onset coincides with the crest of the acid wave of the fractional test meal. The localization is in the epigastrium. With posterior and adherent ulcers it spreads to the back, and may be referred to the point of the shoulder. In duodenal ulcer the tendency is to spread to the right iliac fossa, and in jejunal ulcer to the left side. Radiation with a more constant pain is indicative of complications, while the sudden acute stabbing pain is pathognomonic of perforation. Relief of pain is more often occasioned by vomiting in gastric ulcer, and by more food in duodenal ulcer. The ingenious methods of obtaining relief devised by patients are as numerous as they are inconstant.

The feature of carcinoma following on an ulcer is the change from a fluctuating to a constant pain, which is not relieved by the ingestion of more food; whereas in the primary carcinoma the pain may be so insidious that it is merely described as discomfort, and in a man over 40 should arouse suspicion. The pain caused by non-organic lesions of the stomach is more variable, and tends to come on earlier after food. The average time of onset of pain after food in this series was: In gastric ulcer, 1.7 to 2.4 hours; in duodenal ulcer, 2 to 2.4 hours; in carcinoma, never later than 1½ hours; in non-ulcers, ¾ to 1½ hours.

4. Vomiting.

In gastric ulcer vomiting is a common feature and occurred in 88 per cent. of the cases; it was almost equally common in the cases of carcinoma. In the duodenal ulcers and non-ulcer cases it occurred in about 50 per cent. of the cases. Walton states that vomiting in duodenal ulcer occurs in only 8 per cent. of cases. It is a reflex act conditioned by the pain, and becomes a more constant feature with the onset of complications. The fact that emptying of the stomach so often relieves the pain has encouraged patients to devise various means of inducing it. The incidence of vomiting to the number of cases was: Gastric ulcer, 43/51; duodenal ulcer, 16/33; carcinoma, 20/27; non-ulcer, 14/27.

5. Haemorrhage.

Haemorrhage from a chronic ulcer is always an alarming occurrence, and more so if it is duodenal in origin. Cases are published from time to time where it has been so severe as to warrant immediate operation, and the base of the ulcer has been found eroding one of the larger arterial branches. Death seldom occurs from haemorrhage. Medical treatment is invariably successful, and operation is better deferred. Blood transfusion has enabled us to deal far more effectively with this condition than hitherto. In severe and repeated haemorrhages it should always be resorted to, although I would add that special care is necessary in dealing with Group II cases.

Most perplexing are those cases of gastrostaxis, first described by Hale-White, in which the mucosa literally weeps blood and no demonstrable lesion can be found. I recently treated a young woman for this condition. Her x-ray examination showed a large filling defect on the greater curvature which was reported as an ulcer. The test-meal acidity was low. At operation no sign of ulcer could be found, but the whole mucosa oozed with blood. This patient was subsequently treated with hydrochloric acid. She remained fairly comfortable on 12 minims thrice daily, but any reduction of this amount caused her pain and haemorrhage to reappear.

Haematemesis occurs in about 30 per cent. of gastric ulcers. In duodenal ulcer regurgitation of blood may take place through the pyloric sphincter, and the more common feature of melaena may be overlooked when the blood is passed in small quantities.

Altered blood in the vomit, the so-called "coffee grounds," is found with carcinoma. I can offer no adequate explanation of the cause of the haematemesis or melaena associated with such cases as fibrosis of the appendix, Lane's kinks, adhesions, gall stones, and tuberculous mesenteric glands which have occurred in this series.

	Cases.	Haematemes.	? Haematemes.	Melaena.	? Melaena.
Gastric ulcer	51	15	3	5	2
Duodenal ulcer	33	4	2	7	2
Carcinoma	27	7	1	6	1
Non-ulcers*	26	8	—	2	—

* Haematemesis and melaena in 4.

6. Physical Signs.

It is rare for physical signs to be of any importance in the diagnosis. Of them, tenderness in the epigastrium is by far the commonest, and rigidity of the upper recti comes next in frequency. The most helpful signs when present are visible peristalsis, a succussion splash, dilatation of the stomach, or a tumour. As would be expected, they are most commonly found in carcinoma.

	Tender.	Splash.	Peristalsis.	Rigid.	Lump.
Gastric ulcer	17	8	4	6	—
Duodenal ulcer	14	2	—	4	—
Carcinoma	10	2	2	7	9
Non-ulcers	11	2	—	—	2

7. Appetite and Loss of Weight.

Appetite is so capricious that loss of weight is a more reliable index of the patient's general condition. Macbeth's dictum, "Now, good digestion wait on appetite, and health on both," would be qualified in these more scientific days by saying that good digestion depended more on the free hydrochloric acid in the gastric juice. Thus we find that where the acid is high the appetite is good; where it is low the appetite is poor, as also with the onset of complications. Caesar, quite apart from his mistrust, may have had in his mind the dyspeptic type when he said:

"Let me have men about me that are fat; . . .
Yond' Cassius has a lean and hungry look."

The majority of these patients are certainly of the lean type, and many of them are afraid to eat. That this is so, the following average loss of weight shows:

Gastric ulcer	...	...	1 st. in 9 months.
Duodenal ulcer	...	...	1 st. in 21.5 "
Carcinoma	...	...	1 st. in 5 "
Non-ulcers	...	...	1 st. in 8 "

8. Teeth.

The occurrence of septic teeth as a focus of infection is a diminishing factor, even in the hospital type of patient, since the general public and the medical profession are becoming increasingly alive to its consequences. There is thus little justification for it ever being overlooked in the class of case under consideration. The dental examination should not be confined to septic gums or teeth, but should include radiography of the roots for hidden apical abscesses. At this point the tonsils and accessory sinuses should also receive attention.

9. Constipation.

Constipation *per se* as a symptom of ulcer does not exist, unless there is an organic delay, or an association with some cause of stasis in the lower alimentary tract. That patients frequently suffer from constipation must be regarded more as a direct result of the habitual taking of bismuth.

10. X Rays.

I consider the x-ray examination of the patient after the administration of an opaque barium meal to be the most accurate means of investigation, when performed by an expert radiologist; and it must be remembered that evidence is often obtained by screen examinations which is not always recorded on the films. In unskilled hands the results are often useless and misleading.

Walton estimates that in gastric ulcer the results are correct in 90 to 95 per cent. of cases. In duodenal ulcers this standard is not so high, and the early cases of growth are difficult to distinguish from ulcer. The evidence is either direct or indirect. The classical direct evidence is an ulcer crater on the lesser curve filled with barium, the so-called niche, with a notch on the greater curvature opposite it, or a filling defect, an hour-glass stomach, an obvious delay, or an irregular duodenal cap. With indirect evidence we find such signs as rapid emptying, active peristalsis, spasm, and local tenderness to pressure. Occasionally the evidence may be misleading, the symptoms being due to conditions other than ulcer.

In all the cases under consideration the x-ray reports or films have been checked by the operative findings.

(a) *Gastric Ulcer*.—In this series there were 36 cases, of which the diagnosis was correct in 29—an average of 83 per cent. Of the 7 wrong, 3 were posterior ulcers, 2 being diagnosed as caecal stasis; 1, thought to be a lesser curve ulcer with delay, was a pyloric diverticulum; 1 diagnosed as duodenal ulcer was found to be pyloric; in 1 there was no evidence of ulcer; a recent ulcer had perforated and then healed; 1 showed rapid exit of food for a lesser curve ulcer with severe haemorrhage. The chief difficulty in diagnosis is, therefore, the posterior ulcer, and it is probable that with the improved technique of taking photographs obliquely or laterally in order to silhouette the posterior wall of the stomach, more of these ulcers will be demonstrated.

(b) *Duodenal Ulcer*.—The evidence is less precise in the case of the x-ray reports of duodenal ulcer. Rapid emptying and peristalsis, the classical textbook syndrome, do not figure so prominently as do delay, irregular cap, no cap, or a combination of delay with abnormalities of the cap. Out of 20 cases, 14 were correct—an average of 70 per cent.

(c) *Carcinoma*.—Of 18 cases of carcinoma, 12 were correct and 6 wrong—a percentage of 66. Of the 6 wrong, 4 showed no definite deformity; 1 showed duodenal obstruction; 1 showed duodenal ulcer. It is not uncommon for duodenal lesions to be confused with those of growth.

(d) *Non-Ulcers*.—It is of more interest to study those cases in which the x rays showed some evidence of an organic lesion which was not confirmed operatively. It is impossible to classify these, so I give the results *seriatim*.

	X-Ray Examination.	Operation.	Test Meal.
S. M.	Seven hours' delay and caecal stasis	Appendicectomy and Lane's kink	Ulcer.
W. L.	Rapid exit (? gastric or duodenal ulcer)	Appendicectomy and Lane's kink	
S. C.	Duodenal ulcer	Appendicectomy and adhesions	
H. D.	Duodenal ulcer	Appendicectomy and visceropexia	Ulcer.
W. B.	Hypertonic and irritable stomach	Appendicectomy and adhesions	
N. C.	Abnormal pylorus (? ulcer)	Appendicectomy and adhesions to gall bladder	
M. S.	? Lesser curve ulcer	Adhesions; post-gastro-enterostomy	Ulcer.
F. C.	—	Appendicectomy and adhesions	
W. R.	Eight hours' delay; tender pylorus	Appendicectomy and nodule on pylorus	Ulcer.
S. W.	Lesser curve crater and spasm	Nothing found	
E. W.	Hour-glass	Gall stones and band across stomach to splenic flexure	Low.
J. C.	Seven hours' delay; low dilated	Gall stones and tubercle of ileum	
R. J.	Filling defect; seven hours' delay	Hypertrophic tubercle of ileum with omental band fixed in pelvis	
M. F.	Big ulcer on greater curve	Gastrostaxis	

GASTRIC ANALYSIS.

The diagnosis of gastric lesions by means of the fractional test meal alone is apt to be misleading, but it often gives very valuable corroborative evidence. This is due to the fact that the acidity is subject to a wide range of variation. Ryle and Bennett, in a study of 100 healthy students, found this variation of free hydrochloric acid in the majority of normal stomachs. In those cases where there is any doubt in the diagnosis or a conflict of other evidence, the test meal may be very useful in coming to a conclusion, more especially in cases of duodenal ulcer and carcinoma. In about 75 per cent. of these cases curves are obtained which conform to a fairly uniform standard. In duodenal ulcer the acid content is high, and the feature of the maintenance of its secretion after the second hour is characteristic. In carcinoma there is little or no free hydrochloric acid present. This low type of curve, however, may be found with the anaemias, gall stones, visceropexia, and achlorhydria gastrica.

The curves in gastric ulcer are most variable. Here the acid rises quickly after the meal, but is not continued as in duodenal ulcer. In the majority of cases the acid is increased, but it may be normal, or even low. Curves similar to this group are sometimes found associated with chronic appendicitis, with or without adhesions, a Lane's kink, and occasionally cases of carcinoma will retain their ulcer acidity.

In pyloric stenosis there is a wide disproportion between the free and total acids, the latter being high as a result of fermentation processes.

OPERATIVE TREATMENT.

When a chronic ulcer fails to heal after continued medical treatment, surgical treatment becomes necessary. The operation of gastro-enterostomy has been most frequently practised for this condition. This procedure has a mechanical purpose, the anastomosis enabling the stomach to be more readily drained, and a physiological value, the acidity of the stomach becoming slowly neutralized by the alkaline bile, and the secretions of the pancreas and small intestine. The rest afforded to the ulcer by these means permits it to heal, and, it is said, delays the onset of malignancy. In cases of pyloric stenosis, for which the operation was primarily designed, the results are excellent; and in cases where the ulcer is distal to the anastomosis the results are good. Unfortunately it has been allowed to fall into disrepute through being performed for other conditions which do not warrant it.

Various methods have been practised from time to time which are directed against the ulcer itself, and these should be combined with gastro-enterostomy. They consist

in infolding of the ulcer, excision, or destruction by means of the cautery. This line of attack should be pursued in all cases of haemorrhage. In America Balfour excises the ulcer by means of a diathermy knife, and this destruction of the ulcer is an additional safeguard in forestalling the onset of malignant change. More recently a bolder technique has been employed in the form of partial gastrectomy; the ulcer-bearing portion of the stomach is removed, and the remainder united to the jejunum. This method is particularly advocated by Sir Berkeley Moynihan and the Mayo brothers. Other operations have been devised to meet special contingencies, and many modifications of the technique have been suggested, into the details of which I do not propose to enter.

In dealing with the grosser complications, such as hour-glass stomach and adhesion of the ulcer to the pancreas and to the surrounding structures, no hard-and-fast lines can be laid down. Most of these patients, whose general condition is very poor indeed, have to be treated according to their special requirements, and the ingenuity as well as the finer sense of judgement of the surgeon may be taxed to their utmost limits. Finally, every operation should include a thorough exploration of the abdominal cavity, especially of the gall bladder and appendix, for it is useless to operate on an ulcer and leave such a focus of infection untreated.

Operative Findings.

There were 22 cases of ulcer of the lesser curve of the stomach. Of these, 2 were kissing ulcers, 2 were associated with duodenal ulcers, 4 were posterior, and 2 were hour-glass.

There were 23 cases of pyloric ulcer. Of these, 3 were posterior, 4 were multiple, 3 were pyloro-duodenal, 3 were adherent, and 5 were associated with stenosis.

There was one case of jejunal ulcer; this was due to a silk stitch.

Operative Procedures.

Operation.	Gastric Ulcer.	Duodenal Ulcer.	Carcinoma.
Posterior gastro-enterostomy ...	32	26	12
Anterior gastro-enterostomy ...	2	4	2
Partial gastrectomy ...	7	1	1
Sleeve resection ...	4	—	1
Pylorotomy ...	1	1	—
Ulcers excised ...	3	—	—
Ulcers cauterized ...	2	—	—
Ulcers infolded ...	3	3	—
Double gastro-enterostomy ...	1	—	—
Gastro-gastrostomy ...	1	—	—
With appendicectomy ...	—	5	—
Inoperable cases ...	—	—	11
Total cases ...	51	33	27
Deaths ...	3	1	1

Total mortality=4.5 per cent.

BRIEF REPORTS OF CASES.

Case 1.—C. S. for some years had had intermittent attacks of epigastric pain radiating to the back; it was of a drawing or burning character, and came on one to two hours after food; it was relieved by bismuth. The appetite was good, but the patient was afraid to eat; had lost 8 lb. in four months. There was tenderness over the epigastrium. X-ray examination suggested pyloric ulcer. Test meal showed delayed secretion for one hour, then hyperchlorhydria. At operation an ulcer was found just beyond the pylorus.

Case 2.—Mrs. F. C., aged 60, had suffered from indigestion for fifteen years. Immediately after food there was an intermittent heavy feeling in the epigastrium which radiated to the right iliac fossa; it was relieved by vomiting and hot drinks. Haematemesis had occurred on several occasions. The appetite was poor, and the general condition feeble; she had lost 4 st. in the last few years. X-ray examination revealed an hour-glass stomach. At operation a chronic ulcer was found on the lesser curve; sleeve resection was performed. (See Plate, Fig. 1.)

Case 3.—T. G., aged 63, for twelve months had had a gnawing pain three hours after food; it was relieved by taking food. The patient was awakened between 2 and 3 in the morning, and had been much worse in the nine weeks previous to seeking advice. Vomiting sometimes accompanied the pain. The appetite was good; there had been slight loss of weight; splash. X-ray

examination: seven hours' delay; ulcer crater on lesser curve; notch on greater curve. Operation, partial gastrectomy, revealed an ulcer on the lesser curve and a coexisting ulcer on the anterior inferior portion of the first part of the duodenum. The history suggested duodenal ulcer, but the x rays showed only gastric ulcer. (See Plate, Fig. 2.)

Case 4.—R. J., male, aged 57, had for two years had dull aching pain across the epigastrium and colicky pain across the lower abdomen. The pain came on two hours after food and was relieved by vomiting. The vomitus resembled coffee-grounds. He was afraid to eat, and had lost half a stone in three months. There was general tenderness over the abdomen. The x rays showed a filling defect of the stomach, with seven hours' delay. A test meal was not estimated owing to the presence of bile in each specimen. Operation showed that the filling defect was caused by a rolled up piece of omentum fixed to the pelvis and dragging on the stomach; the ileum and mesenteric glands were tuberculous. (See Plate, Fig. 3.)

(a)

(b)

(c)

Typical Test Meal Curves: (a) Duodenal ulcer; (b) gastric ulcer (no radiographic signs); (c) carcinoma of greater curvature.

Case 5.—E. A., male, aged 51, complained of pain of a tingling and later of a boring character localized to the left epigastrium and radiating downwards; it was relieved by various means. For the past twelve months it had come on once in every twenty-four hours, and he had induced vomiting to relieve it. His appetite was good, but he had lost 2 st. in three months. X rays revealed a typical lesser curve ulcer. The test meal showed very low acidity. At operation a large chronic ulcer was found high up on the lesser curve. (See Plate, Fig. 4.)

Case 6.—J. F., a woman aged 40, had for seven years had intermittent attacks of pain, lasting about a fortnight, with intervals of freedom lasting for several months. Pain of a sharp nature coming on half an hour after food, and aggravated by food, was relieved by vomiting; the pain extended to the shoulders. She had lost a good deal of weight; there was resistance in the epigastrium, and rigidity. Occult blood was present. X rays showed a stomach of small capacity; over seven hours' delay; large filling defect of pyloric antrum. Inoperable carcinoma. (See Plate, Fig. 5.)

Case 7.—Mrs. F., aged 53, for two years had had constant dull aching epigastric pain conducted to the back; it was increased by taking food and relieved by medicine. There was almost daily vomiting after food. The appetite was poor, and she had lost a stone in six months. She was anaemic and wasted, and had enlarged glands in the axilla and groin. A blood count showed lymphatic leukaemia. X-ray examination suggested an ulcer on the lesser curve. A test meal showed hypochlorhydria (see chart). At operation (lesser curve ulcer), posterior gastro-enterostomy, an indurated mass was found in the caecum. Microscopy of appendix—lymphatic leukaemia. (See Plate, Fig. 6.)

Note.—Many of these cases were collected while I was surgical registrar, and I am indebted to the surgical staff, especially Mr. H. S. Clogg, for their kind permission to publish them. Also to Dr. Russell Reynolds and Dr. S. Cochrane Shanks for their willing co-operation in the radiological examination, and to Mr. Jocelyn Patterson for the test meal results.

METHOD OF TREATING ASTHMA BY RADIATION. (Preliminary Note.)

BY

S. GILBERT SCOTT, M.R.C.S., L.R.C.P.,
D.M.R.E.CAMB.,
RADIOLOGIST TO THE LONDON HOSPITAL.

THE amount of medical literature on the subject of asthma, and the number of methods of cure advocated from time to time, illustrate the humiliating fact that little is known of its pathology, and that no satisfactory line of treatment has been discovered. There are patients who pin their faith to one pet drug, which they hold in reserve, as they have found from experience that it gives some temporary relief of their distressing symptoms; others ever ready to try any new remedy; and lastly, those who have given up all remedies and simply "grin and bear it."

Let me state at once that I have no new theories as to the causation of asthma, but the results obtained from the