fetid odour. The caecum lay on the left side of the abdomen, with a retrocaecal appendix. The tense ascending colon passed from left to right across the fourth part of the duodenum, under cover of coils of ileum. The transverse colon lay coiled under the dilated stomach on the left side. The descending and pelvic colons and the rectum were normally situated and attached.

Between the pylorus and the splenic flexure of the colon there was no direct attachment of the intestines to the posterior abdominal wall. The duodenum had a mesentery. There was acute torsion of the third part of the duodenum, and, round this point as axis, there had been a volvulus of the entire gut distal to the third part of the duodenum and proximal to the middle of the transverse colon. The superior mesenteric artery was twisted spirally round the third and fourth parts of the duodenum. The presence of the artery, together with the acute torsion, effectually obstructed the lumen of the gut at this part. The vascular lumen was apparently also partially, or completely, obstructed.

The torsion of the duodenum was readily undone by twisting the mass of intestine from left to right, through an angle of 180 degrees. The duodenum was then seen as a straight tube

Diagram of position of viscera at birth. A volvulus of 180 degrees occurred, secondary to this condition, the caecum passing forward and to the left in front of the duodenum. The superior mesenteric artery was carried across the third part of the duodenum. A=Superior mesenteric artery. B=Duodeno-jejunal flexure. C=Transverse colon.

of the mesentery of the ileum. The ascending colon was also adherent to the root of the mesentery as it passed behind the duodenum. The rest of the gut had a free mesenteric attachment.

The mesentery was drawn tightly round the third part of the duodenum, drawing with it the superior mesenteric artery, which thus passed successively downwards behind the duodenum, forwards round the right side of the duodenum between the duodenum and ascending colon, then transversely across the duodenum, passing into the mesentery of the ileum. A large branch of the artery passed obliquely downwards, crossing anterior to the duodenum, and then behind the duodeno-jejunal flexure, to supply the jejunum. This artery appeared to cause partial obstruction to the duodenal lumen. The lumen of the colon was obstructed by the acute flexure at the junction of ascending and transverse colons. The caecum and ascending colon contained faeces of a normal yellow colour. The ascending colon was tense and smooth.

The specimen suggests a primary malrotation in a direction the reverse of the normal, bringing the colon behind the duodenum, this rotation being arrested, at the stage where the ileum had passed forward across the duodenum, by the firm adhesion of the transverse colon to the mesentery of the ileum. Any further rotation could only occur by torsion of the duodenum and volvulus. After birth the irregular peristalsis induced by the obstruction at the duodeno-jejunal flexure, and more especially at the junction of the ascending and transverse colons, determined the occurrence of the volvulus, which carried the caecum through an angle of 180 degrees across the duodenum to the left, twisted the superior mesenteric artery tightly across the third part of the duodenum, and thus produced acute duodenal obstruction.

Our thanks are due to Professor McIlroy and to Mr. Norbury for permission to publish the case.

A CASE OF EMBOLECTOMY.

BY

SIR CHARLES GORDON-WATSON, K.B.E., C.M.G.,
F.R.C.S.,

SURGEON TO ST. BARTHOLOMEW'S HOSPITAL.

MR. GEOFFREY JEFFERSON is to be heartily congratulated on the successful case of embolectomy recorded in the *BRITISH MEDICAL JOURNAL* of November 28th, 1925. In a leading article in the same issue some comments are made (to which I shall refer later) on a case upon which I operated in 1911; the details of this were published in a clinical lecture on post-operative embolism of the femoral artery in the *Clinical Journal* of December 10th, 1913.

The publication of Mr. Jefferson's case has stimulated me to record a case on which I operated in 1914 at St. Bartholomew's Hospital, and which I should probably have published before but for the war.

A married woman, aged 27, was admitted to the maternity ward at St. Bartholomew's, under Dr. Barris, on March 2nd, 1914. She was suffering from mitral stenosis, and had severe albuminuria with oedema of both legs. On March 4th she was delivered with forceps. On the following day marked dilatation of the heart was noted, followed by dyspnoea. During the morning of March 13th sudden acute pain was felt in the right foot below the knee, and no pulsation could be felt in the leg below the groin.

I saw the patient about five hours after the onset of symptoms. She was desperately ill, with feeble pulse and severe dyspnoea. The right limb was somewhat mottled below the knee, cold and insensitive. No femoral pulse could be felt. I exposed the common femoral artery under local anaesthesia, the patient being propped up in the sitting position as she was unable to lie down. I removed an embolus from the bifurcation of the artery with some recent thrombus from above, passing a probe up the artery until blood flowed. The artery was closed with fine vaselined silk, and pulsation was restored to the limb immediately after the operation. Shortly after the patient was returned to bed it was noted that pulsation in the femoral could not be obtained. The patient died the same evening, about three hours later. Before death pulsation was absent from both femorals.

Post-mortem examination showed that thrombosis had occurred in the right common femoral artery up to the bifurcation of the aorta and down to the junction of the external and internal iliacs on the left side. Mitral stenosis, bilateral pleural effusion, and chronic interstitial nephritis were also found.

It is doubtful if this patient would have recovered in any circumstances in view of the severe kidney and heart disease, but the rapid and extensive thrombosis above the embolism showed that the operation was performed too late. It seems probable that after the embolus had been removed the probe passed through and did not dislodge to any extent the thrombus, which possibly was even then extending up to the bifurcation of the aorta. So feeble was the action of the heart that the channel was soon blocked up again—a block which before death occurred extended to the left common iliac.

This case emphasizes the importance of early operation if any benefit is to accrue.

In the case which I published in 1913 (operation in 1911), referred to in the *BRITISH MEDICAL JOURNAL* of November 28th, 1925, an embolus occurred in the common femoral artery a fortnight after laryngectomy for carcinoma. At the operation the embolus became dislodged when the artery was lifted up in the wound in order to apply a Crile clamp before arteriotomy, and was checked at the bifurcation of the popliteal. I did not expose the popliteal but awaited events. Gangrene occurred below the knee. I subsequently amputated just above the condyles, and the patient made a good recovery.

It has been argued (*BRITISH MEDICAL JOURNAL*, November 28th, 1925, p. 1018) that the lower end of the popliteal should then have been exposed and the clot removed, a procedure which may present considerable difficulties. In the *Clinical Journal* I argued that immediate massage or pétrissage, as described and advocated by M. Paul Claisse (*Bulletins et Mémoires de la Société Médicale des Hôpitaux de Paris*, June 10th, 1910, p. 766) in reporting a successful case, might have been

successful in breaking up the embolus at the time of lodgement and before thrombosis had occurred. I still think that this might have succeeded and ought to have been tried. Such an opportunity of breaking up an embolus at the instant of formation, before thrombosis had commenced, is not very likely to occur again. In this case no serious thrombosis had occurred above the femoral bifurcation, as femoral pulsation was restored after dislodgement of the embolus, and remained intact down to the knee.

Massage, by the breaking up of a peripheral embolus, deserves more attention than it has met with, especially in the aged.

ACUTE SUFFOCATIVE PULMONARY OEDEMA.

PREVIOUS references to acute suffocative pulmonary oedema seem to indicate that this condition is less rare than was previously thought. The etiology is uncertain, but the notes and reports recently received throw a little light on this problem which has an important bearing on treatment. Hitherto such remedies as atropine, strychnine, and pituitary extract have been prescribed for the relief of symptoms in the absence of a clear conception of the underlying cause. Disturbance of the vasomotor mechanism of the lungs was suggested by Dr. F. B. Julian (April 17th, p. 697), while Dr. M. Cohen (March 20th, p. 528) ascribes the occurrence of the symptoms to a sudden transient relative failure of the left ventricle.

I.—ASSOCIATION WITH CARDIAC DISEASE.

ACUTE suffocative oedema of the lungs is an event so serious and so far from being a rarity that I may perhaps be allowed to give my conclusions, based on a series of cases from 1921 to 1925, which I studied in some detail. The one factor common to all is an advanced degree of cardio-vascular sclerosis. Valvular disease was present in some, but not in others; in all of them there was just prior to the attack or had been at some time evidence of auricular fibrillation. The onset is sudden in patients previously comfortable, or at any rate not distressed by their arrhythmia, and in no case was there peripheral oedema. During the attack the patient is only partially conscious and may have no recollection next day of any measures taken during the attack. With one of my series I had several opportunities of examining the chest very early in an attack, and the picture presented to the mind by auscultation is that of an attack of asthma without the lengthening of the expiratory phase; the oedema process creeps over the lung lobule by lobule, and with the extension the distress increases steadily. I have not in any of my cases been able to make out increase in the diameter of the heart over what it had been the day before. As to treatment, all my cases responded to hypodermic administration of atropine, with or without morphine, but in some venesection is necessary in addition.

I think the routine treatment for the cases with free expectoration should be 1/50 grain of atropine followed at once by venesection to about 20 oz.; as the attack subsides a further dose of 1/100 grain of atropine should be given with at least 1/4 grain of morphine to ensure sleep and give the neurones time to recover.

I have used adrenaline and also amyl nitrite, and found them without appreciable effect. The events are intensely dramatic, and the patient, who at 10 p.m. was moribund, at 10 a.m. next day may be enjoying his ordinary breakfast without an added sound in his lungs and without recollection of the events of the night before.

The pathology seems to be some vasomotor derangement. It certainly is not heart failure in the ordinary sense, for in one of my patients the attacks lessened and ceased as the ordinary signs of muscle failure appeared, and in the last few weeks of her life she was free from the attacks, though she had the peripheral oedema and other signs of progressive heart failure.

The late Sir James Mackenzie in a private communication said he had done autopsies on several of these cases without being able to discover the actual cause of death: that the dilatation of right heart theory was not true—enormous dilatation was common enough but never in

association with this peculiar oedema; that he had little doubt it was a vasomotor phenomenon and not explicable on a mechanical basis, but that he had never found or known of any sensible explanation, experimental or otherwise.

It seems that we must wait on the physiologists to enlighten our darkness by further experimental work directed to this particular subject, which may also help to elucidate angina pectoris.

Stafford.

A. E. HODDER, M.B.

OF the cases cited in the JOURNAL of April 17th (p. 696) and by Dr. Cohen, March 20th (p. 528), no fewer than eight out of the eleven were the subjects of valvular heart disease. Dr. Coleman's fifth case may possibly also have been. The cardiac condition is not mentioned in the remaining two cases.

The known association of auricular fibrillation with mitral stenosis raises the question whether acute suffocative pulmonary oedema may be due to paroxysmal auricular fibrillation.

The following case is significant.

An adult woman was admitted to hospital one evening under my care with an acute abdominal condition; there were abundant moist râles all over the chest. The pulse was so rapid that it was impossible to count it with any accuracy. The patient was in the same distress as is seen in acute suffocative pulmonary oedema. Though she was spitting up frothy mucus it was not pink or abundant. The heart beat did not sound irregular, but it was so fast that it was difficult to estimate the rhythm. Something made me suspect that the auricle was fibrillating, and when the ward sister told me that the patient had previously been in the ward with some heart condition I no longer doubted it. Neither a general anaesthetic nor any operation was admissible; either would have been fatal; 1½ drachms of tincture of digitalis was given at once, and at six-hourly intervals m 45, m 30, and m 15. The result was most gratifying, and next evening a laparotomy was performed under local anaesthesia, and a band which was obstructing the bowel was divided. The patient recovered.

I know that dyspnoea and attacks of nocturnal cardiac dyspnoea occur in patients whose ventricular beat is not so irregular as to suggest auricular fibrillation (I do not speak of the actual attack), but the presence of this condition may be demonstrated by a polygraphic tracing. I would suggest that a very careful investigation of the cardiac condition in all cases of acute suffocative pulmonary oedema may lead to an elucidation of the true nature of this condition. Such cannot, of course, be made during the urgent symptoms, but possibly some of the patients who survive the attack may show evidence of fibrillation in a venous tracing made when the urgent symptoms have subsided.

S. M. HEBBLETHWAITE, M.D.Lond., M.R.C.P.

Cheltenham.

THE following case of recurrent attacks of acute pulmonary oedema may be of interest.

The patient is a woman, aged 52 years. She has aortic incompetence, signs of which were first observed three years ago. Between her attacks she is very pale; the heart is moderately enlarged, and the systolic blood pressure in the arm is 160 mm. Visible pulsation in the carotid arteries and water-hammer pulse are well marked. There is oedema of the feet, but no signs in the lungs. She is irritable and difficult to manage.

During the past year she has had nine attacks of acute oedema of the lungs, all of a similar character. The interval between the first two was about three months, but they have gradually become more frequent, only a fortnight intervening between the last two attacks. They all occurred (with one exception) between 10 and 12 p.m., and nearly all on Sundays—perhaps brought on by excitement and overeating on these days. The onset is sudden and without any previous warning. On arrival, she is found making efforts to sit up in bed; she is anxious and terrified and appears to be aware of the danger of the attacks. The breathing is rapid and difficult. The face is pale and covered with a cold sweat. The lips and nose are livid. The fingers are cold and clammy. The pulse is feeble, small, and rapid. The temperature is subnormal. There is a profuse pink frothy expectoration and foaming at the mouth. After a short time consciousness is lost. Over the entire chest are heard numerous coarse bubbling râles. The attack lasts about two hours. On one occasion the patient had two attacks in one night, between which she recovered consciousness and lost her distress. On the day following an attack she feels quite well, and has often been found out of bed.

The treatment adopted in each attack consisted of a hypodermic injection of morphine 1/6 grain and atropine 1/100 grain, aromatic spirit of ammonia (1 drachm) by the mouth, and a hot-water bottle to the feet. If there was not much improvement in fifteen minutes the atropine was repeated.

London, N.

M. ROCKFELT, M.B., B.S.Lond.