Mr. NORMAN PATTERSON (London) asked Professor von Neumann whether, in doing a modified radical operation under a local anaesthetic, it was his practice to remove the incus. Professor VON NEUMANN replied that in about 80 per cent. of cases of chronic otitis the incus was disarticulated from the stapes, and in these circumstances he removed it; its removal did not adversely affect the hearing.

Mr. G. J. JENKINS (London), in reply, supported Professor von Neumann in advising meticulous care in the opening up of all cells of the mastoid and neighbouring regions in operations for acute mastoiditis. He was familiar with the group of cells between the dura of the posterior fossa and the facial nerve, and examples of this group could be seen in Mr. Cheate's collection in the Royal College of Surgeons. He knew of two abscesses of the cerebellum, just above and external to the jugular foramen, which seemed to be associated with the infection of this group of cells.

Mr. J. S. FRASER referred to tubal catarrh being responsible for the continuation of discharge after a radical mastoid operation, and spoke of the impossibility of curing it. His own experience of primary closure in a simple mastoid operation, undertaken against his wish and only from motives of dire necessity, had been followed by a fatal result.

PHARYNGEAL AND OESOPHAGEAL DIVERTICULA.

BY

WILLIAM HILL, M.D.,

Consulting Surgeon, Diseases of the Ear, Nose, and Throat,
St. Mary's Hospital, London.

THERE is probably no group of lesions in the descriptions of which so much ignorance and inaccuracy have been so frequently perpetrated and perpetuated as in those appertaining to the various etiologically, topographically, and clinically distinct types of diverticula of the pharynx and of the oesophagus. The taking stock and the straightening out of this subject which I have undertaken are long overdue. Pouches originating in the former region, the pharynx, are almost invariably described under diseases of the oesophagus, and with great frequency they are actually, but wrongly, designated oesophageal pouches and occasionally pharyngo-oesophageal pouches. They are neither. Moreover the textbooks, both general and special, ignore at least four separate varieties of pouch which I shall emphasize later—two forms being truly of pharyngeal, and the other two of oesophageal origin.

The following is a classification of the various species of diverticula of the upper alimentary tract which, whilst presenting some features more or less in common, are for the most part etiologically, patho-anatomically, and in their clinical signs and symptoms separate and unrelated entities. Their presentation in tabular form will help to clear away many of the prevalent misconceptions which have obscured in the past and still obscure these lesions.

A. Pharyngeal in Origin.

1. Congenital centrally placed anterior diverticula.
2. Congenital lateral post-faucial diverticula. (Synonyms: *Aeroceles*; *branchial diverticula*.) The mouth of these pouches is usually in the region of the palato-pharyngeal fold, but may be lower down in the glosso-epiglottic and in the pyriform fossae or near-by these. These pouches are due to embryonic developmental errors in the region of the branchial clefts. They form swellings on one side of the neck and of considerable size when of long standing. They contain air and mucous secretions, but rarely food, as the orifice on the lateral wall of the pharynx is usually so small.
3. Deep pharyngeal pressure diverticula or pouches. (Synonyms: *Pharyngeal pulsion pouches*; *Rokitansky's pulsion pouches*; *Zenker's pulsion diverticula*.) The mouth of these pouches is invariably situated posteriorly and mesially at the line of junction of the upper oblique and lower transverse sphincteric fibres of the inferior constrictor muscle. They are fairly common. (Pulsion pouches of the pharyngo-oesophageal junction—the *Grenz diverticula* of Rosenthal—and of the Lannier-Hackerman area at the upper end of the gullet are a figment of the imagination of untrustworthy and incompetent observers, and no *post-mortem* confirmation is obtainable

of their occurrence. Most authors misleadingly include these alleged varieties, but on insufficient evidence and faulty observation.)

B. Oesophageal in Origin.

4. Traction diverticula, situated mostly more or less at the level of, posteriorly to, the hilus of the lung—for example, to the bronchial glands area (*Rokitansky*). They may, however, occur elsewhere—for instance, in the supraclavicular gullet (*Chevalier Jackson*), and in the epiphrenic segment not far from the level of the diaphragm (*Starck*, *Luschka*, *Arnold*, *von Brosch*, *Halstead*, *Hurst*, and others). (Synonym: *Rokitansky's traction diverticula*.) They are not really rare, apparently, though not often diagnosed during life, as symptoms are usually absent.

5. The traction-pulsion diverticula of *Starck*. In these a pouch originating by traction subsequently becomes expanded by food pulsion. Food lodging in these pouches was held by *Rokitansky* to be a frequent source of septic mediastinitis and various suppurative diseases of the lungs and pleura of hitherto unexplained origin. Such sequelae, however, have not been considered common by subsequent authorities.

6. *Leugart's* post-bronchial ledge pulsion diverticula, situated behind the level of the left bronchus above *Leugart's* ledge. These diverticula are rare.

7. *Epiphrenic pulsion diverticula*, situated in the lower third of the gullet not far above the diaphragmatic level (*Zenker*, *Bensaude*, *Kraus*, *Starck*, *Hurst*, and others). They are rather rare.

8. Dissecting diverticula.

9. Diverticular sinuses.

CONGENITAL CENTRAL ANTERIOR DIVERTICULUM OF THE PHARYNX.

Considering the frequency of anterior centrally placed fistulae, sinuses, and cysts at the base of the tongue (second branchial cleft survivals) diverticular *aeroceles* might be expected to occur there, and also in connexion with the third and fourth branchial clefts lower down; but the only recorded instance, probably of this nature, with which I am acquainted is a case recently reported by *Hurst*, in which there was stated to be a central opening on the posterior pharyngeal surface of the cricoid plate with a diverticulum passing down between the trachea and oesophagus; this filled up with food, causing dysphagia. This could evidently be capable of surgical extirpation by a lateral incision in the neck. Although of congenital origin it was evidently increased in size by food pulsion. This exceptionally, if not uniquely, placed diverticulum was discovered by radiography, to which recourse was had to explain the dysphagia.

CONGENITAL LATERAL PHARYNGEAL DIVERTICULAR SINUSES AND AEROCELES OF ABNORMAL BRANCHIAL CLEFT ORIGIN.

Branchial fistulae opening both externally and internally, and branchial sinuses ending blindly in the neck, opening either externally only or, less commonly, internally only, are by no means rare lesions, and are well known to be due to embryonic developmental irregularities in the devolution of the branchial clefts; cysts, dermoid and otherwise, can be traced to the same source; it is only very exceptionally, however, that a sinus passing into the neck from an internal pharyngeal orifice expands into an *aerocele*. *Godlee* and *Bucknill*, however, in recording such a case in 1901, referred to a comprehensive paper by *Kostanecki* in 1890 which, whilst dealing generally with lesions of branchial origin, such as cervical fistulae, sinuses, and cysts, mentioned a number of recorded cases in which an internally opening sinus had developed into a well marked diverticular *aerocele*, which presented as an obvious air-filled swelling in the neck.

Watson of Edinburgh recorded the first case of this form of diverticular *aerocele* in 1875; found in a dissecting-room subject, it reached the level of the clavicle and resembled a Florence flask with a very elongated neck, which opened by a small orifice at the favourite spot just behind the tonsil on the palato-pharyngeal fold.

The next record was also British. The elder *Wheeler* described and illustrated an enormous air-filled cervical diverticulum (*aerocele*); it was as large as a fair-sized orange. At operation it was found to extend in front of the sterno-mastoid, and was covered by a thin layer of muscle which I surmise was the platysma; it reached the

clavicle below and the middle line over the thyroid cartilage in front. The stalk was traced upwards between the two carotids, and entered the pharynx in a space bounded by the palato-pharyngeus and the stylo-pharyngeus laterally and the middle and superior constrictors above and below. This diverticulum, Wheeler states, could be traced to the right sinus pyriformis, and the opening was apparently large in contrast to that of the lesions described by Watson and by Godlee. After removing this sac Wheeler unexpectedly discovered another diverticulum lower down, originating posteriorly from the deep pharynx and passing down behind the oesophagus. This true pharyngeal pulsion diverticulum was the size of a walnut, and was removed. Wheeler had, therefore, in 1886, the distinction of being the first to operate on, and to remove successfully, a congenital lateral pharyngeal diverticulum, and to be the first to remove a pharyngeal pulsion diverticulum with survival and cure of the patient; though the latter procedure of excising a pulsion pouch had been previously (in 1884) performed by Niehans and von Burckhardt, but with a fatal result in each instance.

These diverticula, though more commonly having a small opening in the region of the posterior faucial pillar just behind the tonsil, may, as in Wheeler's case, open lower down; the neck in Godlee's case passed through the lateral part of the thyro-hyoid membrane, and in another instance the opening was large enough to admit a finger somewhere in the region of the glosso-epiglottic fossa. Although the records describe these sacs as containing air and some secretions, and in one instance grumous material, it would be imagined that, where the pharyngeal opening happens to be large, food must sometimes get in; but perhaps there is a valvular condition of the orifice which prevents this happening as a rule. Certain it is that the presence of food in the sacs is not mentioned in any of the records I have consulted; but I cannot claim that my search has been exhaustive.

HISTORICAL NOTES ON PULSION DIVERTICULA OF THE DEEP PHARYNX.

According to Zenker the first recorded case of such a pouch was made by an Englishman (Ludlow) in 1764, and the specimen, a large one which extended into the thorax, is preserved in the Hunterian Museum of the Royal College of Surgeons. Zenker and von Ziemssen give references to no fewer than twenty-two cases, in which the diagnosis was rendered unequivocal by *post-mortem* examination, which were recorded between 1746 and 1879, and they add five cases of their own in the same category. In this list are included four English cases—namely, Ludlow's mentioned above, and one by Sir Charles Bell in 1816, one by Worthington in 1846, and one by Ogle (which is in the St. George's Hospital Museum) in 1865. Zenker also refers to seven recorded cases, in which, however, there was no autopsy, including one described by Dendy in 1848 and another described in the *Edinburgh Medical Journal* in 1856.

I have already alluded to the important case reported by Wheeler of Dublin in 1886, which was of especial interest in that it was not only the first one cured by surgical removal by a lateral incision, but the case was complicated by, and operation was undertaken for the removal of, a very large congenital lateral post-faucial diverticulum of congenital embryonic branchial cleft origin. The lower pharyngeal diverticulum was unsuspected, and only discovered during the course of the operation, and of course it was removed at the same time as the large lateral air pouch.

The first description, in any way approaching accuracy, of diverticula in the region under discussion dates from a communication by Mondirre in 1833. In 1840 Rokitsansky placed the subject on a sure basis. He it was who classified these pouches under two main headings—namely: (1) pulsion or pressure pouches, and (2) traction pouches. Although he devoted most attention to, and laid the foundation of our knowledge of, traction pouches of the middle region of the oesophagus, he showed remarkable perspicuity in pointing out that the pulsion pouches found high up really originated from the lower pharynx. Yet in spite of this clear statement, and the fact that the point was empha-

sized in 1887 by Zenker, these pulsion diverticula of Rokitsansky, originating high up in the alimentary tract, have been almost habitually included under the heading of "Diseases of the oesophagus," and have been described as originating either at the extreme upper end of the gullet or else at the pharyngo-oesophageal junction, and the importance of Killian's dehiscence in the inferior constrictor, with hernia of the mucous membrane through the same, has been almost ignored even to the present day.

After Rokitsansky, the next comprehensive and really illuminating account of diverticula of the pharynx and of the gullet was set forth with, on the whole, considerable accuracy by Zenker, writing independently in 1876, and also in collaboration with von Ziemssen in 1877, in the *Encyclopaedia of Medical Practice*, edited by the latter, the English edition of which appeared so far back as 1878. In volume viii of this work, under the section dealing with diseases of the oesophagus, they included, it is true, diverticula of the pharynx, for they write:

"In a careful investigation of the subject we must overstep the true limit of the oesophagus and consider also the lower portion of the pharynx at the same time, since some of these diverticula, strictly speaking, belong to the pharynx. . . . In the literature such pharyngeal diverticula are repeatedly and erroneously described as oesophageal."

This statement is surely sufficiently definite, yet even at the present day this erroneous nomenclature as regards these herniae of the pharynx is perpetuated in our textbooks, both general and special, and in the report of cases in current journals. Zenker, following Rokitsansky, divided diverticula in the upper food tract into those resulting from (1) pulsion and (2) traction, and stated that the two forms "are so totally distinct and different that it is impossible by grouping them to give a general description of the disease. Although this has been done by nearly all writers, excepting only Rokitsansky and Heschl." Zenker thought that pressure—that is, pulsion—diverticula were limited to the upper part of the alimentary tract, corresponding to the true pharyngeal diverticula, and that traction diverticula were limited to the gullet. The latter statement is apparently true, but it is not a fact that pressure diverticula do not occur far remote from the pharynx—that is, unequivocally in the oesophagus, and mostly in the middle third, but exceptionally both high up and quite low down. In reference to pressure diverticula, however, they state later that "they are located, as it appears, exclusively (certainly only with rare exceptions) at the lowest part of the pharynx just at the upper boundary." Although it turns out they were here right in not postulating too absolutely that these pulsion pouches were limited to the pharynx, they were not strictly accurate in placing them "at the lowest part of the pharynx just at the upper boundary of the oesophagus," for the pouches really originate, as was first definitely shown by Killian, at the junction of the oblique and transverse fibres of the inferior constrictor (cricopharyngeus) muscle and at least a centimetre above, not "just at the upper boundary of the oesophagus." Zenker, however, fully realized that the pouch was, in many cases at all events, surrounded at its neck *only* by "the transverse continuous fibres of the inferior constrictor muscle of the pharynx"; and further on in the article Zenker, in alluding to his dissections, disposed of the erroneous statement of others that the pouch has a muscular coat, more especially in the smaller sacs (Klebs); he and von Ziemssen assert this view to "be untenable when we consider the arrangement and origin of the fibres of the inferior constrictor through which the mucous coat passes."

In spite of this clear statement that these herniae of the lower pharynx pass through the fibres of the inferior constrictor, many writers (from Butlin in 1893 down to those of the present day—for example, Chevalier Jackson, Shallow, and the Mayo staff) have taught that they are at times situated at the junction of the pharynx and gullet, and imply that they are as much oesophageal as pharyngeal in origin in some instances. In my opinion they are wrong.

Others, again, have considered that some of these pouches originated definitely just below the lower margin of the pharynx and through the posterior wall of the

extreme upper end of the oesophagus, 'in the alleged Lannier-Hackerman weak area. Killian proved conclusively that there were no known museum specimens which supported this oft-quoted view. Killian also showed that the pouch in the early stages consisted, as held by Zenker and von Ziemssen, of the fibrous submucous and mucous coats only of the pharynx. The fact that muscular fibres are found round the neck of moderately advanced and larger specimens is, in my opinion, explained by the fact that the hernia, when it attains a large size, tugs on the margins of the separated oblique and transverse portions of the inferior constrictor, and thus enlarges the mouth of the sac by traction and pulls down some of the fibres around the neck.

As regards the statement that there is a weak triangle at the postero-superior aspect of the extreme upper end of the gullet, where the muscular fibres are stated to be either sparse or absent, this heresy owes its origin to Lannier and Hackerman, and has been repeated, on absolutely insufficient evidence, by many writers in textbooks and special articles. In 1910 Waggett and E. D. Davis made a number of examinations of excised gullets in the *post-mortem* room, and conducted pressure experiments, and found that there was little warrant for such an assertion; as a matter of curiosity I repeated these observations and confirmed them.

The long and short of the matter is that what are usually known as Zenker's pulsion diverticula are entirely pharyngeal in origin, and do not bulge through either the pharyngo-oesophageal junction or through the extreme upper segment of the gullet in the region of the Lannier-Hackerman triangle, and any nomenclature which suggests any origin other than through Killian's dehiscence—that is, between the two portions of the crico-pharyngeus—is not warranted. Amongst writers who have promulgated the erroneous views that some pressure pouches occur in this situation *below* the lower margin of the inferior constrictor are Morell Mackenzie, Butlin, Zesas, Knott, Halstead, Starck, Stetten, and the majority of British and foreign writers on the subject, down to contributions this year by Shallow and Chevalier Jackson. John Morley, like StClair Thomson in his authoritative *Handbook*, correctly describes these pouches in the text, but Morley, under the wrong title of diverticula of the oesophagus, and Thomson deals with them under the section devoted to oesophageal diseases. A recent notable exception deserves mention—namely, the paper on pharyngeal diverticula by Wilkie and Hartley in the *British Journal of Surgery* (1922), which is refreshingly free from the usual errors of fact and from misleading nomenclature.

DIAGNOSTIC SYMPTOMS AND SIGNS OF PHARYNGEAL PULSION POUCHES.

The classical symptoms recorded by Rokitsansky, and later by Zenker, well back in the last century, were so much to the point that they stand to-day. Dysphagia, intermittent in the early stages and very gradually getting worse and more constant after years; immediate regurgitation of food at times, as in other forms of stenosis of the pharynx and gullet, but associated also with slight regurgitation of portions of food long after a previous or a meal or even two before the last; variable discomfort in swallowing rather than any difficulty in getting down ample nourishment is the rule and this not merely in the early stages, so that extreme loss of flesh is most exceptional and slight loss not so frequent as might be expected, even when the pouch is of long standing; aphagia necessitating gastrostomy appears amongst the records of last century; gurgling noises due to the mingling of swallowed air with the food stagnating in the pouch; in advanced cases there may be an obvious, visible, and palpable swelling on one or other side of the neck, more usually the left; external pressure may lead to ejection of the contents when food is present in the pouch. Discomfort rarely amounting to actual pain may occur from temporary overdistension and imprisonment of ingesta in the sac. Pouch patients are also liable to get distressing fits of choking and coughing sometimes, becoming actually cyanosed when food ejected from the pouch finds its way out suddenly and overflows into the air passages. In a case described and illustrated

by Robinson of Sunderland the pouch extended upwards towards the base of the skull behind the middle and upper pharynx as well as a long distance downwards behind the gullet, and when the patient swallowed water the neck swelled out laterally like the pouter pigeon, and for a few minutes at times presented the appearance of goitre.

A stricture high up in the gullet may exceptionally bear a superficial resemblance in symptoms, for a time, to a pouch, but a definite diagnosis can always be made, when efficiently carried out in the correct position, by an x-ray examination in conjunction with barium or bismuth paste. It is often useful partially to fill the sac through a rubber tube with paste instead of allowing the patient to swallow it. The endoscopic findings are characteristic but rarely really necessary for diagnostic purposes alone. The orifice of the sac is easily seen through the endoscope, but the lumen in the deep pharynx leading to the mouth of the gullet is often difficult to find even by experts. It can, however, be located by the swallowing of a shotted string.

TREATMENT OF DEEP PHARYNGEAL PULSION POUCHES.

When Zenker wrote his articles in 1876 and 1877 the only treatment recommended when the condition of the patient approached aphagia, and when there was failure to insert easily a rubber or gum-elastic feeding tube through the prolapsed pharyngo-oesophageal orifice into the gullet, was the performance of gastrostomy. He and von Ziemssen wrote, however, as follows:

"The radical cure of diverticula by operative procedure from without is one of our vain wishes; yet we should hope that even this operation conducted on Lister's plan might at some future day be performed without danger."

This "vain wish," first suggested by Kunge in 1864, and then by Hamburger in 1871, and attempted, but not with survival of the patient, by Niehans in 1884, was first successfully realized by Wheeler of Dublin in 1886, when he encountered a deep pharyngeal pulsion pouch whilst removing an enormous congenital lateral pouch or aerccele, and he promptly excised both. Zesas and Niehans, von Bergman, Kocher, and Butlin repeated this operation between 1888 and 1892, and primary excision has remained the method of choice in the main ever since. A few operators—for example, Stetten—recommend a preliminary gastrostomy. This appears to me to be neither necessary nor even justifiable; at all events my seven cases were successful without it.

Primary excision of the pouch through a cervical incision along the anterior border of the left sterno-mastoid and dissecting down to the side of the pharynx and gullet, severing the omo-hyoid muscle, tying the superior thyroid vessels, and displacing the left lobe of the thyroid gland, is a comparatively easy procedure, and has been described in detail by Butlin and others. Operation should always be preceded by what the earlier operators called the "toilet of the pouch"—namely, washing it out clean with some mild antiseptic beforehand. A bougie in the sac and another in the gullet considerably helps to define these structures and aids in the subsequent separation of the pouch by blunt dissection. If a bougie does not pass readily into the gullet beforehand, it can easily be manipulated into it when the deep preliminary dissection to the region of the side of the gullet is made. When the pouch has been freed from its bed considerable difference of opinion prevails as to the method of cutting through the neck close to Killian's area of dehiscence. I think, however, that simple incision is sufficient, though some operators still prefer to ligature the neck first, and divide it distally with the electric cautery knife.

The crux of the whole operation depends largely on the method of sewing up the orifice left when the pouch has been removed. Formerly the mucosa was invaginated and the fibrous coat sewn up by Lembert's method of suture. At the present day, however, most operators resort to some form of purse-string suture. The method of Wilkie and Hartley will be alluded to later. John Morley advocates the technique adopted in closing a duodenal stump in a Polya gastrectomy, and considers that excision of a pulsion diverticulum "should only be attempted by those who have

served a long apprenticeship to the technique of gastrointestinal surgery." It would be thought that familiarity with the surgical anatomy and experience in surgical operations in this region of the neck were rather essential to operative confidence and success, and that one need not be an abdominal surgeon to carry out any method of suture for closure of the hiatus, in the lower pharynx, left after excision of the diverticulum.

Several alternatives to simple primary excision have been either advocated merely or else actually carried out. The former include painting the interior of the sac with silver nitrate (Dendy); dissecting out the sac free from its bed by a lateral incision, and then invaginating it into the food tract (Girard); the orifice being stitched up posteriorly to endeavour to prevent recurrence to the original position, which, however, has occurred sooner or later in several cases, and the method has fallen into disfavour on that account. Another method advocated, but not carried out, was that of Chevalier Jackson, of passing a volsella through the endoscope down to the interior surface of the fundus and endeavouring to perform internal traction invagination into the pharynx without making an external cervical approach. As I remarked in reviewing Jackson's larger work twelve years ago, the suggestion showed that he had no experience of dissecting out a pouch free of its adhesions, or he would have known that the latter would not yield to internal traction.

Statistics compiled by Stetten in 1910 showed that there was a mortality of no less than seven in the first thirty recorded cases of simple primary excision alone, though carried out by first-rate general surgeons for the most part. This high mortality of over 23 per cent. would probably have been greatly exceeded if he could have got at the number of unrecorded fatal cases. It is, however, significant that there were only two deaths in the next eighteen recorded cases, and, on the other hand, Stetten found that of five other cases, in which primary excision was preceded by gastrostomy, there was only one death, and that was avoidable probably, because obviously due to very injudicious after-treatment. Stetten recommends this preliminary measure which he himself followed, but I regard it as only exceptionally justifiable.

It was probably the high mortality that led up to the Goldmann operation of removing the pouch in two stages.

The two-stage excision is generally credited with having been first successfully carried out by Goldmann in 1907, but Halstead, in his article in *Surgical Clinics of Chicago* in 1919, claims that he first practised it, though he gives no reference to priority of publication over Goldmann, who recorded two successful cases in 1907. It does not appear to have been performed very frequently on the whole, though some operators prefer it in all cases, but others especially reserve it for very large pulsion pouches (Colledge).

The first stage of Goldmann's operation consisted of freeing the pouch from its bed by an incision along the anterior border of the sterno-mastoid and subsequent dissection as in the operation for the primary excision; instead, however, of cutting off the pouch at its neck, it was turned up with the fundus above, and the latter was stitched up to the upper angle of the operation field, the neck was lightly constricted with catgut, and the skin incision was then sewn up except at its lower end. Ten or more days afterwards, when the cervical lymphatics were presumed to have been sealed and the risks of mediastinal infection reduced, should subsequent leakage occur, the wound was reopened, the pouch excised, the neck suitably sutured, and closed as in the operation of simple primary excision.

Subsequently Goldmann tried a modification of this method: the pouch having been dissected free, a catgut

was tightly—not lightly as in the first method—tied round the neck of the pouch so as to ensure strangulation and necrosis of the pouch, which was packed up with gauze high up in the wound; the skin incision was only partially sewn up, being left open at the upper as well as at the lower end. The second stage consisted in removing the pouch when it had separated at its neck by sloughing a week or two later. This, of course, meant leakage. It was a dirty procedure and proved not to be free from risk, though Marsik and others adopted it for a time. When the Goldmann two-stage method is mentioned to-day it is the first type which is usually referred to.

Wilkie and Hartley modified the first Goldmann operation, which they prefer to primary excision, by incising the pouch and performing a submucous resection of the mucosa, which they remove, leaving the fibrous coat, which they then invaginate and sew up by purse-string sutures.

Bevan's method of obliterating the cavity of the pouch without excision, in order to reduce the risk of leakage, is theoretically ingenious. The sac having been dissected out free from its bed under local anaesthesia, Bevan proceeds, if the pouch is of moderate size, not more than 2 inches long, "to invaginate with three purse-string sutures of black silk or Pagenstecher linen one-half of the diverticulum into the other half, then the rest of the diverticulum should be obliterated by longitudinal sutures, six or eight, running parallel with the long axis of the diverticulum." When the operation is completed the area of the mouth of the pouch at Killian's dehiscence presents as a firm rounded button or knob on the posterior surface of the lower pharynx. In dealing with a large long diverticulum another method is employed by Bevan.

"The diverticulum can be crushed at its centre with a pair of heavy forceps, tied with a silk ligature, and the portion distant to the ligature cut off with the electric cautery. The remains of the diverticulum which are then no larger than one's thumb can be treated by invaginating them into the oesophagus" (meaning pharynx) "with three purse-strings just as we handle the smaller diverticula."

The patient is fed by a feeding tube passed for each meal for ten days.

Bevan does not allude to any cases in which these operations have been

carried out, so we know nothing as to whether they were successful for a time, or whether there was any recurrence of symptoms, as occurs sometimes after Girard's invagination; nor is anything said about subsequent leakage over having happened. The latter would appear to be not unlikely in any but a pouch with an extremely thick wall instead of the usual thin wall; and considering that from nine to eleven sutures are inserted it seems extremely likely that the wall would be perforated, and possibly leakage and sepsis result. In the significant absence of any statistical record of the cases operated on and the immediate and remote results, judgement must be suspended as to the value or otherwise of Bevan's really ingenious proposals.

My operation of upward suspension and fixation of the freed and inverted pouch was described under the name of "diverticulopexy" in connexion with a case in which I carried out this operation, shown at the Laryngological Section of the Royal Society of Medicine in 1917 and recorded in the *Proceedings* in 1918. It consists essentially of the first stage of the Goldmann operation. The pouch having been freed from its bed by dissection and isolated is then turned down-side up and the fundus stitched up as high as possible to the upper angle of the operative field by sutures to the side of the pharynx and to the deep cervical fascia under the sterno-mastoid. In order that stitches shall not perforate the sac wall and shall hold it, it is essential not to make a clean dissection of the sac, but to leave as much areolar tissue as possible adherent to

FIG. 1.—Typical posterior pharyngeal pulsion diverticulum showing the marked prolapse of Killian's lip.

its wall; before stitching the pouch up I pass a catgut ligature around the neck of the sac, but do not draw it sufficiently tightly to produce strangulation. The object of this is to prevent food entering the pouch in the first weeks following operation.

For a very large intrathoracic pouch, really too long for successful suspension, it might be advisable to do the second stage of the Goldmann operation if a radiographic examination showed that the pouch was inclined to prolapse after a short period. I do not recommend the operation for a small walnut-sized pouch, as in the only case in which I did a "pexy" in such circumstances the symptoms recurred from prolapse a year later, and I then resorted to excision. The method, therefore, is probably liable to fail in extremely long pouches extending well down into the chest, and also in rather small pouches. In addition to the case of the small pouch which recurred, I have done four other cases of suspension, with complete relief in each up to the present—that is, for five years or more. I also assisted Sir James Dundas-Grant in dealing with a rather long pouch by this method. There is, of course, no chance of leakage if the wall of the sac has not been perforated by suture and the dreaded mediastinitis is eliminated.

The cause of dysphagia in these diverticular cases is not, as is usually held, so much due to pressure on the oesophagus from behind by food in the pouch, as to prolapse or downward displacement of Killian's lip, which forms the lower border of the mouth of the pouch; this prolapse displaces the plane of the pharyngo-oesophageal junction from a horizontal into a more or less vertical plane, and prevents the normal opening up of the mouth of the gullet during deglutition. The suspension and fixation upwards of the pouch corrects this marked prolapse and draws up Killian's lip together with the posterior wall of the cervical gullet, into its normal position, and the patients are able to swallow normally the same evening in most instances. The prolapse of Killian's lip is well shown in museum specimens, but the point has not, I think, been previously emphasized.

Morley states that I and König advise suspension by suturing the sac to the hyoid bone or to the omo-hyoid muscle; that is the method described in 1922 by König, but has not been hitherto carried out by me. It may be worth while to use these points as additional attachments for the pouch.

Of course, the tendency of most "pexies" is to recur, but this only happened to me in the one instance of the small pouch above mentioned.

TRUE DIVERTICULA OF THE OESOPHAGUS: THEIR ETIOLOGY, SYMPTOMS, AND TREATMENT.

Very few pouches in the gullet have been recorded other than the traction diverticula of Rokitsansky behind the roots of the lungs; whether these latter are as prevalent as he asserted, though latent, and whether they are so often as he believed, the determining factor in serious lung and pleural abscesses and in mediastinitis, when decaying food is retained, is still a moot point.

Traction diverticula are usually unsuspected till found at necropsies; their cause is, on the other hand, evident, as they can always be traced to a previous breaking down of a suppurating lymph gland which forms an inflammatory adhesion to the outside of the wall of the gullet, which gets dragged out of position by subsequent healing and cicatricial contraction. They are rarely observed at any spot other than in the middle third of the gullet behind the area of the glands in relation to the bronchi and trachea. Jackson has, however, recorded a traction diverticulum of the supraclavicular gullet, and they have also been observed low down near the diaphragm, presumably from the adhesion of breaking down mediastinal glands. Traction pouches, which are normally narrow

funnels, may very exceptionally become greatly expanded by food lodging in them, and when this stagnation ectasia happens we have the traction-pulsion diverticula of Starck.

The rare true pulsion diverticula of the gullet, which occur usually in the lower half, though occasionally higher up, may sometimes be due to increased deglutitory pressure when there is either a benign inflammatory or a functional stricture of the gullet, as suggested by Hurst, but this would hardly apply to such pouches higher up, or where there is no obvious stricture. When pressure pouches are associated with and result from functional or from cicatricial stricture at the level of the diaphragm the treatment is obviously that of relieving the causal obstruction. Pouches which have no underlying stricture in the gullet, excepting Leugart's variety, do not themselves appear to have often any recognizable symptoms; they are at times accidentally discovered either at necropsy or when making radiographic or oesophagoscopy investigations, and there is no *ad hoc* surgical or other treatment either necessary or even possible beyond, perhaps, lavage.

Leugart's pouching is extremely rare. I have only seen one well marked example—namely, in a case I was asked to examine with Atkinson at the Central Throat and Ear

Hospital. In this case there was a well marked lateral recess in association with a prominent horizontal ledge projecting internally from the anterior and left lateral wall of the gullet at the level of the left bronchus. Both Leugart himself and Halstead thought that the gullet was actually compressed by the left bronchus itself in these cases; this is obviously wrong, as the ledge passes backwards and to the left, which is not in any way the direction of the left bronchus, of course, and the latter never comes directly in relation with the wall of the oesophagus. I think the probable explanation is that contracting connective tissue bands pass from the bronchus outwards and backwards to the left side of the body of that dorsal vertebra which lies behind the gullet at the level of the left bronchus. I have on many occasions when making an endoscopic examination observed a slight ledge in this position, and it is so frequent as to have been alluded to as marking the

site of the alleged bronchial constriction of the gullet. The enlargement of this not uncommon, though usually slight, ledge so as to intercept large, fairly undivided portions of ingesta—for example, large portions of potato—is very rarely observed, but in time a recess may form in which solid portions of food get arrested, pouching follows, and slight dysphagia is observed; at least, that is what apparently happened in Atkinson's patient. Here, again, cutting endo-oesophageal surgery is not advisable, but the passage of large bougies might possibly be useful, or else stretching by Brüning's dilator.

To render this review comprehensive I must not omit to mention that a dissecting diverticulum has been recorded passing down between the coats of the gullet. This condition probably resulted from ulceration down to the muscular coat, and subsequent separation of the coats by pulsion.

Another rare form is a narrow diverticular sinus of varying length and unknown origin, though traumatism, or ulceration, may be suspected. I had an instance of this lesion an inch and a half long, which passed backwards, downwards, and outwards near to and parallel with the diaphragm, *post mortem* in a case of phreno-cardiac stenosis.

To sum up: diverticula of the gullet, in contrast to diverticula in the pharynx, are rarely suspected during life, rarely cause obvious symptoms, and are rarely susceptible of or influenced by any treatment, even should they accidentally be discovered during life. We hear very little of them, and for the above very obvious reasons.

FIG. 2.

FIG. 2.—Typical traction diverticulum of the oesophagus.

FIG. 3.

FIG. 3.—Interior of the same oesophagus showing the mouth of the diverticulum situated well below the level of the bronchial glands.

I desire to acknowledge my indebtedness to Dr. Irwin Moore for the use of the original drawings from which the figures illustrating this paper have been reproduced.

BIBLIOGRAPHY.

- For references to the following authors, mentioned or consulted—namely, Rokittansky, Monro, Ludlow, Bell, Worthington, Ogle, Heschl, Klebs, Kunge, and Nicoladoni—see the article by Zenker and von Ziemssen in the latter's *Cyclopaedia of Practical Medicine*, English translation, 1878. For references to Kocher, von Bergman, Girard, Brewer, Rosenthal, and many others, see Stetten, *Annals of Surgery*, 1910.
- Back and Colledge: *Lancet*, 1923.
 Bevan: *Journ. Amer. Med. Assoc.*, 1921.
 Butlin: *Med.-Chir. Trans.*, 1893; *BRITISH MEDICAL JOURNAL*, 1898 and 1903.
 Davis, E. D.: *Proc. Roy. Soc. Med., Laryng. Sec.*, 1917.
 Dundas-Grant: *Proc. Roy. Soc. Med., Laryng. Sec.*, 1919.
 Godlee and Bucknill: *Lancet*, 1901. Many references to branchial diverticular sinuses and aerocoeles.
 Goldmann: *Zeit. f. Chir.*, 1907; *Beit. Klin. Chir.*, 1909.
 Halstead: *Annals of Surgery*, 1904, and *Surgical Clinics*, Chicago, 1919.
 Hill, W.: *Proc. Roy. Soc. Med., Laryng. Sec.*, November, 1917, and March, 1919.
 Hurst: *Quart. Journ. Med.*, 1914, and *Guy's Hosp. Reports*, 1925-26. The latter gives references to rare true pulsion pouches of the gullet by Kraus, Bensaude, and others.
 Jackson, Chevalier: *Peroral Endoscopy*, 1914; and *Annals of Surgery*, 1926.
 Keith: *BRITISH MEDICAL JOURNAL*, 1910.
 Kelson: *Proc. Roy. Soc. Med., Laryng. Sec.*, 1919.
 Killian: *Ann. d. Mal. de l'Oreille*, 1908; *Zeit. f. Ohrenheilk.*, 1908.
 Knott: *Med. Press and Circ.*, 1900.
 König: *Arch. f. Klin. Chir.*, 1922; *Deut. med. Woch.*, 1922.
 Kostanecki: *Arch. f. Path. Anat.*, 1890.
 Mayo, Charles: *Annals of Surgery*, 1923, and various reports from the Mayo Clinic.
 Morley, John: *BRITISH MEDICAL JOURNAL*, June, 1926.
 Negus: *Journ. of Laryngol.*, 1925.
 Shallow (with Jackson): *Annals of Surgery*, 1926.
 Starck: *Die divertikel der Speiseröhre*. Leipzig, 1900, and 1911.
- Many references.
- Thomson, St. Clair: *Diseases of the Nose and Throat*, 1926.
 Waggett and E. D. Davis: *BRITISH MEDICAL JOURNAL*, 1912.
 Watson: *Journ. Anat. and Physiol.*, 1875.
 Wheeler: *Dublin Med. Journ.*, 1886.
 Zesas: *Deut. Zeit. f. Chir.*, 1906; for references to Niehans, Burckhardt, and 50 other cases.

LARGE PHARYNGEAL DIVERTICULA.

BY

E. I. SPRIGGS, M.D., F.R.C.P.,

Senior Physician, Ruthin Castle, North Wales.

THE following details relate to two patients with large pharyngeal pouches of long standing which had not been operated upon. The first had been present for over forty years.

CASE I.

The patient was a man, aged 76, who sought admission to Ruthin Castle early in 1924, for an affection of one hip-joint, stating that he was otherwise in good health. When giving his history he said that a pressure pouch of the oesophagus had been diagnosed by Sir (then Mr.) Henry Butlin, thirty-eight years before. He had managed the pouch for the whole of this time, the only serious disadvantage being that occasionally at a meal it might discharge involuntarily through the nose. This accident was rare and often did not occur for several months, but the fear of it is present when he is obliged to dine in public rooms in hotels or on a ship. He had lived a very active life, travelling, riding to hounds, and walking long distances, even recently as much as twenty-five miles a day.

With a barium meal a large pouch of clear outline can be seen extending into the thorax as low as the fourth vertebra and below the level of the aortic arch (Fig. 1). It is a little more to the right side than the left.

The procedure which the patient has evolved for managing the pouch is as follows: he retires after each meal, bends forward and makes an effort to increase intrathoracic pressure, putting the closed hand, the back of the first phalanges, against the left side of the neck above the clavicle, and ejects part of the contents of the pouch through the mouth. He then takes a glass of water, evacuates again, and is comfortable and safe from disturbance.

Since the lower part of the pouch is not mainly on the left, it appears that the placing of the back of the hand on that side probably supports the upper part of the pouch or protects a weak place, and this aids in raising thoracic pressure without discomfort. When this procedure was carried out with a barium and buttermilk meal, the first evacuation got rid of three-quarters to four-fifths of the contents of the diverticulum (Fig. 2).

With subsequent meals the glass of water was taken as usual, but some barium returned at the end of the day. The patient believes, however, that with ordinary food there is complete emptying after each meal. In a recent letter to my colleague Dr. Anderson this patient states, for the benefit of anyone in a similar plight, the result of his forty years' experience as follows:

1. All food left in the pouch must be expelled within a few minutes after a meal and the throat gargled with cold water.
2. The saliva should be expelled also before getting into bed.
3. Alcohol must be regarded as a poison; "I allow myself one glass of Madeira to my dinner, however."
4. Beef should be avoided unless minced.
5. Devonshire cream is recommended as a good lubricant which is well borne.

CASE II.

In this patient, a man aged 70, also not operated upon, the symptoms dated back sixteen years. It was first noticed that food began to be arrested on swallowing, and that a piece of meat would return. Twelve years later—that is, four years ago—the symptoms became worse and weight was lost. At the time of the first consultation twenty months ago he could take only soft food, mainly cereal, with milk, milk pudding, butter, bacon fat, and some fruit pulp. Food tended to come back and had to be swallowed again or rejected. Fits of choking occurred, when food might return through the nose. The pouch could be emptied voluntarily. On one occasion this was done in my presence. About half a pint of brown fluid was produced, with the odour of coffee; some fat was floating on the surface. The fluid was acid to litmus, but contained no free acid.

On x-ray examination with a barium meal a large pouch was seen which, in an antero-posterior view, extended into the thorax well beneath the level of the clavicles as low as the fourth dorsal vertebra, most of it being to the right side of the middle line. It was irregular in outline. In the lateral view it can be seen to come below the level of the aortic arch. The mouth of the diverticulum appeared to arise from the posterior part of the pharynx just above the junction of the gullet. It was found on the x-ray table that food passed into the oesophagus more easily in the prone position (Figs. 3, 4, and 5).

This patient was recommended to take food lying forward on the right side. A year later he was holding his own; the weight was the same and he had little discomfort. He found it better not to empty the pouch during the day, as if it were kept full food would pass over it into the stomach. It was then emptied each evening after the last meal.

These were regarded as pressure pouches. Such diverticula have been called congenital, just as have the diverticula arising from the second part of the duodenum, and with no more evidence. They both arise in adult life at a part of the alimentary tube where there are frequent rises of pressure from the contraction of the muscular tube above and where the tube is weak from the divergence of muscular fibres. There may be a sphincteric contraction below, as described by Wilkie and Hartley.¹ When the pharyngeal pouch is once formed we may add to these factors a definite obstruction below; for the neck of the filled pouch will pull upon the gullet at its origin (Fig. 5, lateral view). During swallowing the lower part of the pharynx then becomes a closed cavity, the pressure in which will distend the diverticulum more and more.

In neither case was there any unpleasantness or putrefaction in the contents of the pouch.

Both of these men would be much better off had the pouch been successfully removed. In the first case operation was refused by the patient many years ago. His last consultation with Mr. Butlin was about 1893. At that time he was told that the mortality of the operation was about 50 per cent. It must be admitted that he has done well by his own methods. He does not suffer, casual acquaintances have no idea of the disability, and he has lived, and still lives, an active interesting life.

In the second case failure of nutrition set in definitely four years ago, but on considering the patient's age and physique, two surgeons advised independently against operation. By experiment and observation, including the finding out in the x-ray room in what position food entered the oesophagus most easily, the patient (who is a retired medical man) and his advisers have evolved simple means of diet and living which have enabled him to hold his own.

The photographs shown are by Mr. O. A. Marxer.

REFERENCE.

- ¹ D. P. D. Wilkie and J. N. J. Hartley: *Brit. Journ. Surg.*, 1922, x, 83.

DISCUSSION.

Dr. IRWIN MOORE (London) agreed with Dr. William Hill that such diverticula should not be described as oesophageal. The site at which a pressure pouch originated was in every case so definite and constant—namely, the posterior wall of the deep pharynx, between the oblique and transverse fibres of the crico-pharyngeus muscle—that there was no reason whatever why such pouches should be described as other than pharyngeal. Again, frequent endoscopic examinations had confirmed the fact that the slit-like opening of the lower pharyngeal orifice was always situated at a slightly lower level than the mouth of the pouch, which was definite proof that such pouches were pharyngeal in origin. Their lateral relationship to the oesophagus was probably the reason why they were frequently and erroneously described as oesophageal.