rather firm consistence, and red on section. Some of this enlargement might have been due to congestion, but some of it was certainly due to myeloid transformation. On microscopical examination the splenic pulp, not the lymphatic follicular tissue, was enlarged. In Giemsa-stained sections the pulp was found to contain great numbers of eosinophil cells and erythroblasts and a certain number of myelocytes; a few megakaryocytes were also found (in other sections). A blood count at the end of February, 1926, had given 3,416,000 erythrocytes and 5,450 white cells to the cubic millimetre of blood (of the white cells 89 per cent. were polymorphonuclear leucocytes, 5 per cent. were lymphocytes, 4 per cent. were monocytes, and 2 per cent. were eosinophils), and in a blood film taken on March 10th (the day before death) a few myelocytes were seen.

Since myeloid transformation (metaplasia) was first discovered by Dominici it has probably been more often observed in the spleen than in any other viscus. O. Naegeli, who gives an excellent review of the subject,⁴ says that though Itami failed to produce myeloid metaplasia by experimental bleeding, Skornjakoff succeeded in obtaining this vital reaction by allowing sufficient intervals for recovery between the bleedings. Myeloid transformation of the spleen may (though of course it does not always) occur as a vital reaction to anaemias and cachexias of various kinds and various infective and toxic agents,⁵ including even mercury perchloride.⁶ It has likewise been observed as a compensatory reaction in cases in which the bone marrow has been reduced by osteo-sclerosis.

In conclusion, I must thank Dr. O. Bode (house-physician) for help in the clinical examination of the case, and Dr. E. Bock, and likewise Dr. J. W. McNee, for help in the pathological and histological examination.

REFERENCES.

- ¹ H. Oertel: *Journ. Med. Research*, Boston, 1904, xii, p. 75; and *Journ. Exper. Med.*, New York, 1906, viii, p. 103.
- ² F. Parkes Weber: *Proc. Roy. Soc. Med.*, Section of Pathology, 1909, ii, p. 109 (with references to Oertel's and Curschmann's cases).
- ³ A. Roccavilla: Fall von herdförmiger biliärer Lebernekrose, *Virchow's Archiv*, 1912, ccviii, pp. 421-425.
- ⁴ O. Naegeli: *Blutkrankheiten und Blutdiagnostik*, fourth edition, Berlin, 1923, p. 200-204. Cf. also H. Fischer's excellent monograph, *Myeloische Metaplasie*, Berlin, 1909.
- ⁵ J. W. McLeod and J. W. McNee: On the Anaemia Produced by the Injection of the Haemolysin obtained from *Streptococci*, *Journ. Path. and Bact.*, Cambridge, 1913, xvii, pp. 524-537. Fanny Albrecht: Zur Entstehung der myeloiden Metaplasie bei experimentellen Blutgiftanämien, *Frankfurter Zeit. f. Path.*, Wiesbaden, 1913, xii, pp. 239-265.
- ⁶ Moulinier: Transformation myéloïde complète de la rate dans une intoxication subaiguë par le bichlorure de mercure, *Arch. de Méd. Navale*, Paris, 1903, lxxx, pp. 347-353.

THREE CASES OF OBSTRUCTION AT THE VESICAL OUTLET AFTER PROSTATECTOMY.

BY

W. K. IRWIN, M.D., F.R.C.S.E.,

SURGEON TO OUT-PATIENTS, ST. PAUL'S HOSPITAL FOR GENITO-URINARY DISEASES.

A GREAT deal is written about the satisfactory results of prostatectomy, but comparatively little about the unsatisfactory cases. Yet it is by no means uncommon for obstruction at the vesical outlet to recur after a short interval, however carefully and apparently successfully the operation has been performed, and when there has been nothing to suggest malignancy. The following typical cases from my own practice may be of interest.

CASE I.

In October, 1922, a man, aged 62, was sent to me complaining of difficult and frequent micturition of twelve months' duration. He stated that he was passing urine ten or a dozen times a day, and two or three times at night, the first few hours of recumbency being untroubled. His eye and knee reflexes were normal, and nothing unusual could be palpated in the renal and vesical areas. His urine was free from blood and from any sign of infection, but the residue left in the bladder after micturition amounted to 4½ ounces. On rectal examination the prostate was of uniform consistency, though not so elastic as usual, and the right lobe was definitely enlarged.

I did the ordinary (Freyer) suprapubic prostatectomy, but

experienced some difficulty in enucleating the enlargement, which was more fibrous than usual. I was particularly careful to divide any shelf existing between the prostatic and vesical cavities, and no abnormal symptoms showed themselves during convalescence. The suprapubic wound was completely closed and the patient was passing water by the twenty-second day. He stated that he had to strain more or less to get the urine away, but bougies and catheters passed readily into the bladder.

Five months later he returned complaining of great difficulty and considerable pollakiuria, so much so that he asked for something further to be done. Instruments would not pass owing to definite obstruction at the internal meatus.

I excised the scar tissue, opened the bladder suprapubically and freed it anteriorly, passed a metal bougie by the urethra into the prostatic cavity, put the patient in the Trendelenburg position, and inserted a Thomson-Walker retractor. I found that the internal meatus was greatly contracted, so, while my assistant pushed up the metal bougie, I cut away the intervening tissue with scissors until the instrument entered the bladder. The bougie was then removed, and I completely dissected away the partition between the vesical and prostatic cavities, leaving a large internal meatus. I then passed a big coude catheter along the urethra, fastened it into the bladder and closed the vesical and abdominal wounds around a small suprapubic tube, which was removed three days later.

The bladder was irrigated with boric lotion and the catheter changed daily for a week, the urethra being irrigated as the instrument was withdrawn. On the eighth, ninth, and tenth days I passed a catheter in order to irrigate the bladder, but did not tie it in. On the eleventh day, when the urethra had had three days' rest, I again fastened a catheter in and continued to do so till the seventeenth day, when the wound was firmly healed and the patient passing urine more freely than for years previously.

I have seen him from time to time since, and introduced urethral instruments with a view to ascertaining whether there was any contraction at the bladder neck. Large coude catheters pass without meeting any obstruction, and micturition is free.

CASE II.

In November, 1924, I was asked to see a man, aged 65, suffering from retention, who gave a history of slight difficulty for about a year and great difficulty for two or three months. By rectal examination both prostatic lobes were moderately enlarged but of uniform consistency. I passed a catheter and emptied the bladder gradually.

I performed a prostatectomy by the Freyer method as in the former case, the enlargement being hard, fibrous, and difficult to remove. Great care was taken to divide any shelf between the prostatic and vesical cavities, and the patient convalesced normally for some time, passing urine by the urethra at stool on the eleventh day. Three days later, and many times afterwards, he micturated independently of defaecation.

On the twenty-second day I found that I could not pass a catheter into the bladder, the point being arrested in the prostatic cavity. Only a small quantity of fluid could be introduced and that very slowly. I tried again on many subsequent occasions, always with the same result. Five weeks after the operation, when I was out of town, the suprapubic wound closed and the patient developed retention. The medical man who was called to see him passed a catheter. He thought the instrument was in the bladder when it was really only in the prostatic cavity, a mistake easily made in these cases. As he could not draw off any urine, he came to the conclusion that he had to deal with a case of suppression, and actually applied the treatment for that condition. After a short interval, however, the suprapubic wound burst open again and discharged a large quantity of urine.

I explored the vesical neck in the way already described, but after a thorough examination failed to find any opening. I completely removed the partition between the prostatic and vesical cavities, inserted a catheter into the bladder by the urethra, and treated the vesical and abdominal wounds as described above. The technique of the after-treatment was the same as in the former case. The patient made a rapid and uninterrupted recovery and has since passed urine without any difficulty.

CASE III.

A man, aged 50, who had been leading a healthy country life, consulted me in April, 1925. He looked very fit, but complained of slight difficulty for four or five years and great difficulty for a month. He was passing small quantities of urine every hour both by day and by night. He found it very hard to begin the act, and the stream was a mere dribble. There was no haematuria, no sign of infection, and no history of gonorrhoea. The eye and knee reflexes were normal. On rectal examination the right lobe of the prostate was definitely enlarged, the left little, if at all. I passed a catheter, after meeting some obstruction in the prostatic region and drew off 6 ounces of residual urine.

I performed prostatectomy by the open method, as recommended by Sir John Thomson-Walker, and again found great difficulty

in enucleating the enlargement, which the pathologist described as a fibroadenoma without any trace of malignancy. I took great care to remove all the shelf between the vesical and prostatic cavities; in fact I took away so much that my assistant was afraid I might open into the rectum.

For a time the patient did very well, and three or four weeks later, when the wound was healed, was passing water without the least trouble. Two months after the operation he began to experience some difficulty, more especially after a tiring day's work. I saw him in August, when he was complaining of great difficulty, the stream being small and intermittent. On passing instruments I encountered some obstruction at the internal meatus, though I managed to get steel bougies through. The dilatation gave him relief for some time, but he returned in February, 1926, when the suprapubic wound had broken down and formed a fistula. He was passing some urine normally with great difficulty, but most of it was coming by the suprapubic route.

I operated again on February 23rd and found a pin-point opening surrounded by dense scar tissue. I removed all the obstructing partition, made a large internal meatus, tied in a big catheter by the urethra and drained the bladder suprapubically. In three weeks the abdominal wound was healed and the patient passing urine without the least difficulty; in fact he does not remember ever micturating so easily and quickly before.

The points of special interest in these cases are:

1. All three patients complained of difficulty as their most prominent symptom before the prostatectomy.
2. The enlargement was in each case of the smaller and more fibrous type, a fact which supports the usually held theory that post-operative obstruction is commoner in this class of case.
3. The rapidity with which the wounds healed and the ease with which the patients subsequently passed water lead me to believe that in dealing with post-operative obstruction it is advisable to tie a catheter into the bladder by the urethra off and on until the suprapubic wound is healed. I have not found that the patient objects to its insertion and retention, if the instrument is changed and the urethra irrigated daily, and a good lubricant used.
4. While there is no doubt that the Thomson-Walker open operation is in the majority of cases greatly to be preferred to the old "blind" method, Case III proves that cicatricial contraction may cause obstruction and difficult micturition even after the more thorough procedure.

STUDY OF AGGLUTINATION OF THE GONOCOCCUS IN MAN.

BY

C. E. JENKINS, M.R.C.S., L.R.C.P.,
PATHOLOGIST TO SALFORD ROYAL HOSPITAL.

MUCH work has been done upon the agglutination of the gonococcus by means of the experimental inoculation of animals, and all workers appear to be unanimous that agglutinins are produced; the titre of such serums is high and there is general agreement that a titre of 1 in 1,600 is relatively easy to obtain.

It is on the subject of differentiation of type that divergence of opinion becomes evident. Some workers have been able to differentiate a number of types and subtypes; the precise number varies. Other workers have concluded that it is not possible to distinguish these types with any certainty by means of the agglutination test. A review of the literature has been given by Tulloch.¹

An attempt to investigate this matter in the human subject presented an initial difficulty in that failure to obtain a positive result might possibly be due to the fact that the patient was infected with a type or strain of gonococcus not present in the polyvalent stock antigen. The only practicable method of overcoming this obstacle was to discard the idea of using a stock antigen and instead to use for each patient his own autogenous strain.

The following technique was therefore employed. Upon a patient presenting himself with a newly contracted urethral discharge a culture was made upon the medium always used in this laboratory and described by me in the *Journal of Pathology and Bacteriology* at various times. Next morning the growth was examined to ascertain that pure culture was possible, and subcultures made. These were incubated for twenty-four hours, examined

for contaminations, and, if satisfactory, made into a suspension of about 2,000 million per cubic centimetre with normal saline. The suspension was then violently shaken for a few minutes to break up any clumps present. The further steps in the preparation varied.

Four methods of treating it were tested. In the first neither antiseptic nor heat was used, the suspension being placed in the ice-chest immediately after shaking. The low temperature is quite sufficient to kill the organism. The second method was to heat at 55° C. for thirty minutes, then place in the chest. The third was to heat as before and afterwards add 0.1 per cent. of formaldehyde before storing in the ice-chest. Lastly, 0.1 per cent. of formaldehyde was added without preliminary heating. Immediately before the actual test was performed a sufficient quantity of antigen was taken from the bulk stock and passed through filter paper or a plug of wool in order to ensure a smooth suspension.

In the case of gonococcal urethritis a period of three or four weeks was allowed to elapse before taking a sample of blood. The purpose of the interval was to allow the patient an opportunity to develop any complications usual to that period—orchitis and so forth. It was thought that a more generalized infection would favour the production of agglutinins.

The test was performed the day after the blood was obtained, which meant that the antigen was then about twenty-four days old. The dilutions ranged from 1 in 10 to 1 in 320. The tubes were incubated at 55° C. for two hours, then put out, on the bench at room temperature for another twenty-four hours. They were inspected at frequent intervals. The antigen and serum were stored and the test repeated at the end of a fortnight and again when the serum was six weeks old. This last procedure was to ascertain if any change in the serum took place with age.

The total number of cases examined was 21; they can be conveniently divided into two groups of 16 and 5 respectively.

The first group comprised the patients in whom the organism had been obtained from the urethra. Most of them had developed posterior urethritis, and four of them testicular lesions also, by the time their serum was collected. The result of the tests showed that the serum of every patient in this group was completely negative in all dilutions. Such a result has been obtained by other workers and has been explained on the assumption that agglutinins are not formed where the infection is localized or semi-localized.

The results obtained in the second group are therefore interesting. This group consisted of five cases of gonococcal arthritis. In each case the organism was obtained in pure culture from the joint fluid. Primary cultures are not difficult to obtain provided the fluid is inoculated as soon as it is withdrawn and not left to lie about the ward for an hour or so. The subcultures were made the next morning when the blood was collected and the serum separated. Immediately the antigen was ready for use the test was performed, and repeated at the same intervals as in the first group. The dilutions used were the same. Four of the serums were negative in all dilutions; the fifth was positive in a dilution of 1 in 10. This particular serum was tested again the next day for confirmation. Fourteen days later it was still positive, but was negative when tested at the age of six weeks.

At the present moment an original method of treating the serum is being tried and appears to offer better results in the second group, but the method is still under observation.

CONCLUSIONS.

1. Agglutinins to the gonococcus are not produced in man when the infection is limited to the genito-urinary system.

2. In a generalized infection such as arthritis the amount of agglutinin produced is so small and uncertain that the agglutination test as ordinarily performed is of no value for diagnostic purposes.

REFERENCE.

¹ Tulloch, W. J.: *Journ. Path. and Bact.*, vol. xxv, p. 353.