A STUDY OF EPHEDRINE.

BY

K. K. CHEN, PH.D.

(From the Laboratory of Pharmacology of Peking Union Medical College, Peking, China.)

MA HUANG has been identified as *Ephedra vulgaris*, var. *helvetica*. It is a low, dioecious, practically leafless shrub, 60 to 90 cm. high. The stem, which is green, ribbed, and channelled, is the part sold in Chinese drug stores. It is usually cut into pieces, 1 to 1.5 cm. long. Ephedrine, a natural secondary amine, is the physiologically active constituent and can be easily isolated from the plant by immiscible solvents. It was first discovered by Nagai in 1889, and its structural formula has been studied by different observers.

The Action of Ephedrine.

Our preliminary reports have shown that ephedrine possesses sympathomimetic actions and that its outstanding feature is its circulatory stimulation.

(a) *On Circulation.*—An intravenous injection of ephedrine in an anaesthetized dog or cat always results in rise in blood pressure and increase in pulse rate for fifteen or more minutes. The site of action is not in the central nervous system, for destruction of brain and cord does not prevent the rise in pressure. The acceleration of pulse is not due to depression of the parasympathetic system, for atropinized animals respond just as well as non-atropinized, and, in fact, the pulse is usually decreased in non-anaesthetized animals. Unlike adrenaline, the rise in pressure is not due to peripheral constriction, for intestinal and kidney volumes are often passively increased. Application of the drug directly to the stellate ganglia causes a marked increase in rate and strength of cardiac contractions, but only little rise in blood pressure. When ephedrine has produced maximal cardiac stimulation by local application to the stellate ganglia, or when the latter are paralysed by intravenous injection of nicotine, ephedrine given intravenously still elevates the blood pressure in the usual manner. It appears clear that the rise in blood pressure is mainly due to the stimulation of structures peripheral to the sympathetic accelerator ganglia—myoneural junction or the muscle itself. Since ephedrine acts on the nervous structures of other muscular organs it is probable that it does the same thing here.

(b) *On Smooth Muscles.*—The action of ephedrine on smooth muscles follows very closely the sympathetic innervation. Thus the pupils are dilated by the application of ephedrine to the conjunctival sac. The light reflex is retained and physostigmine causes prompt miosis. Bronchial spasm due to physostigmine is rapidly relieved by ephedrine. The isolated gut is inhibited and relaxed, the vessels of the perfused kidney show definite and prolonged constriction, and strips of rabbit's uterus suspended in Tyrode's or Locke's solution undergo one or more powerful contractions when ephedrine is added. The site of action is apparently only concerned with the terminal nervous structures, for direct application of ephedrine on respective ganglia, or nicotization to paralyse the latter, does not alter the subsequent action of ephedrine.

(c) *On Secretions.*—Ephedrine causes increase in secretion of several organs. In anaesthetized dogs urine and lymph are found to be constantly increased, submaxillary and pancreatic secretion usually augmented, while sweat is not affected, by ephedrine. In non-anaesthetized dogs in which a Pavlov or Heidenhain pouch is made the gastric secretion is found to be definitely increased by subcutaneous injection of ephedrine. Free and total hydrochloric acid rises at the same time. Intestinal juice and bile do not start to flow after ephedrine in dogs with Thiry-Vella and bile fistulae respectively, but pancreatic juice is sometimes increased in those with pancreatic fistulae. In all cases except the submaxillary gland when ephedrine exerts influence the results may be probably referable to the changes in circulation.

(d) *Absorption and Toxicity.*—Circulatory effects are immediately shown by intravenous injection, less rapidly by intramuscular or subcutaneous injection, or by the mouth. Intraperitoneal injection is also effective. The

minimal effective dose in anaesthetized dogs is as small as 0.25 mg. The minimum lethal dose varies in different animals and with different methods of administration. In frogs, when injected into the anterior lymph sac, it lies between 600 and 650 mg. per kilo of body weight. By intravenous injection the M.L.D. in white rats is 140 mg. per kilo of body weight, in rabbits 70 mg. per kilo, in cats 75 mg. per kilo, and in dogs 70 mg. per kilo. By intramuscular or subcutaneous injection rabbits can tolerate a much larger dose than the M.L.D., and by daily repeated intravenous injections they can stand a dose 840 times the M.L.D. without any pathological changes. There is no doubt that this drug has a low toxicity.

Ephedrine in Experimental Shock and Haemorrhage.

In anaesthetized dogs, ephedrine raises blood pressure in haemorrhage and experimental shock induced by histamine, peptone, anaphylaxis, or surgical violence and trauma. The rise in blood pressure is permanent under favourable conditions, and its effect is due to cardiac stimulation and not to arterial constriction. Ephedrine fails to act when the heart becomes impaired, or respiration ceases, or the degree of shock is too extensive, or when haemorrhage exceeds 25 per cent. of the total volume. It has, however, no harmful effects. Clinical use can probably be made in the early stage and in mild forms of shock, or as a prophylactic drug against surgical shock in long operations.

Clinical Uses.

Ephedrine can replace adrenaline in several instances.

(a) *In Hypotension.*—The only case of hypotension studied was that of Addison's disease, in which intramuscular injections of 100 mg. of ephedrine twice daily raised blood pressure from 70/40 mm. to 110/68 mm. Hg for about two hours, and by mouth a single dose of 150 mg. maintained the pressure at a high level for over twenty hours.

(b) *In Asthma.*—As a broncho-dilator ephedrine appears to be very efficient. In five cases treated the results were most convincing; 80 to 150 mg. given intramuscularly never failed to relieve asthmatic attacks completely. The effects come on usually within fifteen minutes at the same time as blood pressure rises. In one case adrenaline was administered before ephedrine, but the patient's dyspnoea did not totally disappear until the latter was used. This is considered a temporary relief, although none of our treated patients have come back with recurrence of their complaint.

The good features of ephedrine can be most conveniently summarized as follows:

1. Ephedrine is chemically stable; age, exposure to light, or boiling does not alter its action.
2. It has a persistent and uniform action, in contrast with that of adrenaline.
3. It has a low toxicity. The margin of safety is unquestionably very wide.
4. It can be given very conveniently, either per os or by intramuscular injection. There is no local irritation.

Memoranda:**MEDICAL, SURGICAL, OBSTETRICAL.****ESSENTIAL HAEMATURIA.**

THE following details of a case of essential haematuria which occurred in Wuchang, Central China, in June, 1926, seem to me to be of sufficient interest to merit publication.

A young and otherwise healthy man, aged 29, with a negative family history for kidney trouble of any kind, had one attack of albuminuria in Egypt in 1917. This was thoroughly investigated; radiological and cystoscopic examinations were made without result, and no tubercle bacilli were detected. The right kidney only was affected, and the albuminuria was thought to be connected with a concurrent attack of appendicitis. The appendix was removed in 1919, and showed signs of previous inflammation.

The first attack of haematuria occurred at the end of June, 1926. The shade temperature was about 90° F., and the patient was working at high pressure, mentally and physically, but feeling fit. He first noticed blood in his urine after a strenuous game of tennis, followed immediately by a cold-bath. The urine remained red that day, but the following midday it was much