or roasted. Potatoes, mashed or in their jackets. Turnips. Spinach. Artichokes. Mild cheese. Fruit after three months.

Forbidden.—Fat and eggs. Stewed and fried foods. Sweetbreads. Liver. Duck. Kidneys. Pork. Black puddings. Game. High or spiced dishes. Oily fish. Soft roe. Herrings. Salmon. Sardines. Cooked butter. Peas. Cabbage. Raw vegetables. Strong cheeses. Aerated waters. Alcohol.

DIET AFTER GASTRO-ENTEROSTOMY.

Guides to Diet:

1. Easily digested food.
2. Large meals should never be taken.
3. Preferably there should be no drinking during meals.
4. Food and drinks should not be taken hot.
5. Eat slowly, masticate thoroughly, avoid bulky meals.
6. One glass of milk between meals and one before retiring.

Breakfast.—Carefully made porridge. Egg—lightly boiled. White fish, boiled or baked. Cocoa or weak tea with plenty of milk. Cold dry toast and margarine or butter. Cold boiled ham.

Other Meals.—Oysters. A little roast beef, well cooked, mutton or veal. Clear gravy. Boiled beef. Chicken. Tripe. Spinach. Cauliflower. Mashed potatoes. Macaroni. Milk puddings. Junket. Blanc-mange. Custard.

Forbidden.—Fried fish in fat. Smoked fish. Bacon. Sausage. Meat pies. Meat puddings. Stews. Made-up dishes. Re-cooked meats. Cabbage. Brussels sprouts. Carrots. Turnips. Parsnips. Watercress. Fried potatoes. Pepper. Mustard. Vinegar. Pastry. Cake. Brown bread. Jam. Sweets. Cheese. Coffee. Alcohol.

My investigations have shown that success or failure in the treatment of gastric and duodenal ulcers depends far more on proper after-care than the exact type of operation chosen. I am strongly convinced that the attitude of those surgeons who affirm that they have no use for any operation which does not completely cure the patient is entirely wrong.

A change of air greatly assists the convalescent. He gets away from the environment of his illness. He sees new scenes and new people and gets new interests. Routine and regularity should mark the period. Each day should be mapped out with the proper intervals of rest and

graded exercise, and the amount of mental recreation should be apportioned. While it is difficult to estimate accurately the effects of post-operative respiratory exercises and physical exercises on the Swedish principle, there is no doubt that the general circulation is improved, the bodily functions are more normally performed, muscular weakness is diminished, and the patient returns to complete health more quickly. I hold strongly that systematic exercises should be a routine measure in the convalescent period. The various measures of physiotherapy also hold a high place in conducting to rapid recovery.

For the convalescent period the hospital patient's own home does not, as a rule, provide an environment conducive to rapid recovery, and though many may be sent to convalescent homes there will always be a large proportion of patients for whom it is not possible to provide accommodation in these institutions. I strongly believe that there should be a convalescent department attached to each hospital. In this way we should eliminate the neglected period that now prevails between the completion of active surgical treatment and the time of return to full occupation. To this department there could be attached a social service, for it has been estimated that 30 per cent. of patients discharged from hospital require help in this direction. The medical staff in charge should be specially trained in convalescence. Here patients would be completely investigated and graded; their requirements as to diet, discipline, and physical and mental recreation would be determined, and any disabilities that arose or whose presence had been overlooked could be dealt with and the malingerer eliminated. Advantage could be taken in many instances to improve the general health.

THE RESULTS OF INSULIN THERAPY IN DIABETES MELLITUS.*

BY

HUGH MACLEAN, M.D., D.Sc., F.R.C.P.,
Professor of Medicine, St. Thomas's Hospital, University of London.

ABOUT four years have elapsed since insulin became generally available in this country for the treatment of diabetes mellitus. This is not a very long period in which to judge the ultimate effect of insulin on diabetes, but it is sufficient to enable us to come to some definite conclusion as to its more immediate and direct influence on the disease. The theoretical considerations which govern the action of insulin in diabetes are quite clear. The oxidation of our food-stuffs requires the presence of a substance which is normally supplied by the pancreas. When, for some reason, the specialized pancreatic islet cells fail to supply this substance in the necessary amount, an interference with the burning up of sugar and fat results. This is the basis of diabetes mellitus. Since it is now possible to supply this essential substance in the form of insulin, it might be thought that the treatment of diabetes should no longer present any difficulties, and that all that is necessary is to supply sufficient insulin to make up for the deficiency in any particular patient.

The difficulty, however, is to supply the exact amount of insulin necessary for the varying requirements of the organism. In the body there is a mechanism which regulates the production of insulin in accordance with the tissue needs at any given time. The amount of insulin required by the body is dependent on many factors, and varies under different conditions. The artificial use of insulin by injection is attended by the drawback that we cannot regulate the amount required to supply the exact demands at any given time. It is true that in some patients we can get fairly close to the theoretical needs, but this can only be accomplished by paying great attention to

various factors, such as diet and exercise. Often it is impossible to regulate the artificial supply of insulin so as to simulate the normal mechanism. Though we can now supply the necessary insulin when the body can no longer secrete it, we can, so far, do but little to reproduce the results of the normal automatic mechanism which regulates the production of insulin according to the prevailing needs at the moment. This is one of the main difficulties in insulin therapy, but under ordinary conditions this drawback can generally be more or less overcome. It is when diabetes becomes complicated by various infections that our troubles really assume a serious aspect. In certain of these conditions insulin seems to lose its normal activity, and, in spite of greatly increased dosage, the patient may sometimes succumb. The healthy individual when suffering from a similar infection usually shows no disturbance of metabolism such as is seen in diabetes; it must, therefore, be assumed that insulin, when supplied naturally by the body tissues, is more effective in these conditions than it is when given artificially by injection. Indeed, in certain cases of diabetes complicated by severe infection, insulin artificially administered proves of little or no benefit.

Before considering the effects of insulin in diabetes it will be necessary to consider the results obtained by diet in the days before insulin was available.

Results of Dietetic Treatment of Diabetes.

For some years before the isolation of insulin much attention was paid to the dietetic treatment of diabetes, and many claims were made on behalf of this method. While it was generally agreed that the only means of influencing the disease in any way was by means of properly regulated diet, the outlook for the average patient suffering from the disease was gloomy in the extreme. Generally, it was found that suitable diet had a wonderful effect to begin with, but, in spite of the greatest care, the patient after a time gradually began to show his old symptoms, and soon a point was reached at which the strictest dietetic measures no longer prevented the occurrence of severe glycosuria and ketosis. From this point the patient usually went steadily downhill, and soon coma or some intermittent infection terminated the scene. The duration of life under dietetic treatment depended, of course, on the

* A paper opening a discussion in the Section of Medicine at the Annual Meeting of the British Medical Association, Edinburgh, 1927.

severity of the lesion, but, unfortunately, as it is generally seen in young people, the disease is a steadily progressive one, and after a time its course cannot usually be stayed to any appreciable extent by diet.

No doubt so-called diabetes in elderly subjects often presented an entirely different picture, and frequently responded well to diet. In such patients, however, glycosuria is the main feature, and there is generally little or no ketosis. In many of them no evidence of disturbed oxidation can be obtained, and the patient may suffer little or no inconvenience. This type of case is really not a true diabetes, but a simple glycosuria which is probably dependent on an inability on the part of the liver and muscles to store sugar in the normal manner. Consequently hyperglycaemia results, and sugar is passed in the urine. The usual symptoms of diabetes are absent, or, if present at all, are of a very mild nature. These cases should not be classified as true diabetes; there is no tendency to coma, and the glycosuria can often be controlled by dietetic measures. Not infrequently the sugar is detected as the result of a routine examination of the urine. Often the glycosuria does no harm whatever to the patient, and though the condition may sometimes develop into a true diabetes, yet, in my experience, this is a comparatively rare sequel. Such cases do not come under the heading of true diabetes.

In considering the results of dietetic treatment it is essential to remember the difference between these two types, for the ultimate effects of diet depend almost entirely on the nature of the cases treated. Elderly patients suffering from glycosuria with little or no ketosis frequently do exceedingly well on diet. Patients suffering from true diabetes ultimately do very badly, though the results may be satisfactory for a short time. The success or failure of dietetic treatment, as indicated by the experience of any particular observer, depends very largely on the number of cases of ordinary glycosuria he includes under the term "diabetes." If many of the patients were suffering from nothing more serious than the well known glycosuria of the elderly individual, then dietetic treatment may appear to give excellent results. When, on the contrary, only cases of true diabetes with the usual symptoms are included, the results of diet are absolutely unsatisfactory in the great majority of patients.

When considering the effects of insulin, it is really with diabetes as it shows itself in its typical form that we have to deal. Usually diabetes is a disease of youth or of middle age, though not infrequently it may be seen in old age. On the whole, true diabetes is more common in patients under 45 years of age, while glycosuria due to lack of storage generally occurs after this age. In arriving at a definite conclusion as to the effects of insulin in diabetes it is necessary to bear in mind that it is its beneficial effect on true diabetes that is really important. In the other class of case coma does not intervene, and death is usually due, not to the effects of the glycosuria, but to some other disease.

Before the days of insulin many observers claimed excellent results from dietetic treatment. Others frankly confessed that diet, however carefully followed by the patient, was quite useless after a short time. Several years ago I carefully examined my figures for patients suffering from typical diabetes who were treated by dietetic measures. The results showed that death took place on an average in about five years or rather less after the dietetic treatment began. I have had communications with various authorities in other countries on this subject, and the information I obtained was to the effect that the average duration of life in patients on dietetic treatment was from four to six years. These cases, of course, did not include patients suffering from glycosuria without ketosis. We may, therefore, take the view that any patient suffering from diabetes in the true sense of the term is unlikely to survive for more than five or six years on an average when the treatment is restricted to dietetic measures. Indeed, in severe cases, two to three years is often all that may be reasonably expected. Consideration of these figures will enable us to appreciate the changes brought about by insulin therapy.

The Results of Insulin Therapy.

In the four years that have elapsed since insulin became generally available for the treatment of diabetes 291 patients suffering from typical diabetes have been treated in a special clinic in the Medical Unit of St. Thomas's Hospital. All these patients showed definite ketosis, and the majority of them were young or middle-aged. Cases of ordinary glycosuria without ketosis are not included in this list. Each patient was first treated as an in-patient in the ward, the diet and insulin dosage being carefully worked out. After being discharged from the ward the patient attended the clinic at definite intervals. As an outside patient he was usually being looked after by his medical practitioner. When any change in diet or in insulin dosage was considered advisable, the practitioner was informed of the result of the examination, and the suggested changes indicated to him. The complete treatment of the case, however, was, as far as possible, controlled by the special clinic. The following figures show the results of insulin treatment of these cases up to date. In this connexion I should like to thank Dr. Crozier of the medical unit for his help in keeping records of these cases and for assisting me in compiling the results.

Year of admission of patient.	No. of patients admitted.	No. of deaths to date.
1923-24	77	5
1924-25	76	2
1925-26	68	3
1926-27	70	2
Total	291	12

Several of these patients were admitted in coma; three of them developed coma on three different occasions and a few more were in coma on two occasions. All these patients are now doing well. Two patients with tubercle bacilli in their sputum are still alive and in fairly good health.

Of the twelve deaths, one, a girl of 19, developed coma outside, and was treated by her local doctor. Another was treated in hospital, but was admitted at a rather late stage, and though apparently a case of uncomplicated diabetic coma, she failed to respond to insulin. Three of these cases recovered from coma, but died of cardiac failure. In all the others coma developed as the result of some such complication as influenza, pneumonia, or septic lesion.

These figures show a death rate of almost exactly 4 per cent. As they were all well marked cases the figures indicate a very extraordinary improvement over the results of dietetic treatment. Judging from the death rate prior to the use of insulin, the average number of deaths in these 291 patients would have been much greater if treatment had been restricted to diet alone. While it is quite impossible to give anything like exact figures, it is nevertheless possible to arrive at some probable conclusions as to what would have happened if no insulin had been used in these cases.

If we take six years as the average duration of life on diet alone, it appears that of the 77 patients admitted in 1923-24 at least half of them would now be dead. Taking everything into consideration, it seems probable that more than one-third of the total number of these 291 cases would not be alive now. In addition to the above patients, I have seen a fair number regarding whom I have no definite statistics. The general impression conveyed, however, is that the average results of insulin treatment were somewhat similar. So far, I have only seen one diabetic patient who failed entirely to respond to insulin administration. As no *post-mortem* examination took place in this case it is possible that some other complication was present.

These results point to the conclusion that practically all uncomplicated cases of diabetes in young or middle-aged people can be controlled by the judicious use of insulin. It must also be remembered that the general physical condition of diabetic patients treated by insulin is altogether different from that which obtained under dietetic treatment. In many cases dietetic treatment was so severe that the patients could not be induced to continue with it. Some of them quite frankly stated that they preferred death to the physical agony which they

endured. In almost all bad cases the period of a year or so preceding death was usually most distressing and hopeless. Now all is changed. The disease which was formerly so grave has been shorn of many of its terrors, and, in my opinion, it ought to be possible to control any ordinary case of uncomplicated diabetes and to restore the patient to a fair degree of health and comfort.

Some Difficulties encountered in the Use of Insulin.

In uncomplicated cases of diabetes insulin has proved to be a very excellent remedy. With a well balanced diet and suitable dosage of insulin patients usually get on all right and give little trouble as long as infections of various kinds are avoided. When a diabetic patient becomes ill from some acute disease it often happens that the insulin loses its effect to some extent, and a marked tendency to coma sets in. In these circumstances the insulin dose must be very materially increased, perhaps to three or four times what the patient was having previously. Sometimes even much larger doses than this are necessary for a time. The secret of successful treatment of diabetes by insulin lies largely in a recognition of this fact. It cannot be too strongly emphasized that very large doses of insulin may be required when acute infections supervene. Very often, unfortunately, the medical man in attendance on the patient takes the view that some temporary complication, which perhaps makes the patient unable to take his ordinary diet, indicates a decrease in the dose of insulin. Such treatment is often fatal. The dose of insulin should in the great majority of these patients be increased at an early stage of the complication. When this is done in a judicious manner it is my experience that grave consequences can very often be avoided. No doubt the complication itself may sometimes kill, irrespective of the diabetes, but probably the most essential point to bear in mind in the routine treatment of diabetes by insulin is the necessity to give a sufficient amount of insulin when complications supervene.

The Results of Insulin Therapy in Young Children.

It is well recognized that diabetes in the child is a very serious condition, and, in the absence of insulin, is liable to prove fatal in a very short time. On the other hand, the disease is usually well controlled by insulin when a sufficient dose is given. Since nearly all children contract some of the various ailments common to children at some time or other, it is obvious that in the case of very young people there are special difficulties and dangers to contend with. In my experience, however, these dangers are not necessarily so formidable as they might appear to be. Undoubtedly very great care is necessary, and the dose of insulin must at once be increased when the onset of any of these ailments is accompanied by an increase in the diabetic symptoms. When this is carefully done the results on the whole are good. I cannot claim to have seen very many cases of diabetes in young children, but since 1923 I have treated over thirty children varying in age from 2½ to 10 years, and all of these are alive to-day and in quite good health. We have not had a single death from diabetic coma in a child under 10 years of age. Many of these children were first seen while in coma, and a few of them developed coma on at least two subsequent occasions. Some of them have contracted and safely weathered such infections as whooping-cough, measles, and chicken-pox. While it must be admitted that the outlook from the theoretical point of view is not good in the case of children, practical experience suggests that the young diabetic can very often be safely guided through the storms which are peculiar to childhood, and that very satisfactory results can often be obtained.

An interesting question arises in connexion with the possible cure of diabetes by insulin in the young. Definite evidence has been brought forward that regeneration of the islet cells of the pancreas may take place in a young diabetic when the disease is controlled by insulin. From this it might be argued that the administration of insulin in very young people might sometimes effect a cure by allowing new pancreatic cells to develop. Several of my cases now require much smaller doses of insulin than they did at the earlier stages of treatment, but

I have not yet seen a single patient whose recovery of tolerance was sufficiently marked to enable him to discard insulin. In some of these young patients there has been so sign of increased tolerance, and the amount of insulin had sometimes to be increased as the child grew older. It seems fairly certain that all these children will find it necessary to continue the administration of insulin throughout life. It is quite certain that the lives of many children who would otherwise die of diabetic coma can be saved by insulin, and that the outlook is not necessarily as black as it is sometimes considered to be. Given a child otherwise healthy, there is really good ground for the view that such a child may reach adult life, and be no worse off than the average patient in whom diabetes first showed itself at a much later stage in life.

Insulin in Elderly and Old Patients.

In old patients, as already indicated, true progressive diabetes with ketosis is not very common, though it is certainly met with from time to time. By far the most frequent condition in these patients is the type of glycosuria already discussed. In an uncomplicated case of diabetes in an elderly patient insulin appears to act as well as it does in the young or middle-aged, but, unfortunately, it often happens that other ailments are present. In these circumstances good results could hardly be expected in general from the use of insulin in old patients, and this is in accordance with experience. Insulin, however, often gives excellent results in many old people who have glycosuria with such symptoms as weakness, neuritis, and local irritation, but whose condition is otherwise good. It all depends whether the lesion is or is not complicated with degenerative diseases, and other changes incident to old age. There is obviously an absence of the recuperative power of the young individual, and allowances must be made for this.

The Use of Insulin in Surgical Conditions Complicating Diabetes.

Insulin has revolutionized diabetic surgery. Before insulin was available the surgeon hesitated as a rule to incur the grave risk associated with an operation on a diabetic patient. In desperate cases, when an operation had to be attempted, coma often supervened and the patient died. Since insulin has become available for the treatment of these patients it is safe to say that, under good conditions, and when treated by someone who understands the use of insulin, the diabetic has about as good a chance as a normal patient who undergoes an operation. This is well borne out by the statistics of the Mayo Clinic. In 667 operations performed on diabetic patients only 20 deaths occurred. Taking into consideration the fact that many of these patients were suffering from arterial sclerosis and other degenerative lesions the results appear to be very satisfactory.

Some Anomalous Statistical Results.

From the excellent results obtained by the use of insulin it might be thought that diabetes would figure no longer in official statistics as an important cause of death. At any rate, it might be considered that the number of deaths officially returned as caused by diabetes would be much reduced, and that this reduction would continue for several years. Since the majority of diabetic patients develop the disease when comparatively young, the natural inference would be that the lives of these patients would be prolonged for many years by insulin; ultimately, when death took place in the usual course of events, though the patient might die of something quite different from diabetes, there would be a tendency on the part of the medical man to return the cause of death as diabetes. Theoretically, then, there should be a marked diminution in the number of deaths from diabetes for several years, while after a certain period an increase might be looked for, but the majority of these deaths should be those of elderly people.

So far as official statistics go, however, insulin has apparently not decreased the number of deaths from diabetes. According to these returns, insulin has failed to affect to any material extent the death rate from diabetes. We are therefore faced with the curious result

that apparently the number of patients dying from diabetes is as high as ever, in spite of the undisputed fact that many lives are saved by insulin. What is the explanation of this? One possible explanation might be that the incidence of diabetes has largely increased during the last few years, and that, despite the beneficial action of insulin, the increase of the disease is so marked as to keep the death rate up to the old level. This view really means that in the absence of insulin the death rate from diabetes would be very materially higher than it now is. Against this view is the fact that insulin is such a potent measure in the treatment of uncomplicated diabetes that even a marked increase in the incidence of the disease should not materially increase the death rate. There appears to be some evidence pointing to the conclusion that there is a tendency for some slight increase in the number of diabetic cases, but this does not explain the difficulty.

An examination of the age of death reveals the fact that the age incidence has shifted, and that for the last few years the majority of diabetic deaths occurred in elderly people. The number of diabetic deaths in young people has very materially decreased. Now, since true diabetes is comparatively rare in elderly people, while simple glycosuria is exceedingly common, it is but reasonable to suppose that the majority of persons returned as having died from diabetes really suffered from simple glycosuria. Since this type of glycosuria does not as a rule produce coma, it is more likely that the majority of these deaths were caused by some condition other than diabetes. The fact that glycosuria happened to be present resulted in "diabetes" being returned as the cause of death. No doubt the recent introduction of insulin has tended to focus attention on glycosuria, and so its presence is more frequently determined. Glycosuria, when present, is such a definite symptom that naturally it would nearly always be mentioned under the term "diabetes" in the official certificate of death.

Again, it is certain that insulin is often not nearly so effective in the treatment of coma in the elderly as it is in the young. Coma is exceedingly detrimental to the heart, and since the heart muscle in old people frequently suffers from some degree of degeneration, it is not unusual for the patient to recover from coma and to die of cardiac failure.

All these facts may play a part in explaining the anomalous official figures which apparently suggest that insulin has failed to reduce the diabetic death rate. These considerations indicate the marked importance of differentiating between cases of true diabetes and those of simple glycosuria.

Results of Insulin Treatment in Coma.

When coma has actually developed much depends on the quickness and energy of the medical man in attendance. The patient should be treated at the earliest possible moment, for the strain on certain of the organs, such as the heart and kidneys, is very great, and, if prolonged, may result indirectly in death.

If the patient is seen at a fairly early stage and no complications are present, recovery should be the rule. I have not had a single death from coma in a young and otherwise healthy patient since insulin became available. In patients suffering from acute infections recovery is more doubtful, but if sufficient insulin is used a successful result should be frequently obtained. Generally speaking, the young diabetic patient, healthy in other respects, should not die from diabetic coma if insulin is judiciously administered at an early stage. In old people the case is very different, for reasons already discussed. Coma in a vigorous sound patient generally responds to insulin, whatever the age happens to be.

Some Points in the Treatment of Coma.

It is generally agreed that the most important measure in the treatment of coma is to give insulin in sufficient amount at the earliest possible moment. Whether or not this should be combined with the administration of sodium bicarbonate is a matter on which opinion is divided. From some experiments carried out in my laboratory by Dr. Goldblatt, it would appear that the use of alkali, at any

rate in large doses, is definitely contraindicated in diabetic coma. Goldblatt induced a ketosis in himself by starving for forty-eight hours. At this stage the urine contained fairly large amounts of acetone and diacetic acid, as indicated by Rothera's reaction and the ferric chloride test. This ketosis rapidly disappeared after taking some glucose. Careful observation showed that after the ingestion of 50 grams of glucose dissolved in water the ketosis disappeared entirely in almost exactly one hour. When, in addition to the glucose, a large dose of sodium bicarbonate was also taken, the ketosis persisted for a long time, and was still present in definite amounts after eight hours. Many experiments of a similar kind were carried out, both on man and on animals; the results were always the same, and tended to show quite conclusively that sodium bicarbonate administered in the course of an acidosis markedly prolonged the condition. In the light of these experiments there can be little doubt that the ingestion of large doses of alkali during diabetic coma tends to hinder the disappearance of the coma. Sodium bicarbonate should, therefore, not be given when treating coma by insulin, since it does no good and definitely retards recovery.

Insulin Treatment and the Persistence of Glycosuria.

It might be thought that the results of insulin therapy in any given patient would depend on keeping the blood sugar normal and the urine free from sugar. Experience shows quite definitely that this is not always so. While it is good practice in all cases to endeavour to render the patient aglycosuric, yet it may sometimes be impossible to do so without having recourse to very frequent injections of insulin. In my experience patients whose blood sugar was kept within normal limits at all times, and whose urines were constantly free from sugar, did no better than others whose blood sugar could not be always kept low. In the case of a small boy under my care it was quite impossible to control the blood sugar and glycosuria, for this patient had a real genius in the way of procuring all sorts of prohibited food. While in hospital it was almost impossible to regulate his diet, and after going out he was entirely beyond control. His urine nearly always contained a large amount of sugar—sometimes a very large amount—which generally depended on liberal supplies of chocolate which he bribed his young friends to procure for him. He seemed a quite hopeless case, and I must admit that I anticipated disaster, yet in spite of all he has done quite well and is now well developed, vigorous, and healthy. Indeed, the result in the case of this young patient makes one wonder whether certain of our theories connected with the treatment of diabetes are sound. Certainly it is best to keep the urine free from sugar, and the blood sugar as near the normal as possible, but the medical man need not despair if he finds this ideal difficult. Many patients do exceedingly well even when some glycosuria is more or less constantly present.

The Administration of Insulin.

Insulin is usually given twice daily—in the morning before breakfast and at night before the evening meal. In some patients two doses are not sufficient to keep the blood sugar within normal limits, and the question arises whether in such cases we are justified in giving three or more injections. I am quite definitely convinced that the injection of more than two doses usually does more harm than good. Though frequent injections may control the glycosuria, the mental effect is so disturbing that many patients cannot stand it. There is a feeling that they can never get away from injections for any period of time, and they sometimes suffer from a definite nervous condition. I have seen a patient in a hopeless condition of nervous dread as the result of the administration of insulin three times a day; all the symptoms disappeared when the patient was put on two daily injections.

Diet and Insulin.

There is no doubt that the effect of insulin therapy is largely dependent on the balancing of the insulin dosage with the diet. Experience, however, shows that, provided sufficient insulin is taken, there are fairly wide limits of

diet between which the patient can live and thrive. Many of our accepted ideas with regard to diet are theoretical, and do not always work out well in actual practice. When sufficient insulin is given it is quite possible for a patient to get along satisfactorily without weighing his food. Patients who are feeling fit as the result of insulin treatment are usually engaged in some business or manual labour and cannot be expected to carry a balance about with them. Indeed, it is seldom, if ever, necessary to weigh the food constantly if the patient is at all intelligent. All that is required is to teach the patient to observe how much food is represented by a definite weight. After a very little practice he is able to select and gauge his diet with ease. The average patient very soon learns to eat approximately the same amount of food each day without weighing it, and I am beginning to think that the added comfort induced by this method results in the patient being in better condition than is the case when his food is scrupulously weighed out for him. The simpler we make all the arrangements connected with insulin therapy the better results we obtain, provided that the main principles are always kept in view.

On the whole, insulin can, in general, control the symptoms of diabetes in uncomplicated cases. There are drawbacks to its use, but when judiciously administered it usually results in restoring the patient to a satisfactory state of health. This is especially so when insulin is used at an early stage of diabetes. So far no substitute has been discovered which can be taken by the mouth. None of the pancreatic mouth preparations at present on the market appear to have the slightest effect on the disease, in spite of the claims constantly advanced on their behalf. Dr. Fuller, who recently carried out an investigation on this point in my laboratory, came to the conclusion that there was no evidence of any action on carbohydrate metabolism as the result of taking these preparations.

The latest mouth preparation is a synthetic product called "synthalin." There can be no doubt that this substance when taken by the mouth does produce a definite effect in certain cases of diabetes. The glycosuria is lessened and the ketosis is often markedly reduced. The product, however, has marked toxic properties, and tends to produce gastro-intestinal and other disturbances, so that its use in the treatment of diabetes is at present not practical. It is conceivable that some preparation of a somewhat similar constitution may be found which will have a similar action without the toxic effects of synthalin. Meanwhile we must accept the view that, for the present, the only useful substance in the treatment of diabetes is insulin.

STATISTICAL EVIDENCE OF THE VALUE OF INSULIN.*

BY

THE LATE PROFESSOR K. A. PETRÉN,
University of Lund, Sweden.

WHEN we are dealing with a chronic disease we cannot come to a true knowledge of the prognosis merely by observation in the clinic. As regards diabetes, we have all seen in the old days how many patients died from coma, and in this respect we have obtained a certain knowledge of prognosis; but full knowledge can only come from following up in a systematic way a sufficient number of unselected cases. Very few investigations of that kind have been made; I know of only two—one by Horace Gray in America, the other by Malmnos, using the material from my clinic. Malmnos dealt only with patients after at least five years had passed since they were first treated in the clinic. The results of this investigation were bad; 70 per cent. of cases showing a fasting blood sugar value of 0.18 per cent. on admission ended fatally. The interest of these investigations is the greater because the doors to the country where we could get knowledge of this kind are now closed and no human force can open them; they were closed by

* Communicated to the Section of Medicine at the Annual Meeting of the British Medical Association at Edinburgh in a discussion on the results of insulin therapy in diabetes mellitus. News of Professor Petré's recent death reaches us as we go to press with his paper.

the discovery of insulin, and the noble way in which our colleagues in Toronto have given the fruits of the discovery to the whole world.

It will in time be of the greatest interest to compare the corresponding figures of the insulin period, but the time for this is not yet ripe. Professor Maclean has given us his statistics for four years of the insulin epoch, and we must thankfully recognize that the figures speak for a very hopeful view on the prognosis of this disease; only the future, however, can give us the definite answer.

In the figures I am giving here of my cases during the insulin epoch I am for the present dealing only with cases which were admitted to the clinic with a clear picture of coma.

Seventy-three Attacks of Coma in 57 Cases (May, 1923, to July, 1927).

Year.	No. of Cases.	Living.	Fatal Issue.		
			In Coma.		Other Diseases.
			In Hospital.	At Home.	
1923 ...	10	6	—	—	4 (3 tuberculosis)
1924 ...	15	9	1	4	1
1925 ...	10	10	—	—	—
1926 ...	17	14	1	—	2 (sepsis)
1927 ...	5	3	2 (21)	—	—
Total ...	57	42	4 (23)	4	7

Analysing the cases of death in coma in the clinic, we find that in 1924 we had not formulated our present rules for treatment, and the patient who died in that year was treated with too small doses of insulin. The 1926 case and one of the 1927 cases were patients who had refused insulin treatment in the clinic earlier in the illness; in the other 1927 case the necropsy revealed a broncho-pneumonia which the pathologist considered to be antecedent to the coma.

With regard to the treatment of coma, it is my rule to try to bring down the blood sugar to a normal value in a very short time—twelve hours at longest, preferably in six or eight; in my opinion it is unnecessary to wait for the report on the blood sugar before giving a further dose of insulin; if the respiration continues to be increased a fresh dose of insulin may be given after half an hour's interval.

As to diet, I agree with Joslin and give the same diet as in other cases of grave diabetes; consequently I give 1.0 glucose. Campbell in Toronto originally spoke in favour of large intravenous injections of glucose; I am glad to see that in his later publication he has abandoned this treatment and says nothing more about it. The following table gives the dates of the latest information about my cases.

For the 42 living cases the latest information is:

From June, 1927 ...	9 cases
May, 1927 ...	17 "
April, 1927 ...	6 "
March, 1927 ...	4 "
February, 1927 ...	6 "

For comparison I give the corresponding figures of Joslin's cases from his latest publication.

Joslin's Cases: Sixty-three Attacks of Coma in 57 Cases (April, 1923, to February, 1927).

Year.	No. of Cases.	Living.	Fatal Issue.		
			In Coma.		Other Diseases.
			In Hospital.	At Home.	
1923 ...	11	8	2	1	—
1924 ...	18	13	—	—	5
1925 ...	14	12	—	—	2
1926 ...	17	16	—	—	1
1927 ...	3	2	1	—	—
Total ...	57* (63)	45* (51)	3	1	8

* Joslin has not made a distinction between the number of attacks and the number of cases. I have had some difficulty therefore, and I hope I have made no mistake in quoting his figures.