Published in final edited form as:

J Am Chem Soc. 2006 July 26; 128(29): 9268–9269. doi:10.1021/ja061132g.

Reversible O–O bond cleavage in copper-dioxygen isomers: impact of anion basicity

Xavier Ottenwaelder, Deanne Jackson Rudd, Mary C. Corbett, Keith O. Hodgson, Britt Hedman, and T. Daniel P. Stack

Department of Chemistry, Stanford University, Stanford, CA 94305 and Stanford Synchrotron Radiation Laboratory, SLAC, Stanford, CA 94309

μ - η^2 : η^2 -peroxodicopper(II) (**P**) and bis(μ -oxo)dicopper(III) (**O**) complexes are valence isomers that differ by the degree of O₂ reduction and the presence of an O–O bond.^{1,2} These isomers can exist in a measurable equilibrium with a small activation energy.^{3–6} This facile isomerization is significant to the processes of making and breaking an O–O bond, which are key steps in photosynthesis, respiration, and the catalytic cycle of tyrosinase, a binuclear copper enzyme that *ortho*-hydroxylates phenols. The characterized **P** species of oxygenated tyrosinase is accepted as the active oxidant in the oxygen atom transfer reaction, but a transient **O**-type species in which the O–O bond is cleaved prior to oxygen insertion cannot be overlooked.⁷ Understanding these steps in detail is important to the design of synthetic catalysts that use O₂ as a terminal oxidant.

A systematic study of the influence of the Lewis basicity of various anions, i.e. their coordinating ability, on the **P/O** equilibrium was undertaken as a model of substrate binding to the **P** core in tyrosinase.^{5,6} **P/O** mixtures were prepared with the ligand *N,N,N',N'*-tetraethylpropane-1,3-diamine (TEPD) by injecting a CH₂Cl₂ solution of [(TEPD)Cu(CH₃CN)_{*n*}](X), where X[–] = SbF₆[–], CF₃SO₃[–], TsO[–] (*p*-toluenesulfonate) or CH₃SO₃[–],⁸ into a precooled, preoxygenated volume of CH₂Cl₂, tetrahydrofuran (THF) or acetone (–85°C, 1 atm O₂, 1 mM in Cu). **P** and **O** isomers were stable only below –75°C and were identified by their characteristic charge-transfer absorptions (Table 1· Figure 1).¹ In THF or CH₂Cl₂, [(TEPD)₂Cu₂O₂](CF₃SO₃)₂ exhibits rapid, reversible interconversion between equilibrium positions upon temperature change.⁹ A Van't Hoff analysis yields $\Delta H^\circ = -4.3(2) \text{ kJ mol}^{-1}$ and $\Delta S^\circ = -24(2) \text{ J K}^{-1} \text{ mol}^{-1}$ for this **P** $\rightleftharpoons$ **O** equilibrium in THF: **O** is favored enthalpically and **P** is favored entropically, as previously determined for other systems.^{4,5}

More strongly coordinating counteranions bias the **P:O** equilibrium position towards **P**, from ~10:90 with SbF₆[–] to ~100:0 with CH₃SO₃[–] (Figure 1a). The **P:O** ratio follows anion basicity regardless of size: e.g. CH₃SO₃[–] is slightly smaller than CF₃SO₃[–], yet the more compact **O** isomer is not observed with CH₃SO₃[–]. Such a basicity effect is counter-intuitive, as more electron donation to the Cu₂O₂ core is anticipated to stabilize the higher oxidation state of the copper centers and hence favor the **O** isomer.¹⁰

Titration experiments with competing anions highlight the importance of anion basicity and reveal the existence of specific anion/dication interactions. Addition of a more coordinating anion Y[–] (CF₃SO₃[–], TsO[–], CH₃SO₃[–], CF₃CO₂[–], PhCO₂[–]) to a preformed **P/O** solution with a “weaker” anion X[–] (SbF₆[–], CF₃SO₃[–]) results in a rapid, isosbestic isomerization in the direction **O** $\rightarrow$ **P**. Spectroscopically pure **P** species are obtained by addition of 1.0 equivalent of TsO[–], CH₃SO₃[–], CF₃CO₂[–] or PhCO₂[–] per binuclear complex (Figure 1b).^{6,11} In all cases, no significant spectral changes occur upon addition of more than 1.0 equivalent of Y[–] per Cu₂O₂ species, and the final **P:O** ratio is the same as per direct oxygenation of [(TEPD)Cu(CH₃CN)_{*n*}](Y) (Y[–] = CF₃SO₃[–], TsO[–] or CH₃SO₃[–]). In contrast, the reverse titration of a

$[(\text{TEPD})_2\text{Cu}_2\text{O}_2](\text{Y})_2$ solution with a “weaker” X^- anion is incomplete even with 200 equivalents of X^- per dimer.¹² Overall, these experiments suggest that each dicationic Cu_2O_2 species is associated with *one* anion intimately.¹³

Extended X-ray Absorption Fine Structure (EXAFS) analysis of a frozen THF solution of $[(\text{TEPD})_2\text{Cu}_2\text{O}_2](\text{CH}_3\text{SO}_3)_2$ provides structural evidence of the close association between the anion and the Cu_2O_2 core in this spectroscopically pure **P** species. The EXAFS data are consistent with a side-on peroxo-bridged copper dimer having four Cu–N/O¹⁴ interactions at 1.94 Å, one Cu–O at 2.26 Å, and one Cu···Cu at 3.51 Å (Figure 2a).¹ The scattering atom at 2.26 Å is required for a good fit and is ascribed to a CH_3SO_3^- oxygen atom,¹⁵ consistent with the titration experiments. Coordination of CH_3SO_3^- would place the sulfur atom within 3.3–3.8 Å of the copper centers, which corresponds to the poorly fitted region in the 4-component model. A 5-component EXAFS fit with a Cu···S interaction at a refined distance of 3.47 Å¹⁶ provides a better match to the data (Figure 2b, Table S1). The dual requirement for a Cu–O interaction at 2.26 Å and a Cu···S interaction at 3.47 Å in the EXAFS fit establishes that the CH_3SO_3^- anion is well-ordered at close range from the Cu_2O_2 core.

Density Functional Theory (DFT) calculations¹⁷ support the EXAFS structure. A plausible model consists of the association between a $[(\text{TEPD})_2\text{Cu}_2\text{O}_2]^{2+}$ molecule and one CH_3SO_3^- anion, in accord with the experimental 1:1 anion-to-dimer stoichiometry. Electronic optimization of this model with a Cu···Cu distance fixed to the experimental value of 3.51 Å converges to the **P** species depicted in Figure 3. The CH_3SO_3^- anion bridges the two copper (II) ions through axial positions, a ligation mode documented for the weaker triflate anion in copper dimers.¹⁸ The calculated distances about the copper ions agree closely with the EXAFS results, notably for the axial oxygen atoms. The Cu_2O_2 core exhibits a slight butterfly distortion with a dihedral angle of $\sim 150^\circ$ between the two CuO_2 planes that may account for the weak feature at ~ 500 nm in the UV-Vis spectrum of this **P** species (Figure 1a).¹⁹

The axial ligation mode of the anion is substantiated experimentally. The ~ 600 nm charge-transfer feature of the **P** isomers shifts to lower energies by 12% as the basicity of the counteranion increases from CF_3SO_3^- to PhCO_2^- (Table 1). Since this transition originates from the *out-of-plane* π^* orbital of the peroxo moiety, this large shift strongly suggests that the anion is positioned above the Cu_2O_2 plane. This indirect characterization of the anion-core association may become useful for probing substrate binding to a **P** core.

In conclusion, axial binding of an anion induces an electronic/electrostatic, not steric, preference for the **P** isomer: the O–O bond is not cleaved with an anion positioned axially on the Cu centers. In oxytyrosinase, although the substrate presumably approaches the **P** core from above,²⁰ its subsequent deprotonation yields a phenolate anion, which is much more basic than the anions used here. Such a strong ligand can potentially redefine the equatorial planes of the copper centers with minimal reorganization of the O_2 -derived ligands. Such change could trigger O–O bond cleavage and yield a reactive **O**-type species, as suggested recently.⁷

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Grants from the NIH are gratefully acknowledged: GM-50730 (T.D.P.S.) and RR-01209 (K.O.H.). The SSRL is supported by the DOE-BES and the SSRL XAS facilities by DOE-BER and NIH-NCRR-BTP.

References

1. Mirica LM, Ottewald X, Stack TDP. *Chem Rev* 2004;104:1013–1045. [PubMed: 14871148]
2. Lewis EA, Tolman WB. *Chem Rev* 2004;104:1047–1076. [PubMed: 14871149]
3. Halfen JA, Mahapatra S, Wilkinson EC, Kaderli S, Young VG, Que L Jr, Zuberbühler AD, Tolman WB. *Science* 1996;271:1397–1400. [PubMed: 8596910] Henson MJ, Vance MA, Zhang CX, Liang HC, Karlin KD, Solomon EI. *J Am Chem Soc* 2003;125:5186–5192. [PubMed: 12708870]
4. Cahoy J, Holland PL, Tolman WB. *Inorg Chem* 1999;38:2161–2168. [PubMed: 11671001] Liang HC, Henson MJ, Hatcher LQ, Vance MA, Zhang CX, Lahti D, Kaderli S, Sommer RD, Rheingold AL, Zuberbühler AD, Solomon EI, Karlin KD. *Inorg Chem* 2004;43:4115–4117. [PubMed: 15236520]
5. Mahadevan V, Henson MJ, Solomon EI, Stack TDP. *J Am Chem Soc* 2000;122:10249–10250.
6. Stack TDP. *Dalton Trans* 2003;10:1881–1889.
7. Mirica LM, Vance M, Jackson Rudd D, Hedman B, Hodgson KO, Solomon EI, Stack TDP. *Science* 2005;308:1890–1892. [PubMed: 15976297]
8. The 1:1 TEPD–Cu complex is prepared in a drybox by mixing equimolar amounts of TEPD and $[\text{Cu}(\text{CH}_3\text{CN})_n](\text{X})$ ($n = 3, 3.5, \text{ or } 4$) in CH_2Cl_2 . This clear, colorless solution is used directly. The salts $[\text{Cu}(\text{CH}_3\text{CN})_3](\text{TsO})$ and $[\text{Cu}(\text{CH}_3\text{CN})_{3.5}](\text{CH}_3\text{SO}_3)$ were prepared by adapting the procedure of Kubas GJ, Monzyk B, Crumbliss AL. *Inorg Synth* 1979;19:90–92.
9. In cases where the **P:O** ratio differs significantly from 1:1, the equilibrium presumably occurs, but is difficult to detect due to the predominance of one isomer and the broadness of the UV-Vis features.
10. Hatcher LQ, Vance MA, Sarjeant AAN, Solomon EI, Karlin KD. *Inorg Chem* 2006;45:3004–3013. [PubMed: 16562956]
11. Strong anions are detrimental to the stability of the Cu_2O_2 species. The use of PhCO_2^- was successful only in acetone at -87°C with slow addition of up to 1.0 equivalent of anion per dimer. The acetate anion behaves similarly, but its **P** species is too unstable to allow a complete titration.
12. The **P:O** ratio is fairly insensitive to ionic strength: only 10% of the total Cu_2O_2 concentration is converted from **P** to **O** with 200 equivalents of CF_3SO_3^- added per $[(\text{TEPD})_2\text{Cu}_2\text{O}_2](\text{CF}_3\text{SO}_3)_2$ (cf. supporting information).
13. The **P:O** ratios of $[(\text{TEPD})_2\text{Cu}_2\text{O}_2](\text{X})_2$ ($\text{X}^- = \text{SbF}_6^-$ or CF_3SO_3^-) in the more polar solvent acetone are consistently $<10:90$, suggesting that anion association is weak in these cases. The more basic TsO^- and CH_3SO_3^- anions yield solutions of the same composition in all three solvents, suggesting that they form an intimate association with the Cu_2O_2 species.
14. Scatterers differing by $Z \pm 1$ are not distinguishable in an EXAFS analysis. For this data, the resolution of such scatterers is $\sim 0.12 \text{ \AA}$.
15. Mahadevan V, Dubois JL, Hedman B, Hodgson KO, Stack TDP. *J Am Chem Soc* 1999;121:5583–5584.
16. $\text{Cu}\cdots\text{S}$ scattering at $3.4 \text{ \AA}$ has the same frequency but opposite phase as $\text{Cu}\cdots\text{Cu}$ scattering at $3.5 \text{ \AA}$. This coincidence prevents independent determination of these two components. Therefore, their σ^2 parameters were constrained to have equal values during refinement.
17. DFT optimizations were performed in vacuo with a B3LYP function, a 6-31G* basis set, and a broken-symmetry wavefunction for the singlet ground state.
18. Van Albada GA, Mutikainen I, Turpeinen U, Reedijk J. *Inorg Chim Acta* 2001;324:273–277. Van Albada GA, Smeets WJJ, Spek AL, Reedijk J. *Inorg Chim Acta* 1997;260:151–161. Dedert PL, Sorrell T, Marks TJ, Ibers JA. *Inorg Chem* 1982;21:3506–3517.
19. Pidcock E, Obias HV, Abe M, Liang HC, Karlin KD, Solomon EI. *J Am Chem Soc* 1999;121:1299–1308.
20. Matoba Y, Kumagai T, Yamamoto A, Yoshitsu H, Sugiyama M. *J Biol Chem* 2006;281:8981–8990. [PubMed: 16436386]

Figure 1.

(a) UV-Vis spectra of $[(\text{TEPD})_2\text{Cu}_2\text{O}_2](\text{X})_2$, with $\text{X}^- = \text{SbF}_6^-$, CF_3SO_3^- and CH_3SO_3^- in THF at -85°C . (b) Addition of $[\text{N}(n\text{-Bu})_4](\text{PhCO}_2)$ (0–0.9 equivalent per dicopper species) to an acetone solution of $[(\text{TEPD})_2\text{Cu}_2\text{O}_2](\text{CF}_3\text{SO}_3)_2$ at -87°C .¹¹

Figure 2. EXAFS data (inset) and Fourier transforms with offset fit residuals (bottom) of $[(\text{TEPD})_2\text{Cu}_2\text{O}_2](\text{CH}_3\text{SO}_3)_2$ (—). **(a)** 4-component fit (---); **(b)** 5-component fit (---): 4 Cu–N/O = 1.94 Å, 1 Cu–O = 2.26 Å, 6 Cu···C = 2.88 Å, 1 Cu···S = 3.47 Å, 1 Cu···Cu = 3.51 Å.

Figure 3. DFT-optimized geometry of $\{[(\text{TEPD})_2\text{Cu}_2\text{O}_2](\text{CH}_3\text{SO}_3)\}^+$. $\text{Cu}-\text{O}_{\text{eq}} = 1.94 \text{ \AA}$ (avg), $\text{Cu}-\text{N} = 2.02 \text{ \AA}$ (avg), $\text{Cu}-\text{O}_{\text{ax}} = 2.26 \text{ \AA}$, $\text{Cu}\cdots\text{C} = 2.87 \text{ \AA}$ (avg), $\text{Cu}\cdots\text{S} = 3.36 \text{ \AA}$ and $\text{Cu}\cdots\text{Cu}$ fixed to $3.51 \text{ \AA}$.

Scheme 1. Equilibrium between TEPD-based **P** and **O** species

Table 1
[(TEPD)₂Cu₂O₂](X)₂ solutions: UV-Vis features and **P:O** ratios

X⁻	solvent	P:O^a	UV-Vis features: O	P	λ: nm (ε: mM⁻¹ cm⁻¹)^b	P
SbF ₆ ⁻	THF	10:90	326 (12)	~360 (sh)	422 (23)	—
CF ₃ SO ₃ ⁻	THF	50:50	~320 (sh)	363 (16)	416 (13)	605 (0.5)
TsO ⁻	THF	100:0 ^c	—	363 (26)	—	636 (1.3)
CH ₃ SO ₃ ⁻	THF	100:0 ^c	—	362 (28)	—	640 (1.1)
CF ₃ CO ₃ ^{-d}	THF	100:0 ^c	—	364 (20)	—	660 (1.2)
PhCO ₂ ^{-d}	acetone	100:0 ^c	—	362 (21)	—	690 (1.0)

^a Approximate **P:O** ratio at -85°C, estimated from the average molar extinction coefficients of the 360 and 400 nm bands (26 and 24 mM⁻¹ cm⁻¹, respectively).¹

^b A weak band is also present at ~500 nm for the **P** species (Figure 1a); sh = shoulder.

^c At the experimental detection limit (< 5% in **O**).

^d Solutions formed by titration (see text).