Published in final edited form as:

J Org Chem. 2005 December 23; 70(26): 10792–10802. doi:10.1021/jo0517848.

Squaric Acid-Based Peptidic Inhibitors of Matrix Metalloprotease-1 (MMP-1)

M. Burak Onaran[†], Anthony B. Comeau[†], and Christopher T. Seto^{*}

Department of Chemistry, Brown University, Providence, RI 02912

Abstract

A series of squaric acid-peptide conjugates were synthesized and evaluated as inhibitors of MMP-1. The cyclobut-3-enedione core was substituted at the 3-position with several functional groups, such as -N(alkyl)OH, -NHOH and -OH, that are designed to bind to the zinc atom in the active site of the metalloprotease. The 4-position of the cyclobut-3-enedione was derivatized with mono- or dipeptides that are designed to bind in the S1' and S2' subsites of the enzyme, and position the metal chelating group appropriately in the active site for binding to zinc. Positional scanning revealed that -N(Me)OH provided the highest level of inhibition among the chelating groups that were tested, and Leu-Tle-NHMe was the preferred amino acid sequence. A combination of these groups yielded an inhibitor with an IC₅₀ value of 95 μM. For one inhibitor, conversion of one of the carbonyl groups on the cyclobut-3-enedione core to a thiocarbonyl group resulted in a 18-fold increase in potency, and yielded a compound with an IC₅₀ value of 15 μM.

Introduction

Matrix Metalloproteases (MMPs)

Matrix metalloproteases are a family of structurally related endopeptidases that degrade and remodel components of the extracellular matrix (ECM).¹ These enzymes regulate structure and sustain a balanced composition of the ECM, two processes that are important for maintaining normal physiology in a number of tissues. For example, MMPs play a crucial role in embryonic development, healing and reproduction.

The activity of MMPs is normally regulated at three levels: 1) gene transcription, 2) activation of MMP propeptides, and 3) inhibition of MMPs by tissue inhibitors of metalloproteases (TIMPs).^{2,3} Overexpression of MMPs and deregulation of their activity is associated with a variety pathological conditions including tumor growth and metastasis,⁴ angiogenesis,⁵ destruction of joints that causes osteoarthritis⁶ and rheumatoid arthritis,⁷ periodontal disease⁸ and multiple sclerosis.⁹ Therefore, there is significant interest in developing MMP inhibitors for therapeutic applications.

Biological Applications of Squaric Acids

Squaric acid is a molecule that has significant aromatic character. In one resonance form it has two π electrons and a negative charge on each of the carbonyl oxygen atoms (Figure 1). The conjugate base of squaric acid can serve as an electrostatic mimic of negatively charged groups that are common in biology including carboxylates and phosphate mono- and diesters. As a

Christopher_Seto@brown.edu.

[†]These two authors contributed equally to this work.

result, derivatives of squaric acid have been used as a replacement for these groups in a number of medicinal applications.

Our research group has used squaric acid derivatives to mimic the phosphate group in phosphotyrosine residues. We prepared a number of 3-hydroxy-4-aryl-cyclobut-3-enediones as non-hydrolyzable isosteres of aryl phosphate esters and found that these compounds are effective inhibitors of protein tyrosine phosphatases.¹⁰ Kim and coworkers have used phosphonocyclobutenedione as a mimic of pyrophosphate and found that it is a selective inhibitor of DNA polymerases from several viruses.¹¹ Sekine has used a diamide of squaric acid to replace a phosphate diester linkage in an oligodeoxynucleotide.¹²

Derivatives of squaric acid have also been used to mimic carboxylates by a number of investigators. Shinada replaced the γ -carboxylic acid of a glutamate residue within a polyamine toxin with a squaric acid derivative. The resulting compound was a selective agonist of ionotropic glutamate receptors.¹³ Sun and coworkers used a derivative of squaric acid in their investigations of an NMDA antagonist that regulates the activation of glutamate receptors.¹⁴ In this example, squaric acid mimics the natural glutamate agonist of the neuronal receptors. In its application as a guanidinium isostere, Butera used diaminocyclobutenedione as a replacement for the *N*-cyanoguanidine group in a bladder-selective potassium channel opener that is used as a treatment for urge urinary incontinence.¹⁵

Hydroxamic acids are potent inhibitors of zinc metalloproteases. X-ray crystal structures show that hydroxamic acids chelate to the active site zinc atom as shown in structure **I** (Figure 2).¹⁶ In addition, Bruckner and coworkers have demonstrated that vinylogous hydroxamic acids that are derived from squaric acid are good metal chelators (see structure **II** in Figure 2).¹⁷ These two observations prompted us to investigate the potential of vinylogous hydroxamic acids that are based upon squaric acid as inhibitors of MMPs. We have derivatized the vinylogous hydroxamic acids with peptides in order to target them to the active site of the proteases.

Results and Discussion

Synthesis of Compounds 4a–4f

We began our studies by making a series of simple derivatives to determine what functional group is preferred at the R¹ position of the inhibitors (Scheme 1). Squaric acid **1** was converted to its dimethyl ester **2** by treating it with trimethyl orthoformate. Reaction of compound **2** with a series of hydroxylamines gave vinylogous hydroxamic acids **3a–e**. Substitution of the remaining methyl ester with several primary amines gave inhibitors **4a–f**.

These compounds were screened for activity against MMP-1. Assays were performed in a buffer of 200 mM NaCl, 50 mM Tris, 5 mM CaCl₂, 20 μ M ZnSO₄, 0.05% Brij 35 at pH 7.6 using the fluorogenic substrate Dnp-Pro-Cha-Gly-Cys(Me)-His-Ala-Lys(Nma)-NH₂. In this substrate, *N*-methylantranilic acid (Nma) is a fluorophore and the dinitrophenyl (Dnp) group is a quencher.^{18,19} The progress of the reactions was monitored by fluorescence spectroscopy with excitation at 340 nm and emission at 460 nm. For inhibitors **4a–b**, R¹ = H while the R² position was varied between a branched and a straight-chain alkyl group. Between these two compounds, inhibitor **4b**, which incorporates a simple n-hexyl chain, had the better activity. We next screened Me, cyclohexyl and benzyl groups at the R¹ position, and found that large groups are unfavorable at this site. Compounds **4e** and **4f**, where R¹ is cyclohexyl or benzyl, were poor inhibitors that showed no activity up to 10 mM concentration. Compound **4c**, which incorporates a smaller methyl group at the R¹ position, was approximately three times more active than compound **4b**, where R¹ = H. Both inhibitors **4c** and **4d** had IC₅₀ values of 310 μ M.

Monopeptide Inhibitors

We next turned our attention to inhibitors that incorporated a single amino acid in order to determine which amino acid side chain is preferred in the S1' subsite within the context of these inhibitors. As shown in Scheme 2, inhibitors **6a–g** were synthesized by reaction of vinylogous hydroxamic acids **3b** or **3e** with a variety of amino acid methyl esters. We chose amino acids with hydrophobic side chains since data from the literature indicated that such structures are preferred in the S1' subsite.¹

Comparison of inhibitors **6a** and **6b** shows that both methyl and isopropyl groups at the R¹ position are small enough to be tolerated by the enzyme (Table 2). In addition, the enzyme appears to be fairly insensitive to the identity of the R³ side chain, since all of the inhibitors in this series gave IC₅₀ values that range from 190 to 380 μ M. Among these compounds inhibitor **6a** had the best activity with an IC₅₀ value of 190 μ M.

Dipeptide Inhibitors

To improve the potency of the inhibitors, we extended their structure by incorporating a second amino acid that is designed to bind in the S2' subsite. X-ray crystallographic studies have shown that peptide-based hydroxamic acid inhibitors bind in the active site of MMPs by occupying the primed subsites.²⁰ The synthesis of the inhibitors in this series is shown in Scheme 3. For these studies, we used the dibutyl ester of squaric acid **7** as the starting point. We found that this compound is more convenient to work with than the corresponding dimethyl ester, since the dibutyl ester has increased solubility in most common organic solvents.

Squaric acid was treated with tributyl orthoformate to give dibutyl ester **7**. Subsequent reaction of **7** with *N*-methylhydroxylamine yielded compound **8**. The dipeptide building blocks were prepared by coupling *N*-Boc amino acids **9a–g** with methylamine to give the corresponding *N*-methyl amides. These were treated with trifluoroacetic acid to give compounds **10a–g**. Coupling of **10a–g** with *N*-Boc-Ile or *N*-Boc-Leu, followed by Boc deprotection gave dipeptides **11a–h**. Finally, reaction of the dipeptides with compound **8** in methanol at room temperature provided the desired inhibitors **12a–h**.

Most of the inhibitors in this series incorporate the side chain of Ile at the R³ position because the data presented in Table 2 indicated that Ile is preferred in the S1' subsite. The side chains at the R⁴ position were chosen based upon literature precedent, which suggest that aromatic and hydrophobic amino acids are preferred at the S2' subsite.¹ For inhibitors **12a–g**, which incorporate an Ile side chain at R³, we did not observe any improvement in activity over the related mono-peptide inhibitor **6a**. In addition, the activity did not depend strongly on the structure of R⁴, since we observe a difference in IC₅₀ values of only 2.5-fold among these seven compounds.

Compounds **12a** and **12h** provide an interesting comparison. Both incorporate a bulky *t*-butyl group at R⁴. Compound **12a** incorporates the side chain of Ile at R³, and was the least active among **12a–g**. By contrast, **12h** has a Leu side chain at R³ and it is 2–5-fold more active than any of the other mono- or dipeptide-based inhibitors. These data suggest that binding interactions in the S1' and S2' subsites are dependent on one another. For the mono-peptide inhibitors that leave the S2' subsite empty (Table 2, compare inhibitors **6a** and **6d**), Ile is preferred over Leu at S1'. This selectivity is reversed in the dipeptide inhibitors. With Ile in the S2' subsite, Leu is preferred over Ile at S1' (compare **12a** and **12h**).

Squaric Acid Monoamides

We also made a cursory examination of a simple monoamide of squaric acid as a potential MMP-1 inhibitor. Scheme 4 shows the synthesis of compound **14**, which is the squaric acid

analog of the vinylogous hydroxamic acid inhibitor **12h**. Reaction of H-Leu-Tle-NHMe (**11h**) with dimethyl squarate **2** gave compound **13**, which was subsequently hydrolyzed under acidic conditions to give compound **14**.

When compound **14** was assayed against MMP-1, we observed no inhibition up to a concentration of 200 μ M. In retrospect, this result is not surprising since the bite angle between the two oxygen atoms in squaric acid is known to be too large to form a strong five-membered chelate with Zn^{+2} (right side of Figure 3).²¹ By comparison, the vinylogous hydroxamic acid inhibitors have the potential to form a six-membered chelate with zinc (left side of Figure 3). This structure has a reduced bite angle and a shorter distance between the two chelating oxygen atoms. Both of these factors favor binding to the metal center.²²

Thiocarbonyl- Vs. Carbonyl-Containing Inhibitors

Despite the fact that hydroxamic acids are potent inhibitors of metalloproteases, they have met with limited success in the clinic because of unfavorable oral bioavailability, stability *in vivo* and side effects.²³ As a result, investigators have been working to discover alternate zinc-binding motifs as a substitute for hydroxamic acids. One such example has been published by Cohen and coworkers, who reported hydroxypyridinone and pyrone ligands as promising alternatives to hydroxamic acids.²⁴ Among the compounds assayed, the zinc chelators that incorporated a thiocarbonyl group had lower IC_{50} values when compared to their non-sulfur containing analogues.

To apply this strategy to the squaric/hydroxamic acid hybrids, we first needed to develop a reliable method for incorporating sulfur into the inhibitors. Squaric acid monoester **3b** (Scheme 2) did not react with a variety of thionating agents including P_2S_5 /hexamethyldisiloxane and Lawesson's reagent. By contrast, the squaric acid monoamide **6a** reacted with both P_2S_5 and Lawesson's reagent to give a product that incorporated a single sulfur atom in place of oxygen, as determined by mass spectrometry. Since Lawesson's reagent provided the cleaner of the two reactions, this method was used to convert **6a**, **12d**, **12g** and **12h** into the corresponding thiocarbonyl compounds (Scheme 5).

Determining the Position of Sulfur Incorporation

The inhibitors in Scheme 5 contain three or four different carbonyl groups, any of which could be the site of reaction with Lawesson's reagent. In order to determine the specific site that sulfur was incorporated into the molecules, we prepared two model compounds to aid us with this analysis (Scheme 6). Dibutyl squarate **7** was treated with two equivalents of NH_2 -Phe-OMe to give compound **19**. When compound **19** was treated with less than one equivalent of Lawesson's reagent, the reaction yielded compound **20** in which one carbonyl group on the cyclobutenedione core had been replaced by a thiocarbonyl as determined by mass spectrometry. Ester carbonyl groups do not react with Lawesson's reagent under the reaction conditions that we employed (25 $^\circ\text{C}$). In a similar manner, compound **6e** was converted to **21**.

The ^1H NMR spectrum of compound **19** (Figure 4) has a resonance at 5.0 PPM that corresponds to the α -protons of the two Phe residues (H^a). When this compound is treated with Lawesson's reagent to give compound **20**, one of the α -protons remains at 5.0 PPM (H^c) while the other is shifted to 6.1 PPM (H^b shown in red). H^b shifts downfield since the neighboring nitrogen atom of this Phe residue is in conjugation with the thiocarbonyl group, which is a better electron acceptor than a standard carbonyl. In a similar manner, compound **6e** has a resonance at 5.0 PPM that corresponds the Phe α -proton (H^d). When this compound is converted to the monothiocarbonyl adduct **21**, this proton shifts downfield to 6.2 PPM (H^e shown in red). This observation, along with the fact that compound **3b** (Scheme 2) that also incorporates a – N

(OH)Me group does not react with Lawesson's reagent, suggests that the carbonyl group opposite the Phe residue in **6e** was the site of reaction with Lawesson's reagent. Since we observe a similar shift for the α -proton of the amino acid that is attached directly to the cyclobutenedione ring in compounds **6a**, **12d**, **12g** and **12h**, we infer that all of these compounds react with Lawesson's reagent on the opposite side of the ring from the peptide chain (see the Supporting Information for the spectra).

Additional evidence for the site of sulfur incorporation comes from the chemical shift of the –OH proton of the vinylogous hydroxamic acid in compounds **6e** and **21**. This proton appears at 10.7 ppm in compound **6e**, since it participates in a strong hydrogen bond with the neighboring carbonyl group on the cyclobutenedione ring. However, the analogous proton in compound **21** appears at 8.5–9.0 ppm. This change in chemical shift is partly due to the fact that the thiocarbonyl group is a weaker hydrogen bond acceptor than the carbonyl group.

Effect of the Thiocarbonyl Group on Inhibition

As shown in Table 4, two of the compounds show improved activity against MMP-1 on conversion to their thiocarbonyl analogs, while the activity of the other two remain unchanged. Compound **15** is 2–3 times more potent than **6a**, while the potency of **16** is increased by 18-fold compared to **12d**. By contrast, compounds **12g** and **h** have similar IC_{50} values when compared to their thiocarbonyl analogs **17** and **18**. These results suggest that conversion of a carbonyl group on the cyclobutenedione core of the inhibitors to a thiocarbonyl can be an effective method for improving inhibition. However, this improvement is dependent on the specific structure of the inhibitor. One plausible explanation for the observation that the potencies of compounds **12g** and **h** do not change upon thionation is that the improvement in chelation between the cyclobutene core and the active site zinc atom changes the position of the bound inhibitor in the active site. This geometry change could decrease binding interactions between the peptide portion of the inhibitors and the S1' and S2' subsites, and offset the improved binding to zinc. By contrast, conversion of compounds **6a** and **12d** to their thiocarbonyl analogs could lead to improved binding with the zinc atom, and also may reposition the inhibitor in the active site so that it makes more favorable interactions with the distal enzyme subsites.

Conclusions

We have investigated the potential of a hybrid between squaric and hydroxamic acids to serve as a metal binding motif for the design of metalloprotease inhibitors. While hydroxamic acids are commonly used as MMP inhibitors, to the best of our knowledge this report represents the first use of a squaric acid derivative as a warhead for the design of inhibitors of metalloproteases. The squaric/hydroxamic acid hybrids are generally not as potent as hydroxamic acid-based inhibitors, many of which have inhibition constants in the nM range. However, since hydroxamic acids have not met with much success in the clinic, these hybrids could serve as an alternate starting point for the design of inhibitors with perhaps improved pharmacological properties. The structure of the peptidic portion of the inhibitors helps target them to the active site of zinc proteases, rather than to other classes of enzymes for which squaric acid derivatives can serve as inhibitors.

Among the alkyl groups that we have examined at the –N(OH)alkyl position of the inhibitors, small substituents such as Me and *i*-Pr are well accommodated in the active site of MMP-1, while sterically demanding groups such as Bn and cyclohexyl are too large and lead to poor activity. Among the inhibitors that do not contain a sulfur atom, we found that the dipeptide Leu-Tle-NHMe provided the highest level of activity when attached to the cyclobutenedione core.

We also developed a regioselective method for converting one of the carbonyl groups on the core into the corresponding thiocarbonyl compound. This reaction occurs specifically at the position on the opposite side of the cyclobutene ring from the peptide substituent. Thionation of the inhibitors leads to improved activity in some cases, but not in others. Compound **12d** gave the largest increase in potency upon thionation. The 18-fold improvement resulted in an inhibitor with an IC₅₀ value of 15 μ M against MMP-1.

Experimental Section

Full characterization for compounds **3b**, **3c** and **3e** has been reported in the literature.¹⁷

3-(Hydroxyamino)-4-methoxy-3-cyclobutene-1,2-dione (**3a**)

3,4-Dimethoxy-3-cyclobutene-1,2-dione **1** (200 mg, 1.4 mmol) was added to 10 mL of MeOH and stirred until all of the solids had dissolved. To this mixture was added hydroxylamine hydrochloride (104 mg, 1.5 mmol), the reaction mixture was cooled to 0 °C, and KOH (94.3 mg, 1.7 mmol) dissolved in 5 mL of MeOH was added. At this point the reaction contained a white precipitate. The reaction was monitored by TLC and upon disappearance of the starting material the mixture was filtered and the solid that was collected was washed with cold MeOH. The filtrate and washes were combined and solvent was evaporated. The crude material was purified by column chromatography (1:9 MeOH/CH₂Cl₂) to give compound **3a** as a yellow solid (0.093 g, 0.650 mmol, 47%). ¹H NMR (300 MHz, CD₃OD) δ 4.37 (s, 3 H); ¹³C NMR (100 MHz, CD₃OD) δ 184.9, 181.5, 174.4, 170.1, 60.1; HRMS-ESI (M + H⁺) calcd for C₅H₆NO₄ 144.0297, found 144.0292.

3-[(Benzyl)hydroxyamino]-4-methoxy-3-cyclobutene-1,2-dione (**3d**)

This compound was prepared using *N*-benzylhydroxylamine hydrochloride (0.367 g, 2.3 mmol) and **1** (0.30 g, 2.1 mmol) according to the procedure described above for the preparation of compound **3a**. Purification of **3d** was performed by column chromatography (5:95 MeOH/CH₂Cl₂) to obtain a white solid (0.318 g, 1.37 mmol, 65%). ¹H NMR (300 MHz, CD₃OD) δ 7.39 (m, 5 H), 4.88 (s, 3 H), 4.38 (s, 3 H); ¹³C NMR (75 MHz, CD₃OD) δ 185.0, 181.3, 175.0, 169.0, 135.0, 129.0, 128.8, 128.5, 60.3, 57.3; HRMS-ESI (M + Na⁺) calcd for C₁₂H₁₁NO₄Na 256.0586, found 256.0590.

3-(Hydroxyamino)-4-[(2-methylpropyl)amino]-3-cyclobutene-1,2-dione (**4a**)

To a solution of compound **3a** (0.055 g, 0.383 mmol) dissolved in MeOH (1 mL), isobutylamine (0.062 g, 0.844 mmol) was added and the mixture was stirred at room temperature for 25 h. The solvent was evaporated and the product was purified by column chromatography (1:9:90 30% aqueous NH₄OH/MeOH/CH₂Cl₂) giving **4a** (0.009 g, 0.049 mmol, 13%) as a white solid. ¹H NMR (400 MHz, CD₃OD) δ 3.39 (d, *J* = 6.8 Hz, 2 H), 1.85 (m, 1 H), 0.96 (d, *J* = 5.2 Hz, 6 H); ¹³C NMR (100 MHz, CD₃OD) δ 181.1, 180.3, 168.9, 167.8, 52.8, 31.4, 20.0; HRMS-ESI (M + H⁺) calcd for C₈H₁₂N₂O₃ 185.0926, found 185.0932.

3-(Hexylamino)-4-(hydroxyamino)-3-cyclobutene-1,2-dione (**4b**)

Compound **3b** (100 mg, 0.70 mmol) was dissolved in MeOH (10 mL) and hexylamine (84.9 mg, 0.84 mmol) was added to the solution. The reaction was stirred for 12 h at room temperature, after which time TLC showed that all the starting material was consumed. The solvent was then removed by rotary evaporation. The crude material was purified by column chromatography (1:9:90 30% aqueous NH₄OH/MeOH/CH₂Cl₂) to yield **4b** as a yellow oil (17.8 mg, 0.084 mmol, 12%). ¹H NMR (300 MHz, DMSO-*d*₆) δ 3.35 (m, 2 H), 1.50 (m, 2 H), 1.27 (m, 6 H), 0.86 (m, 3 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 184.2, 183.6, 170.2, 169.6,

44.1, 31.7, 31.6, 26.4, 23.0, 14.8; HRMS-ESI ($M + H^+$) calcd for $C_{10}H_{17}N_2O_3$ 213.1239, found 213.1250.

3-(Hexylamino)-4-(hydroxymethylamino)-3-cyclobutene-1,2-dione (**4c**)

To a solution of **3b** (0.157 g, 1.0 mmol) in MeOH (1 mL), hexylamine (0.111 g, 1.1 mmol) was added and the mixture was stirred at room temperature for 4 h. The solvent was evaporated and the crude product was purified by column chromatography (1:9:90 30% aqueous $NH_4OH/MeOH/CH_2Cl_2$) giving **4c** (0.19 g, 0.84 mmol, 84%) as a white solid. 1H NMR (400 MHz, $CDCl_3$) δ 11.19 (s, 1 H), 7.50 (s, 1 H), 3.59 (m, 5 H), 1.64 (m, 2 H), 1.32 (m, 6 H), 0.88 (t, $J = 6.5$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 178.9, 177.5, 166.8, 164.9, 45.0, 41.4, 31.4, 30.9, 26.1, 22.6, 14.0; HRMS-FAB ($M + Na^+$) calcd for $C_{11}H_{18}N_2O_3Na$ 249.1215, found 249.1220.

3-(Hydroxymethylamino)-4-(pentylamino)-3-cyclobutene-1,2-dione (**4d**)

Compound **4d** (0.058 g, 0.27 mmol, 37%) was prepared from **3b** (0.118 g, 0.75 mmol) and pentylamine (0.65 g, 0.75 mmol) according to the procedure used for preparing **4c**. 1H NMR (400 MHz, $CDCl_3$) δ 7.16 (s, 1 H), 3.56 (m, 5 H), 1.64 (m, 2 H), 1.33 (m, 4 H), 0.89 (t, $J = 6.8$ Hz, 3 H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 178.9, 177.4, 166.7, 165.1, 45.0, 41.4, 30.7, 28.6, 22.3, 14.0; HRMS-FAB ($M + Na^+$) calcd for $C_{10}H_{16}N_2O_3Na$ 235.1059, found 235.1066.

3-[(Cyclohexyl)hydroxyamino]-4-(hexylamino)-3-cyclobutene-1,2-dione (**4e**)

Compound **4e** (0.0047 g, 0.016 mmol, 13%) was prepared as a white solid from **3c** (0.027 g, 0.12 mmol) and hexylamine (0.015 g, 0.144 mmol) according to the procedure used for preparation of **4b**. 1H NMR (400 MHz, $DMSO-d_6$) δ 7.34 (br s, 2 H), 3.76 (br s, 1 H), 3.32 (s, 4 H), 3.52 (s, 2 H), 1.88 (m, 2 H), 1.70 (m, 2 H), 1.57 (s, 3 H), 1.27 (m, 10 H), 1.53 (m, 3 H), 0.86 (m, 3 H); ^{13}C NMR (100 MHz, $DMSO-d_6$) δ 181.7, 181.4, 167.3, 166.4, 51.5, 42.7, 33.2, 30.3, 30.2, 25.0, 24.3, 23.5, 21.5, 13.4; HRMS-ESI ($M + H^+$) calcd for $C_{16}H_{27}N_2O_3$ 295.2022, found 295.2018.

3-[(Benzyl)hydroxyamino]-4-(hexylamino)-3-cyclobutene-1,2-dione (**4f**)

Compound **4f** (0.232 g, 0.77 mmol, 89%) was prepared as a white solid from **3d** (0.200 g, 0.86 mmol) and hexylamine (0.101 g, 1.00 mmol) according to the procedure used for the preparation of **4b**. 1H NMR (300 MHz, CD_3OD) δ 7.36 (m, 5 H), 4.94 (s, 2 H), 3.58 (t, $J = 7.0$ Hz, 2 H), 1.60 (dd, $J = 13.9, 6.9$ Hz, 2 H), 1.34 (m, 6 H), 0.91 (t, $J = 6.7$ Hz, 3 H); ^{13}C NMR (75 MHz, CD_3OD) δ 180.4, 178.6, 168.2, 166.1, 135.6, 129.1, 128.7, 128.3, 57.5, 44.4, 31.6, 26.1, 22.7, 13.4; HRMS-ESI ($M + H^+$) 303.1709, found 303.1711.

3-(Hydroxymethylamino)-4-(L-isoleucine methyl ester)-3-cyclobutene-1,2-dione (**6a**)

To a solution of **3b** (0.219 g, 1.39 mmol) dissolved in MeOH (5 mL), the hydrochloride salt of L-isoleucine methyl ester (**5a**) (0.229 g, 1.26 mmol) was added. To this stirred solution KOH (0.0713 g, 1.27 mmol) was added, and immediately a white precipitate formed. The reaction was stirred at room temperature for 12 h and solvent was evaporated. Purification was performed by column chromatography (1:9:90 30% aqueous $NH_4OH/MeOH/CH_2Cl_2$) to yield a yellowish flaky solid (0.238 g, 0.88 mmol, 70%). 1H NMR (300 MHz, CD_3OD) δ 5.04 (br s, 2 H), 4.81 (d, $J = 7$ Hz, 2 H), 3.78 (s, 3 H), 3.50 (s, 3 H), 2.03 (m, 1 H), 1.54 (m, 1 H), 1.30 (m, 1 H), 0.97 (m, 6 H); ^{13}C NMR (75 MHz, CD_3OD) δ 180.4, 180.0, 173.0, 167.6, 167.0, 62.4, 52.8, 41.4, 39.4, 25.8, 15.6, 11.6; HRMS-ESI ($M + H^+$) calcd for $C_{12}H_{19}N_2O_5$ 271.1294, found 271.1287.

3-[Hydroxy(1-methylethyl)amino]-4-(L-isoleucine methyl ester)-3-cyclobutene-1,2-dione (6b)

Compound **6b** (0.034 g, 0.11 mmol, 26%) was prepared as a colorless solid starting from **3e** (0.080 g, 0.432 mmol) and **5a** (0.120 g, 0.56 mmol) according to the procedure used for the preparation of **4b**. ¹H NMR (300 MHz, CD₃OD) δ 7.28 (br s, 1 H), 4.83 (br s, 1 H), 4.52 (br s, 1 H), 3.73 (s, 3 H), 2.01 (s, 1 H), 1.49 (m, 1 H), 1.31 (m, 7 H), 0.93 (m, 6 H); ¹³C NMR (75 MHz, CDCl₃) δ 179.1, 179.0, 172.0, 166.7, 165.5, 61.8, 55.5, 52.7, 38.7, 25.1, 19.8, 15.5, 11.8; HRMS-ESI (M + H⁺) calcd for C₁₄H₂₃N₂O₅ 299.1607, found 299.1602.

3-(Hydroxymethylamino)-4-(D,L-norleucine methyl ester)-3-cyclobutene-1,2-dione (6c)

Compound **6c** (0.09 g, 0.33 mmol, 45%) was prepared from **3b** (0.118 g, 0.75 mmol), D,L-norleucine methyl ester HCl salt (0.136 g, 0.75 mmol) and KOH (0.042 g, 0.75 mmol) according to the procedure used for preparing **6a**. ¹H NMR (400 MHz, CDCl₃) δ 7.15 (d, *J* = 8.4 Hz, 1 H), 4.79 (dt, *J* = 9.1, 4.8 Hz, 1 H), 3.73 (s, 3 H), 3.55 (s, 3 H), 1.95 (m, 1 H), 1.82 (m, 1 H), 1.36 (m, 4 H), 0.89 (t, *J* = 7.1 Hz, 3 H); ¹³C NMR (100 MHz, CDCl₃) δ 178.8, 178.2, 172.0, 165.9, 165.2, 57.0, 52.5, 41.3, 32.5, 27.6, 22.2, 13.8; HRMS-ESI (M + H⁺) calcd for C₁₂H₁₉N₂O₅ 271.1294, found 271.1285.

3-(Hydroxymethylamino)-4-(L-leucine methyl ester)-3-cyclobutene-1,2-dione (6d)

Compound **6d** was prepared as a white solid (0.023 g, 0.076 mmol, 38%) starting from **3b** (0.031 g, 0.20 mmol) and **5c** (0.047 g, 0.26 mmol) according to the procedure used to prepare **6a**. ¹H NMR (300 MHz, CDCl₃) δ 7.14 (br s, 1 H), 4.87 (br s, 1 H), 3.72 (s, 3 H), 3.55 (s, 3 H), 1.75 (m, 1 H), 0.96 (d, *J* = 4.8 Hz, 6 H); ¹³C NMR (100 MHz, CD₃OD) δ 184.5, 183.5, 174.1, 170.3, 168.7, 54.8, 53.0, 31.3, 25.9, 23.4, 21.7; HRMS-ESI (M + Na⁺) calcd for C₁₂H₁₈N₂O₅Na 293.1113, found 293.1118.

3-(Hydroxymethylamino)-4-(L-phenylalanine methyl ester)-3-cyclobutene-1,2-dione (6e)

Compound **6e** (0.135 g, 0.44 mmol, 59%) was prepared from **3b** (0.118 g, 0.75 mmol), L-phenylalanine methyl ester HCl salt (0.216 g, 1.0 mmol) and KOH (0.056 g, 1.0 mmol) according to the procedure used for preparing **6a**. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.75 (s, 1 H), 7.79 (s, 1 H), 7.25 (m, 5 H), 5.06 (s, 1 H), 3.70 (s, 3 H), 3.35 (s, 3 H), 3.28 (dd, *J* = 14.0, 3.9 Hz, 1 H), 3.18 (m, 1 H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ 180.0, 179.2, 171.9, 166.9, 166.0, 137.7, 129.6, 128.8, 127.0, 57.6, 52.8, 41.2, 37.7; HRMS-FAB (M + Na⁺) calcd for C₁₅H₁₆N₂O₅Na 327.0957, found 327.0948.

3-(Hydroxymethylamino)-4-(L-tryptophan methyl ester)-3-cyclobutene-1,2-dione (6f)

Compound **6f** (0.16 g, 0.47 mmol, 62%) was prepared from **3b** (0.118 g, 0.75 mmol), L-tryptophan methyl ester HCl salt (0.255 g, 1.0 mmol) and KOH (0.056 g, 1.0 mmol) according to the procedure used for preparing **6a**. ¹H NMR (300 MHz, CD₃OD) δ 7.56 (d, *J* = 7.7 Hz, 1 H), 7.34 (d, *J* = 8.0 Hz, 1 H), 7.07 (m, 3 H), 5.13 (dd, *J* = 7.7, 5.1 Hz, 1 H), 3.76 (s, 3 H), 3.43 (m, 5 H); ¹³C NMR (75 MHz, CD₃OD) δ 179.6, 179.0, 172.4, 166.9, 166.2, 137.0, 127.7, 123.7, 121.5, 119.0, 118.2, 111.3, 109.1, 57.7, 52.1, 40.3, 28.7; HRMS-ESI (M + Na⁺) calcd for C₁₇H₁₇N₃O₅Na 366.1066, found 366.1075.

3-(Hydroxymethylamino)-4-(L-methionine methyl ester)-3-cyclobutene-1,2-dione (6g)

Compound **6g** (0.082 g, 0.28 mmol, 38%) was prepared from **3b** (0.118 g, 0.75 mmol), L-methionine methyl ester HCl salt (0.15 g, 0.75 mmol) and KOH (0.042 g, 0.75 mmol) according to the procedure used for preparing **6a**. ¹H NMR (400 MHz, CDCl₃) δ 7.30 (d, *J* = 7.8 Hz, 1 H), 4.93 (dd, *J* = 12.7, 8.5 Hz, 1 H), 3.75 (s, 3 H), 3.55 (s, 3 H), 2.60 (m, 2 H), 2.27 (m, 1 H), 2.17 (td, *J* = 14.7, 7.2 Hz, 1 H), 2.10 (s, 3 H); ¹³C NMR (100 MHz, CDCl₃) δ 178.8, 178.3,

171.6, 165.9, 165.3, 56.0, 52.8, 41.4, 31.8, 30.1, 15.4; HRMS-FAB ($M + Na^+$) calcd for $C_{11}H_{16}N_2O_5SNa$ 311.0678, found 311.0685.

3-Butoxy-4-(hydroxymethylamino)-3-cyclobutene-1,2-dione (**8**)

To a stirred solution of HONHMe HCl salt (2.631 g, 31.5 mmol) in MeOH (30 mL), KOH (1.767 g, 31.5 mmol) and compound **7** (4.766 g, 21.0 mmol) were added. The mixture was stirred overnight at room temperature, the solvent was evaporated, and the crude product was washed with H_2O , and then dissolved in EtOAc. The aqueous layer was extracted with EtOAc. The organic extracts were combined, dried over $MgSO_4$, and the solvents were evaporated under reduced pressure. The crude product was purified by column chromatography (3% MeOH/ CH_2Cl_2) to yield compound **8** as a pale yellow solid (3.416 g, 17.2 mmol, 82%). 1H NMR (400 MHz, $CDCl_3$) δ 4.70 (t, $J = 6.6$ Hz, 2 H), 3.52 (s, 3 H), 1.77 (m, 2 H), 1.44 (m, 2 H), 0.97 (t, $J = 7.4$ Hz, 3 H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 184.0, 180.7, 174.3, 169.1, 74.2, 41.6, 32.3, 18.9, 14.0; HRMS-FAB ($M + Na^+$) calcd for $C_9H_{13}NO_4Na$ 222.0742, found 222.0740.

L-tert-Leucine methyl amide (**10a**)

Compound **10a** was synthesized starting from *N*-Boc-protected amino acid **9a**. To a stirred solution of *N*-hydroxysuccinimide (NHS, 0.925 g, 8.04 mmol) in DMF (20 mL) under N_2 , NH_2Me (4.02 mL, 2.0 M in MeOH) was added via syringe and the mixture was stirred for 30 min at room temperature. Formation of a white precipitate was observed. Addition of *N*-Boc-protected amino acid **9a** (1.55 g, 6.7 mmol) and cooling the mixture to 0 °C was followed by addition of 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC, 1.54 g, 8.04 mmol). The mixture was stirred for 1 h at 0 °C and 2 h at room temperature and then diluted with EtOAc. The solution was washed with H_2O , the organic layer was separated, and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed with saturated $NaHCO_3$, H_2O , and brine, dried over $MgSO_4$, and the solvents were evaporated under reduced pressure. The crude product was purified by column chromatography (3% MeOH/ CH_2Cl_2), washed with heptane, dried *in vacuo* and dissolved in CH_2Cl_2 (15 mL). To this stirred solution, TFA (15 mL) was added and the mixture was stirred for 40 min at room temperature. The solvents were removed *in vacuo*, and the residue was partitioned between CH_2Cl_2 (90 mL) and saturated aqueous $NaHCO_3$ (90 mL). The organic phase was separated and the aqueous phase was extracted with CH_2Cl_2 . The organic extracts were combined and dried over Na_2SO_4 , and the solvents were evaporated. Purification was performed by column chromatography (5:95 MeOH/ CH_2Cl_2) yielding **10a** as a colorless oil (0.637 g, 4.42 mmol, 66%). 1H NMR (300 MHz, CD_3OD) δ 2.99 (s, 1 H), 2.75 (s, 3 H), 0.98 (s, 9 H); ^{13}C NMR (75 MHz, CD_3OD) δ 175.2, 63.9, 34.0, 25.9, 24.9; HRMS-FAB ($M + Na^+$) calcd for $C_7H_{16}N_2ONa$ 167.1160, found 167.1163.

L-Cyclohexylglycine methyl amide (**10b**)

Compound **10b** was synthesized starting from *N*-Boc-protected amino acid **9b**. To a stirred solution of *N*-Boc-protected amino acid **9b** (1.16 g, 4.52 mmol) in CH_2Cl_2 (20 mL), HOBt (760 mg, 4.97 mmol) was added and then EDC (911 mg, 4.75 mmol) and mixture was stirred at 0 °C for 30 min. Formation of a white precipitate was observed. To this solution NH_2Me (2.49 mL, 2.0 M in MeOH) was added via syringe and reaction was allowed to stir overnight. The solution was diluted with CH_2Cl_2 (50 mL) and washed once with 1M citric acid (30 mL), once with saturated $NaHCO_3$ (30 mL) and then with brine (30 mL). The organic layer was then dried over $NaSO_4$ and the solvent was evaporated under reduced pressure. The crude product was purified by column chromatography (3% MeOH/ CH_2Cl_2) and the product was dissolved in CH_2Cl_2 (15 mL). To this stirred solution TFA (15 mL) was added and the mixture was stirred at room temperature for 40 min. The solvents were removed *in vacuo*, and the residue was

partitioned between 1:1 CH₂Cl₂/saturated aqueous NaHCO₃. The organic phase was separated and the aqueous phase was extracted with CH₂Cl₂. The organic extracts were combined and dried over Na₂SO₄, and solvents were evaporated. Purification was performed by flash chromatography (5% MeOH/CH₂Cl₂) yielding **10b** as a white solid (440 mg, 2.80 mmol, 62%). ¹H NMR (CDCl₃, 300 MHz) δ 7.34 (br s, 1 H), 3.22 (d, *J* = 3.8 Hz, 1 H), 2.80 (d, *J* = 5.0 Hz, 3 H), 1.27 (m, 14 H); ¹³C NMR (75 MHz, CDCl₃) δ 175.5, 60.5, 41.5, 30.6, 27.0, 26.7, 26.6, 26.5, 26.0; HRMS-FAB (*M* + Na⁺) calcd for C₉H₁₈N₂NaO 193.1317, found 193.1320.

O-Methyl-L-tyrosine methyl amide (10c)

The synthesis and the ¹H NMR spectrum for this compound have been reported previously.²⁵ ¹³C NMR (75 MHz, CD₃OD) δ 175.9, 158.6, 129.9, 129.3, 113.5, 56.5, 54.2, 40.3, 24.7; HRMS-ESI (*M* + Na⁺) calcd for C₁₁H₁₆N₂O₂Na 231.1109, found 231.1115.

L-Phenylalanine methyl amide (10d)

Compound **10d** (105 mg, 0.590 mmol, 52%) was prepared as a white solid from **9d** (300 mg, 1.13 mmol), HOBt (190 mg, 1.24 mmol), EDC (228 mg, 1.19 mmol) and NH₂Me (624 μL of a 2.0 M solution in MeOH) according to the procedure used to prepare **10b**. ¹H NMR (300 MHz, CDCl₃) δ 7.26 (m, 5 H), 3.60 (dd, *J* = 9.5, 4.0 Hz, 1 H), 3.28 (dd, *J* = 13.7, 4.0 Hz, 1 H), 2.82 (d, *J* = 5 Hz, 3 H), 2.67 (dd, *J* = 13.7, 9.5 Hz, 1 H), 1.39 (br s, 2 H); ¹³C NMR (75 MHz, CDCl₃) δ 174.8, 138.0, 129.3, 128.7, 126.8, 56.5, 41.0, 25.8; HRMS-FAB (*M* + Na⁺) calcd for C₁₀H₁₄N₂NaO 201.1004, found 201.1007.

L-Phenylglycine methyl amide (10e)

Compound **10e** (0.836 g, 5.1 mmol, 51%) was prepared from **6e** (2.513 g, 10 mmol), NH₂Me (6.0 mL of a 2.0 M solution in MeOH), NHS (1.381 g, 12 mmol), EDC (2.3 g, 12 mmol) and TFA (15 mL) according to the procedure used for preparing **10a**. The ¹H NMR and ¹³C NMR spectra, and the HRMS for this compound have been reported previously.²⁶

L-Tryptophan methyl amide (10f)

The synthesis and the ¹H NMR spectrum for this compound have been reported previously.²⁴ ¹³C NMR (75 MHz, CD₃OD) δ 174.8, 135.3, 125.9, 121.8, 119.6, 116.9, 116.5, 109.4, 108.3, 54.0, 29.3, 23.3; HRMS-FAB (*M* + Na⁺) calcd for C₁₂H₁₅N₃ONa 240.1113, found 240.1117.

L-Leucine methyl amide (10g)

Compound **10g** (0.548 g, 3.8 mmol, 42%) was prepared from **9g** (2.245 g, 9 mmol), NH₂Me (5.4 mL of a 2.0 M solution in MeOH), NHS (1.243 g, 10.8 mmol), EDC (2.068 g, 10.8 mmol) and TFA (12 mL) according to the procedure used for preparing **10a**. ¹H NMR (300 MHz, CD₃OD) δ 3.33 (dd, *J* = 7.7, 6.5 Hz, 1 H), 2.76 (s, 3 H), 1.68 (qt, *J* = 12.9, 6.5 Hz, 1 H), 1.54 (ddd, *J* = 13.8, 7.4, 6.4 Hz, 1 H), 1.39 (m, 1 H), 0.95 (t, *J* = 6.6 Hz, 6 H); ¹³C NMR (75 MHz, CD₃OD) δ 177.0, 53.2, 44.2, 24.8, 24.5, 22.0, 21.1; HRMS-FAB (*M* + Na⁺) calcd for C₇H₁₆N₂ONa 167.1160, found 167.1166.

L-Isoleucyl-L-*tert*-leucine methyl amide (11a)

To a stirred solution of NHS (0.138 g, 1.2 mmol) in DMF (5 mL) under N₂, **10a** (0.144 g, 1.0 mmol) was added and the mixture was stirred at room temperature for 30 min. Addition of *N*-Boc-L-isoleucine (0.288 g, 1.2 mmol) and cooling the mixture to 0 °C was followed by addition of EDC (0.23 g, 1.2 mmol). The mixture was stirred for 1 h at 0 °C and 2 h at room temperature and then diluted with EtOAc. The solution was washed with H₂O, the organic layer was separated, and the aqueous layer was extracted with EtOAc. The combined organic extracts

were washed with saturated NaHCO₃, H₂O, and brine, dried over MgSO₄, and the solvents were evaporated under reduced pressure. The crude product was purified by column chromatography (3:97 MeOH/CH₂Cl₂), washed with heptane, dried *in vacuo* and dissolved in CH₂Cl₂ (2 mL). To this solution TFA (2 mL) was added and the mixture was stirred at room temperature for 40 min. The solvents and TFA were removed, and the residue was partitioned between CH₂Cl₂ (15 mL) and saturated aqueous NaHCO₃ (15 mL). Organic phase was separated and the aqueous phase was extracted with CH₂Cl₂. The organic extracts were combined and dried over Na₂SO₄, and the solvent was evaporated. Purification was performed by flash chromatography (5:95 MeOH/CH₂Cl₂) yielding **11a** as a colorless oil (0.162 g, 0.63 mmol, 63%). ¹H NMR (300 MHz, CD₃OD) δ 5.51 (s, 1 H), 4.19 (s, 1 H), 2.73 (s, 3 H), 1.79 (m, 1 H), 1.48 (m, 1 H), 1.16 (m, 1 H), 0.95 (m, 15 H); ¹³C NMR (75 MHz, CD₃OD) δ 175.5, 171.8, 60.6, 59.4, 38.6, 33.8, 25.7, 24.6, 23.8, 14.8, 10.7; HRMS-FAB (M + Na⁺) calcd for C₁₃H₂₇N₃O₂Na 280.2001, found 280.2000.

L-Isoleucyl-L-cyclohexylglycine methyl amide (11b)

N-Boc-L-Isoleucine (140 mg, 0.584 mmol) was dissolved in CH₂Cl₂ (20 mL) and the solution was cooled to 0 °C. To this solution HOBt (89.5 mg, 0.584 mmol) and EDC (201 mg, 0.558 mmol) were added and the mixture was stirred for 30 min. To this mixture compound **10b** (83.5 mg, 0.531 mmol) was then added and reaction was stirred overnight. The solution was diluted with CH₂Cl₂ (50 mL) and washed once with 1M citric acid (30 mL), once with saturated NaHCO₃ (30 mL) and then with brine (30 mL). The organic layer was dried over Na₂SO₄ and the solvents evaporated under reduced pressure. The crude product was purified by column chromatography (3% MeOH/CH₂Cl₂) and then dissolved in CH₂Cl₂ (15 mL). To this solution TFA (15 mL) was added and the mixture was stirred at room temperature for 40 min. The solvents and TFA were evaporated, and the residue was partitioned between a 1:1 mixture of CH₂Cl₂:saturated aqueous NaHCO₃. The organic phase was separated and the aqueous phase was extracted with CH₂Cl₂. The organic extracts were combined and dried over Na₂SO₄, and the solvent was evaporated. Purification was performed by flash chromatography (5% MeOH/CH₂Cl₂) yielding **11b** as a white solid (69 mg, 0.244 mmol, 46%). ¹H NMR (400 MHz, CD₃OD) δ 7.94 (d, *J* = 9.0 Hz, 1 H), 6.98 (m, 1 H), 4.26 (m, 1 H), 3.28 (d, *J* = 4.0 Hz, 1 H), 2.78 (d, *J* = 4.8 Hz, 3 H), 1.93 (m, 1 H), 1.75 (m, 7 H), 1.39 (m, 2 H), 1.25 (m, 2 H), 1.13 (m, 4 H), 0.96 (d, *J* = 6.9 Hz, 3 H), 0.90 (t, *J* = 7.3 Hz, 3 H); ¹³C NMR (100 MHz, CD₃OD) δ 174.6, 171.9, 59.8, 57.8, 39.7, 38.1, 29.8, 28.8, 26.1, 25.9, 25.8, 25.8, 24.0, 16.1, 11.9 HRMS-FAB (M + Na⁺) calcd for C₁₅H₂₉N₃NaO₂ 306.2157, found 306.2150.

L-Isoleucyl-O-methyl-L-tyrosine methyl amide (11c)

Compound **11c** (0.265 g, 0.82 mmol, 58%) was prepared as a white solid from **10c** (0.295 g, 1.42 mmol), NHS (0.196 g, 1.7 mmol), *N*-Boc-L-isoleucine (0.408 g, 1.7 mmol), EDC (0.326 g, 1.7 mmol) and TFA (3 mL) according to the procedure used for preparing **11a**. ¹H NMR (300 MHz, CD₃OD) δ 7.16 (d, *J* = 8.5 Hz, 2 H), 6.84 (d, *J* = 8.5 Hz, 2 H), 4.55 (dd, *J* = 8.6, 6.3 Hz, 1 H), 3.76 (s, 3 H), 3.17 (d, *J* = 5.0 Hz, 1 H), 3.07 (dd, *J* = 13.8, 6.2 Hz, 1 H), 2.86 (dd, *J* = 13.8, 8.8 Hz, 1 H), 2.69 (s, 3 H), 1.66 (m, 1 H), 1.18 (m, 1 H), 0.99 (m, 1 H), 0.83 (m, 6 H); ¹³C NMR (75 MHz, CD₃OD) δ 175.6, 172.7, 158.7, 129.9, 128.8, 113.5, 59.6, 54.7, 54.2, 38.5, 37.0, 24.9, 23.5, 14.7, 10.6; HRMS-FAB (M + Na⁺) calcd for C₁₇H₂₇N₃O₃Na 344.1950, found 344.1942.

L-Isoleucyl-L-phenylalanine methyl amide (11d)

Compound **11d** (373 mg, 1.28 mmol, 61%) was prepared as a white solid from **10d** (374 mg, 2.1 mmol), HOBt (322 mg, 2.1 mmol), EDC (383 mg, 2.0 mmol) and *N*-Boc-L-Isoleucine (456 mg, 1.9 mmol) according to the procedure that was used to prepare **11b**. ¹H NMR (400 MHz, CDCl₃) δ 8.04 (d, *J* = 8.8 Hz, 1 H), 7.18 (m, 5 H), 4.78 (dt, *J* = 8.7, 6.3 Hz, 1 H), 2.98 (dd, *J*

= 13.8, 8.8 Hz, 1 H), 2.68 (d, J = 4.8 Hz, 3 H), 1.77 (m, 1 H), 0.99 (m, 1 H), 0.88 (m, 1 H), 0.81 (d, J = 7.0 Hz, 3 H), 0.75 (t, J = 7.3 Hz, 3 H); ^{13}C NMR (100 MHz, CDCl_3) δ 175.0, 171.9, 137.1, 129.2, 128.4, 126.7, 59.8, 54.0, 38.4, 26.0, 23.4, 15.9, 11.8; HRMS-FAB ($\text{M} + \text{Na}^+$) calcd for $\text{C}_{16}\text{H}_{25}\text{N}_3\text{NaO}_2$ 314.1844, found 314.1850.

L-Isoleucyl-L-phenylglycine methyl amide (11e)

Compound **11e** (0.032 g, 0.115 mmol, 53%) was prepared from **10e** (0.036 g, 0.217 mmol), NHS (0.03 g, 0.26 mmol), *N*-Boc-L-isoleucine (0.062 g, 0.26 mmol), EDC (0.05 g, 0.26 mmol) and TFA (0.35 mL) according to the procedure used for preparing **11a**. ^1H NMR (400 MHz, CD_3OD) δ 7.44 (m, 2 H), 7.34 (m, 3 H), 5.42 (s, 1 H), 3.29 (d, J = 5.3 Hz, 1 H), 2.74 (s, 3 H), 1.77 (m, 1 H), 1.47 (m, 1 H), 1.15 (m, 1 H), 0.96 (d, J = 6.9 Hz, 3 H), 0.89 (t, J = 7.4 Hz, 3 H); ^{13}C NMR (100 MHz, CD_3OD) δ 175.1, 171.4, 137.7, 128.3, 127.9, 127.1, 59.2, 57.0, 38.8, 25.0, 23.9, 14.6, 10.6; HRMS-FAB ($\text{M} + \text{Na}^+$) calcd for $\text{C}_{15}\text{H}_{23}\text{N}_3\text{O}_2\text{Na}$ 300.1688, found 300.1680.

L-Isoleucyl-L-tryptophan methyl amide (11f)

Compound **11f** (0.103 g, 0.313 mmol, 55%) was prepared from **10f** (0.124 g, 0.573 mmol), NHS (0.079 g, 0.688 mmol), *N*-Boc-L-isoleucine (0.165 g, 0.688 mmol), EDC (0.132 g, 0.688 mmol) and TFA (1 mL) according to the procedure used to prepare **11a**. ^1H NMR (300 MHz, CD_3OD) δ 7.57 (d, J = 7.8 Hz, 1 H), 7.29 (dd, J = 8.1, 0.9 Hz, 1 H), 7.02 (m, 3 H), 4.59 (t, J = 6.5, 1 H), 3.2 (d, J = 6.5, 1 H), 2.62 (d, J = 1.9, 3 H), 1.60 (m, 1 H), 1.14 (m, 1 H), 0.9 (m, 1 H), 0.75 (m, 6 H); ^{13}C NMR (75 MHz, CD_3OD) δ 174.3, 172.1, 135.6, 126.2, 122.1, 119.9, 117.2, 116.8, 109.7, 108.3, 58.3, 52.9, 37.2, 26.7, 23.8, 22.3, 13.4, 9.4; HRMS-ESI ($\text{M} + \text{H}^+$) calcd for $\text{C}_{18}\text{H}_{27}\text{N}_4\text{O}_2$ 331.2134, found 331.2138.

L-Isoleucyl-L-leucine methyl amide (11g)

Compound **11g** (0.219 g, 0.851 mmol, 43%) was prepared from **10g** (0.285 g, 1.98 mmol), NHS (0.273 g, 2.375 mmol), *N*-Boc-L-isoleucine (0.57 g, 2.375 mmol), EDC (0.455 g, 2.375 mmol) and TFA (2.5 mL) according to the procedure used to prepare **11a**. ^1H NMR (300 MHz, CD_3OD) δ 4.4 (t, J = 7.5, 1 H), 3.23 (d, J = 5.5, 1 H), 2.73 (s, 3 H), 1.61 (m, 5 H), 1.18 (m, 1 H), 0.94 (m, 12 H); ^{13}C NMR (75 MHz, CD_3OD) δ 175.9, 174.2, 59.9, 51.9, 41.2, 39.3, 25.4, 24.9, 24.4, 22.5, 21.1, 15.1, 11.0; HRMS-FAB ($\text{M} + \text{Na}^+$) calcd for $\text{C}_{13}\text{H}_{27}\text{N}_3\text{O}_2\text{Na}$ 280.2001, found 280.2006.

L-Leucyl-L-tert-leucine methyl amide (11h)

Compound **11h** (0.054 g, 0.208 mmol, 65%) was prepared from **10h** (0.046 g, 0.322 mmol), NHS (0.045 g, 0.386 mmol), *N*-Boc-L-leucine (0.096 g, 0.386 mmol), EDC (0.074 g, 0.386 mmol) and TFA (0.75 mL) according to the procedure that was used to prepare **11a**. ^1H NMR (400 MHz, CD_3OD) δ 4.22 (s, 1 H), 3.46 (t, J = 5.3, 1 H), 2.74 (s, 3 H), 1.76 (m, 1 H), 1.57 (m, 1 H), 1.41 (m, 1 H), 1.01 (m, 9 H), 0.97 (m, 6 H); ^{13}C NMR (100 MHz, CD_3OD) δ 176.8, 172.1, 60.8, 53.6, 44.5, 34.5, 26.2, 25.1, 24.9, 22.7, 21.2; HRMS-FAB ($\text{M} + \text{Na}^+$) calcd for $\text{C}_{13}\text{H}_{27}\text{N}_3\text{O}_2\text{Na}$ 280.2001, found 280.2010.

General Procedure for the Synthesis of Compounds 12a–h

To a stirred solution of **8** (1.0 eq) dissolved in MeOH, **11a–h** (1.0 eq) was added and the mixture was stirred at room temperature overnight. The solvent was evaporated and the product was purified by column chromatography (3:97 MeOH/ CH_2Cl_2).

3-(Hydroxymethylamino)-4-(L-isoleucyl-L-*tert*-leucine methyl amide)-3-cyclobutene-1,2-dione (12a)

Compound **12a** (0.03 g, 0.077 mmol, 71%) was prepared from **8** (0.022 g, 0.109 mmol) and **11a** (0.028 g, 0.109 mmol) according to the general procedure. ¹H NMR (300 MHz, CD₃OD) δ 4.58 (d, *J* = 8.7 Hz, 1 H), 4.23 (s, 1 H), 3.49 (s, 3 H), 2.73 (s, 3 H), 1.93 (m, 1 H), 1.62 (m, 1 H), 1.23 (m, 1 H), 0.99 (s, 9 H), 0.94 (m, 6 H); ¹³C NMR (75 MHz, CD₃OD) δ 179.5, 178.9, 172.0, 171.8, 166.5, 166.3, 62.4, 61.3, 40.2, 37.7, 34.2, 26.1, 24.9, 24.4, 14.5, 10.0; HRMS-ESI (M + Na⁺) calcd for C₁₈H₃₀N₄O₅Na 405.2114, found 405.2126.

3-(Hydroxymethylamino)-4-(L-isoleucyl-L-cyclohexylglycine methyl amide)-3-cyclobutene-1,2-dione (12b)

Compound **12b** (0.0106 g, 0.026 mmol, 11%) was prepared from **8** (0.038 g, 0.24 mmol) and **11b** (0.068 g, 0.24 mmol) as a white solid according to the general procedure. ¹H NMR (300 MHz, CD₃OD) δ 4.56 (d, *J* = 9 Hz, 1 H), 4.14 (d, *J* = 6 Hz, 1 H), 3.49 (s, 3 H), 2.74 (s, 3 H), 1.92 (m, 1 H), 1.67 (m, 7 H), 1.22 (m, 5 H), 0.94 (m, 7 H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ 183.8, 182.7, 171.7, 171.1, 169.1, 168.1, 61.3, 58.4, 40.7, 39.2, 39.1, 31.1, 30.0, 29.2, 26.9, 26.8, 26.7, 24.4, 15.8, 12.0; HRMS-FAB (M + Na⁺) calcd for C₂₀H₃₂N₄NaO₅ 431.2270, found 431.2261.

3-(Hydroxymethylamino)-4-(L-isoleucyl-O-methyl-L-tyrosine methyl amide)-3-cyclobutene-1,2-dione (12c)

Compound **12c** (0.054 g, 0.12 mmol, 76%) was prepared from **8** (0.032 g, 0.16 mmol) and **11c** (0.051 g, 0.16 mmol) according to the general procedure. ¹H NMR (400 MHz, CD₃OD) δ 7.11 (d, *J* = 8.5 Hz, 2 H), 6.77 (d, *J* = 8.6 Hz, 2 H), 4.60 (dd, *J* = 9.6, 5.5 Hz, 1 H), 4.41 (d, *J* = 8.7 Hz, 1 H), 3.75 (s, 3 H), 3.49 (s, 3 H), 3.06 (dd, *J* = 13.8, 5.5 Hz, 1 H), 2.81 (dd, *J* = 13.9, 9.7 Hz, 1 H), 2.73 (s, 3 H), 1.87 (m, 1 H), 1.53 (m, 1 H), 1.18 (m, 1 H), 0.90 (m, 6 H); ¹³C NMR (100 MHz, CD₃OD) δ 179.1, 178.5, 172.4, 171.3, 165.9, 165.7, 158.6, 129.8, 128.6, 113.4, 62.0, 54.5, 54.2, 40.0, 37.0, 36.7, 24.9, 24.2, 14.1, 9.5; HRMS-FAB (M + Na⁺) calcd for C₂₂H₃₀N₄O₆Na 469.2063, found 469.2078.

3-(Hydroxymethylamino)-4-(L-isoleucyl-L-phenylalanine methyl amide)-3-cyclobutene-1,2-dione (12d)

Compound **12d** (0.224 g, 0.54 mmol, 75%) was prepared from **8** (0.113 g, 0.72 mmol) and **11d** (0.210 g, 0.72 mmol) as a white solid according to the general procedure. ¹H NMR (300 MHz, CD₃OD) δ 7.21 (m, 5 H), 4.63 (dd, *J* = 9.6, 9.3 Hz, 1 H), 4.43 (d, *J* = 8.6 Hz, 1 H), 3.49 (s, 3 H), 3.12 (dd, *J* = 6.3, 5.3 Hz, 1 H), 2.88 (dd, *J* = 9.7, 9.3 Hz, 1 H), 2.71 (s, 3 H), 1.86 (m, 1 H), 1.52 (m, 1 H), 1.16 (m, 1 H), 0.90 (m, 6 H); ¹³C NMR (75 MHz, CD₃OD) δ 179.6, 178.9, 172.7, 171.7, 166.4, 166.2, 137.2, 129.2, 128.4, 126.7, 62.3, 54.9, 40.4, 38.1, 37.3, 25.4, 24.5, 14.5, 9.92; HRMS-FAB (M + Na⁺) calcd for C₂₁H₂₈N₄NaO₅ 439.1957, found 439.1968.

3-(Hydroxymethylamino)-4-(L-isoleucyl-L-phenylglycine methyl amide)-3-cyclobutene-1,2-dione (12e)

Compound **12e** (0.021 g, 0.053 mmol, 65%) was prepared from **8** (0.016 g, 0.08 mmol) and **11e** (0.023 g, 0.083 mmol) according to the general procedure. ¹H NMR (300 MHz, CD₃OD) δ 7.42 (m, 2 H), 7.35 (m, 3 H), 5.42 (s, 1 H), 4.6 (d, *J* = 6 Hz, 1 H), 3.47 (s, 3 H), 2.73 (s, 3 H), 1.96 (m, 1 H), 1.63 (m, 1 H), 1.26 (m, 1 H), 1.0 (m, 6 H); ¹³C NMR (75 MHz, CD₃OD) δ 179.1, 178.5, 171.1, 171.0, 166.1, 165.9, 137.3, 128.4, 128.0, 127.2, 61.8, 57.3, 40.0, 37.7, 25.0, 24.2, 14.1, 9.8; HRMS-FAB (M + Na⁺) calcd for C₂₀H₂₆N₄O₅Na 425.1801, found 425.1810.

3-(Hydroxymethylamino)-4-(L-isoleucyl-L-tryptophan methyl amide)-3-cyclobutene- 1,2-dione (12f)

Compound **12f** (0.057 g, 0.125 mmol, 78%) was prepared from **8** (0.032 g, 0.16 mmol) and **11f** (0.053 g, 0.16 mmol) according to the general procedure. ¹H NMR (400 MHz, CD₃OD) δ 7.57 (d, *J* = 7.8 Hz, 1 H), 7.31 (d, *J* = 8.1 Hz, 1 H), 7.07 (m, 2 H), 6.99 (m, 1 H), 4.66 (dd, *J* = 8.1, 6.4 Hz, 1 H), 4.46 (d, *J* = 8.2 Hz, 1 H), 3.47 (s, 3 H), 3.25 (dd, *J* = 14.6, 6.3 Hz, 1 H), 3.11 (dd, *J* = 14.5, 8.1 Hz, 1 H), 2.67 (s, 3 H), 1.86 (m, 1 H), 1.50 (m, 1 H), 1.13 (m, 1 H), 0.88 (m, 6 H); ¹³C NMR (100 MHz, CD₃OD) δ 179.1, 178.5, 172.8, 171.3, 166.0, 165.8, 136.6, 127.3, 123.3, 120.9, 118.4, 117.9, 110.9, 109.3, 62.0, 54.3, 40.0, 37.3, 27.8, 25.0, 24.1, 14.1, 9.7; HRMS-FAB (*M* + Na⁺) calcd for C₂₃H₂₉N₅O₅Na 478.2066, found 478.2060.

3-(Hydroxymethylamino)-4-(L-isoleucyl-L-leucine methyl amide)-3-cyclobutene-1,2- dione (12g)

Compound **12g** (0.051 g, 0.133 mmol, 82%) was prepared from **8** (0.032 g, 0.16 mmol) and **11g** (0.042 g, 0.163 mmol) according to the general procedure. ¹H NMR (400 MHz, CD₃OD) δ 4.57 (d, *J* = 8.1 Hz, 1 H), 4.41 (dd, *J* = 9.4, 5.4 Hz, 1 H), 3.50 (s, 3 H), 2.74 (s, 3 H), 1.96 (m, 1 H), 1.63 (m, 3 H), 1.53 (m, 1 H), 1.22 (m, 1 H), 0.94 (m, 12 H); ¹³C NMR (100 MHz, CD₃OD) δ 179.1, 178.6, 173.5, 171.6, 166.3, 165.9, 61.9, 51.9, 40.6, 40.1, 37.6, 25.0, 24.5, 24.3, 22.0, 20.7, 14.3, 9.9; HRMS-FAB (*M* + H⁺) calcd for C₁₈H₃₁N₄O₅ 383.2295, found 383.2304.

3-(Hydroxymethylamino)-4-(L-leucyl-L-tert-leucine methyl amide)-3-cyclobutene- 1,2-dione (12h)

Compound **12h** (0.028 g, 0.074 mmol, 74%) was prepared from **8** (0.02 g, 0.1 mmol) and **11h** (0.026 g, 0.1 mmol) according to the general procedure. ¹H NMR (400 MHz, DMSO-*d*₆) δ 4.76 (br s, 1 H), 4.2 (d, *J* = 9.6 Hz, 1 H), 3.38 (s, 3 H), 2.57 (d, *J* = 4.4 Hz, 3 H), 1.76 (m, 1 H), 1.62 (m, 1 H), 1.51 (m, 1 H), 0.87 (s, 15 H); ¹³C NMR (75 MHz, CD₃OD) δ 179.8, 178.9, 172.7, 171.9, 166.5, 166.3, 61.3, 56.6, 41.1, 40.4, 34.4, 26.1, 25.0, 24.9, 22.5, 20.8; HRMS-FAB (*M* + Na⁺) calcd for C₁₈H₃₀N₄O₅Na 405.2114, found 405.2105.

3-(L-Leucyl-L-tert-leucine methyl amide)-4-methoxy-3-cyclobutene-1,2-dione (13)

To a stirred solution of **2** (0.021 g, 0.15 mmol) in MeOH, **11h** (0.042 g, 0.165 mmol) was added and the mixture was heated at reflux overnight. The solvent was evaporated and the crude product was purified by column chromatography (3:97 MeOH/CH₂Cl₂) giving compound **13** (0.052 g, 0.136 mmol, 95%). ¹H NMR (300 MHz, CD₃OD) δ 4.40 (s, 4 H), 4.25 (d, *J* = 9 Hz, 1 H), 2.74 (s, 3 H), 1.68 (m, 3 H), 0.99 (d, *J* = 6 Hz, 15 H); ¹³C NMR (75 MHz, CD₃OD) δ 188.7, 187.9, 184.2, 183.7, 178.0, 176.9, 173.1, 172.6, 172.0, 171.9, 171.4, 60.8, 60.7, 59.9, 56.8, 56.1, 25.6, 24.6, 24.5, 22.1, 20.0; HRMS-ESI (*M* + Na⁺) calcd for C₁₈H₂₉N₃O₅Na 390.2005, found 390.2010.

3-Hydroxy-4-(L-leucyl-L-tert-leucine methyl amide)-3-cyclobutene-1,2-dione (14)

To a stirred solution of **13** (0.05 g, 0.136 mmol) in MeOH (3 mL), 0.15 N HCl (1 mL) was added and the mixture was heated at reflux overnight. The solvent was evaporated and the crude product was purified by column chromatography (3:97 MeOH/CH₂Cl₂) giving **14** (0.032 g, 0.091 mmol, 67%). ¹H NMR (300 MHz, CD₃OD) δ 4.22 (s, 1 H), 3.32 (s, 1 H), 2.74 (s, 3 H), 1.69 (m, 3 H), 0.96 (s, 15 H); ¹³C NMR (75 MHz, CD₃OD) δ 196.3, 187.8, 180.7, 172.5, 171.9, 61.3, 55.8, 40.6, 34.5, 26.1, 25.1, 24.8, 22.5, 20.9; HRMS-ESI (*M* - H⁺) calcd for C₁₇H₂₆N₃O₅ 352.1872, found 352.1880.

3-(Hydroxymethylamino)-2-(L-isoleucine methylester)-4-thioxo-2-cyclobuten-1-one (15)

Compound **6a** (0.100 g, 0.370 mmol) was dissolved in 2 mL of CH₂Cl₂ and allowed to stir while Lawesson's reagent (0.150 g, 0.370 mmol) was added. The reaction was monitored by TLC until all the starting material had been consumed. The solvent was then evaporated and the crude material was purified by column chromatography (5:95 MeOH/CH₂Cl₂) providing **15** as a yellow solid (0.062 g, 0.23 mmol, 62%). ¹H-NMR (300 MHz, CDCl₃) δ 9.02 (br s, 1 H), 8.19 (br s, 1 H), 5.93 (br s, 1 H), 3.76 (s, 3 H), 3.67 (s, 3 H), 2.20 (m, 1 H), 2.07 (m, 1 H), 1.48 (m, 1 H), 1.26 (m, 1 H), 1.00 (d, *J* = 6.8 Hz, 3 H), 0.93 (t, *J* = 7.3 Hz, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 205.4, 201.6, 170.9, 170.8, 170.6, 61.2, 52.9, 39.4, 31.6, 25.3, 15.8, 12.2; ESI-MS (*M* + *H*⁺) calcd for C₁₂H₁₉N₂O₃S₂ 271.1294, found 271.1287.

3-(Hydroxymethylamino)-2-(L-isoleucyl-L-phenylalanine methylamide)-4-thioxo-2-cyclobuten-1-one (16)

Compound **16** (0.093 g, 0.216 mmol, 71%) was prepared as a yellow solid according to the procedure used to prepare **15** starting from **12d** (0.127 g, 0.305 mmol) and Lawesson's reagent (0.123 g, 0.304 mmol). The crude material was purified by column chromatography (3:97 MeOH/CH₂Cl₂). ¹H-NMR (300 MHz, CD₃OD) 7.15 (m, 5 H), 5.56 (d, *J* = 7.6 Hz, 1 H), 4.90 (s, 3 H), 4.68 (m, 1 H), 3.63 (s, 2 H), 3.10 (m, 1 H), 2.72 (d, *J* = 0.6 Hz, 3 H), 1.91 (m, 1 H), 1.49 (m, 1 H), 1.17 (m, 1 H), 0.87 (m, 6 H); ¹³C NMR (75 MHz, CD₃OD) δ 206.7, 204.0, 172.7, 171.1, 170.7, 170.4, 137.1, 129.3, 128.4, 126.5, 60.8, 54.9, 38.3, 37.6, 30.4, 25.5, 24.4, 14.5, 10.3; ESI-MS (*M* + Na⁺) calcd for C₂₁H₂₈N₄O₄SNa 455.1729, found 455.1738.

3-(Hydroxymethylamino)-2-(L-isoleucyl-L-leucine methyl amide)-4-thioxo-2-cyclobuten-1-one (17)

Compound **17** was prepared as a yellow solid (0.013 g, 0.032 mmol, 59%) from **12g** (0.021 g, 0.055 mmol) and Lawesson's reagent (0.022 g, 0.055 mmol) according to the procedure used to prepare compound **15**. The crude material was purified by column chromatography (1.3:98.7 MeOH/CH₂Cl₂). ¹H NMR (300 MHz, CD₃OD) δ 5.67 (d, *J* = 6.8 Hz, 1 H), 4.39 (dd, *J* = 9.5, 5.2 Hz, 1 H), 3.63 (s, 3 H), 2.73 (s, 3 H), 2.00 (m, 1 H), 1.56 (m, 4 H), 1.24 (m, 1 H), 0.96 (m, 12 H); ¹³C NMR (75 MHz, CD₃OD) δ 206.6, 203.9, 173.5, 171.2, 170.9, 170.2, 60.5, 51.9, 40.6, 37.9, 29.9, 24.9, 24.5, 23.9, 22.0, 20.5, 14.2, 10.2; HRMS-FAB (*M* + Na⁺) calcd for C₁₈H₃₀N₄O₄SNa 421.1885, found 421.1880.

3-(Hydroxymethylamino)-2-(L-leucyl-L-tert-leucine methyl amide)-4-thioxo-2-cyclobuten-1-one (18)

Compound **18** was prepared as a yellow solid (0.014 g, 0.035 mmol, 41%) from **12h** (0.033 g, 0.086 mmol) and Lawesson's reagent (0.035 g, 0.086 mmol) according to the procedure used to prepare compound **15**. The crude material was purified by column chromatography (1.3:98.7 MeOH/CH₂Cl₂). ¹H NMR (400 MHz, CD₃OD) δ 5.92 (s, 1 H), 4.21 (s, 1 H), 3.62 (s, 3 H), 2.74 (s, 3 H), 1.72 (m, 3 H), 0.98 (m, 15 H); ¹³C NMR (100 MHz, CD₃OD) δ 206.1, 203.5, 171.7, 171.4, 170.7, 170.0, 61.2, 54.8, 40.7, 34.2, 30.2, 25.9, 24.8, 24.5, 22.2, 20.6; LRMS-FAB (*M* + Na⁺) calcd for C₁₈H₃₀N₄O₄SNa 421.0, found 421.1.

3,4-Di-(L-phenylalanine methyl ester)-3-cyclobutene-1,2-dione (19)

To a stirred solution of L-phenylalanine methyl ester HCl salt (0.646 g, 3.0 mmol) in MeOH (10 ml), KOH (0.168 g, 3.0 mmol) and compound **7** (0.256 g, 1.13 mmol) were added. Formation of a white precipitate was observed. The mixture was stirred for 3 h at room temperature, the solvent was evaporated, EtOAc was added and the solution was washed with H₂O. The organic layer was dried over MgSO₄, and the solvent was evaporated. The crude product was recrystallized from MeOH. The resulting white crystals were washed with cold MeOH and dried *in vacuo* to give **19** as a white solid (0.363 g, 0.831 mmol, 74%). ¹H NMR

(400 MHz, DMSO- d_6) δ 7.91 (br s, 1 H), 7.27 (m, 6 H), 7.14 (d, J = 6.4 Hz, 4 H), 5.00 (d, J = 6.1 Hz, 2 H), 3.70 (s, 6 H), 3.17 (dd, J = 13.5, 4.9 Hz, 2 H), 3.06 (m, 2 H); ^{13}C NMR (100 MHz, DMSO- d_6) δ 183.3, 171.6, 167.5, 136.4, 129.8, 128.9, 127.3, 57.2, 55.3, 52.8; HRMS-ESI ($M + H^+$) calcd for $\text{C}_{24}\text{H}_{25}\text{N}_2\text{O}_6$ 437.1713, found 437.1722.

2,3-Bis(L-phenylalanine methyl ester)-4-thioxo-2-cyclobuten-1-one (20)

To a stirred solution of **19** (0.112 g, 0.257 mmol) in dry CH_2Cl_2 (5 ml), Lawesson's reagent (0.021 g, 0.052 mmol) was added. After TLC analysis indicated that the reaction was complete, the crude product was purified by column chromatography (gradient of 0.25–1.0% MeOH/ CH_2Cl_2) to yield compound **20** as a yellow solid (0.038 g, 0.084 mmol, 40%). ^1H NMR (400 MHz, DMSO- d_6) δ 8.76 (d, J = 7.9 Hz, 1 H), 8.31 (d, J = 9.1 Hz, 1 H), 7.21 (m, 10 H), 6.12 (dd, J = 15.0, 6.3 Hz, 1 H), 5.01 (dd, J = 12.7, 6.7 Hz, 1 H), 3.73 (s, 3 H), 3.70 (s, 3 H), 3.18 (m, 4 H); ^{13}C NMR (100 MHz, DMSO- d_6) δ 204.0, 180.4, 172.0, 171.4, 170.8, 168.9, 136.0, 129.8, 129.7, 129.0, 128.9, 127.5, 127.4, 58.1, 55.8, 53.1, 52.9; HRMS-ESI ($M + H^+$) calcd for $\text{C}_{24}\text{H}_{25}\text{N}_2\text{O}_5\text{S}$ 453.1484, found 453.1490.

3-(Hydroxymethylamino)-2-(L-phenylalanine methyl ester)-4-thioxo-2-cyclobuten-1-one (21)

To a stirred solution of **6e** (0.171 g, 0.562 mmol) in dry CH_2Cl_2 (5 ml), Lawesson's reagent (0.227 g, 0.561 mmol) was added. After TLC analysis indicated that the reaction was complete, the solvent was evaporated and the crude product was purified by column chromatography (CH_2Cl_2) to yield compound **21** as a yellow solid (0.104 g, 0.325 mmol, 58%). ^1H NMR (400 MHz, DMSO- d_6) δ 8.87 (br s, 1 H), 8.56 (br s, 1 H), 7.24 (m, 5 H), 6.23 (t, J = 6.0, 1 H), 3.72 (s, 3 H), 3.50 (s, 3 H), 3.21 (m, 2 H); ^{13}C NMR (100 MHz, DMSO- d_6) δ 206.5, 203.2, 171.0, 170.7, 170.6, 135.7, 129.9, 129.0, 127.5, 56.3, 53.1, 30.8; HRMS-FAB ($M + \text{Na}^+$) calcd for $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_4\text{SNa}$ 343.0729, found 343.0740.

Procedure for the Assay of MMP-1

MMP-1, prepared from a culture medium of human rheumatoid synovial fibroblasts, was obtained from Calbiochem. The assays were based on the enzymatic hydrolysis of the peptide substrate Dnp-Pro-Cha-Gly-Cys(Me)-His-Ala-Lys(Nma)- NH_2 ,¹⁸ using the procedure described by Le Diguarher.¹⁹ This is a fluorescence quenching assay in which *N*-methylantranilic acid (Nma) is the fluorophore and dinitrophenyl (Dnp) is the quencher.

Pro-MMP-1 was dissolved in assay buffer (200 mM NaCl, 50 mM Tris, 5 mM CaCl_2 , 20 μM ZnSO_4 , 0.05% Brij 35, pH 7.6) at a concentration of 1.25 $\mu\text{g/mL}$. APMA solution (*p*-aminophenylmercuric acetate, 2 mM in 0.1 N NaOH) was prepared. Proenzyme activation was performed by mixing the proenzyme solution and the APMA solution in a 10:1 ratio. This reaction was incubated at 37 °C for 30 min and subsequently transferred to ice and then to a freezer. The substrate was dissolved in DMSO at a concentration of 2 mM, and then diluted to 0.2 mM with H_2O . Inhibitors were dissolved in a 10:90 DMSO/buffer solution. Fluorescence measurements were performed in 96-well plates by incubating assay buffer (76 μL), activated enzyme (4 μL), and the inhibitor solution (or buffer for the blank) (10 μL) at 37 °C for 30 min and then adding the substrate (10 μL) to the wells. The change in fluorescence was measured using excitation and emission wavelengths of 340 nm and 460 nm, respectively. IC_{50} values were calculated using the commercial graphing package Grafit (Erithacus Software Ltd.). Data was obtained for assays with at least 5 different concentrations, in duplicate, of each inhibitor.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank Mr. Jian Xie for helpful discussions. This research was supported by the NIH NIGMS (Grant R01 GM057327).

References

1. Whittaker M, Floyd CD, Brown P, Gearing AJH. *Chem Rev* 1999;99:2735–2776. [PubMed: 11749499]
2. Jones CB, Sane DC, Herrington DM. *Cardiovasc Res* 2003;59:812–823. [PubMed: 14553821]
3. Chakraborti S, Mandal M, Das S, Mandal A, Chakraborti T. *Mol Cell Biochem* 2003;253:269–285. [PubMed: 14619979]
4. Westermarck J, Kahari VM. *FASEB J* 1999;13:781–792. [PubMed: 10224222]
5. Fisher C, Gilbertson-Beadling S, Powers EA, Petzold G, Poorman R, Mitchell MA. *Dev Biol* 1994;162:499–510. [PubMed: 7512058]
6. Billingham RC, Dahlberg L, Ionescu M, Reiner A, Bourne R, Rorabeck C, Mitchell P, Hambor J, Diekmann O, Tschesche H, Chen J, Van Wart H, Poole AR. *J Clin Invest* 1997;7:1534–1545. [PubMed: 9119997]
7. Jackson C, Nguyen M, Arkell J, Sambrook P. *Inflamm Res* 2001;50:183–186. [PubMed: 11392606]
8. Lee W, Aitken S, Sodek J, McCulloch CA. *J Periodont Res* 1995;30:23–33. [PubMed: 7722844]
9. Liedtke W, Cannella B, Mazzaccaro RJ, Clements JM, Miller KM, Wucherpfennig KW, Gearing AJH, Raine CS. *Ann Neurol* 1998;44:35–46. [PubMed: 9667591]
10. Xie J, Comeau AB, Seto CT. *Org Lett* 2004;6:83–86. [PubMed: 14703356]
11. Kim CU, Misco PF. *Tetrahedron Lett* 1992;33:3961–3962.
12. Sato K, Seio K, Sekine M. *J Am Chem Soc* 2002;124:12715–12724. [PubMed: 12392419]
13. Shinada T, Nakagawa Y, Hayashi K, Corzo G, Nakajima T, Ohfuné Y. *Amino Acids* 2003;24:293–301. [PubMed: 12707812]
14. Sun L, Chiu D, Kowal D, Simon R, Smeyne M, Zukin RS, Olney J, Baudy R, Lin S. *J Pharmacol Exp Ther* 2004;310:563–570. [PubMed: 15075380]
15. Butera JA, Antane MM, Antane SA, Argentieri TM, Freeden C, Graceffa RF, Hirth BH, Jenkins D, Lennox JR, Matelan E, Norton NW, Quagliato D, Sheldon JH, Spinelli W, Warga D, Wojdan A, Woods M. *J Med Chem* 2000;43:1187–1202. [PubMed: 10737752]
16. Lovejoy B, Welch AR, Carr S, Luong C, Broka C, Hendricks T, Campbell JA, Walker KAM, Martin R, Van Wart H, Browner MF. *Nat Struct Biol* 1999;6:217–221. [PubMed: 10074939]
17. Lim NC, Morton MD, Jenkins HA, Bruckner C. *J Org Chem* 2003;68:9233–9241. [PubMed: 14629141]
18. Bickett DM, Green MD, Berman J, Dezube M, Howe AS, Brown PJ, Roth JT, McGeehan GM. *Anal Biochem* 1993;212:58–64. [PubMed: 8368516]
19. Le Diguarher T, Chollet AM, Bertrand M, Hennig P, Raimbaud E, Sabatini M, Guilbaud N, Pierre A, Tucker GC, Casara P. *J Med Chem* 2003;46:3840–3852. [PubMed: 12930146]
20. Borkakoti N, Winkler FK, Williams DH, D'Arcy A, Broadhurst MJ, Brown PA, Johnson WH, Murray EJ. *Nat Struct Biol* 1994;1:106–110. [PubMed: 7656013]
21. Solans X, Aguilo M, Gleizes A, Faus J, Julve M, Verdaguer M. *Inorg Chem* 1990;29:775–784.
22. An X-ray crystal structure of a hydroxamic acid/zinc complex shows that the bite angle in this complex is 81.1°. Ruf M, Weis K, Brasack I, Vahrenkamp H. *Inorg Chim Acta* 1996;250:271–281. For comparison, we used HyperChem 3D to calculate the bite angle in a zinc complex of the squaric acid/hydroxamic acid hybrids, which was calculated to be 117.9°
23. Reich R, Katz Y, Hadar R, Breuer E. *Clin Cancer Res* 2005;11:3925–3929. [PubMed: 15897594]
24. Puerta DT, Lewis JA, Cohen SM. *J Am Chem Soc* 2004;126:8388–8389. [PubMed: 15237990]
25. Kortylewicz ZP, Galaray RE. *J Med Chem* 1990;33:263–273. [PubMed: 2153207]
26. Reichard GA, Stengone C, Paliwal S, Mergelsberg I, Majmundar S, Wang C, Tiberi R, McPhail AT, Piwinski JJ, Shih NY. *Org Lett* 2003;5:4249–4251. [PubMed: 14601972]

FIGURE 1.
Resonance structures of squaric acid.

FIGURE 2. Known binding mode of hydroxamic acid **I** versus proposed binding mode by squaric acid derivative **II**.

FIGURE 3.
6-membered vs. 5-membered zinc chelation models.

FIGURE 4.

Changes that occur in the ^1H NMR spectra of compounds **19** and **6e** upon monothiation of the cyclobutenedione core.

SCHEME 1a.

^aReagents: (a) $CH(OCH_3)_3$, MeOH, Δ ; (b) $HONHR^1 \cdot HCl$, KOH, MeOH; (c) H_2NR^2 , MeOH.

SCHEME 2a.

^aReagents: (a) MeOH, KOH. ^bRacemic Nle was used to prepare **5b** and **6c**.

SCHEME 3a.

^aReagents: (a) $(\text{BuO})_3\text{CH}$, BuOH, reflux; (b) HONHMe·HCl, KOH, MeOH; (c) NH_2Me , *N*-hydroxysuccinimide (NHS), 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC), DMF; (d) 50% TFA, CH_2Cl_2 ; (e) NHS, *N*-Boc-Ile or *N*-Boc-Leu, EDC, DMF; (f) 50% TFA, CH_2Cl_2 ; (g) MeOH. Tle = *tert*-leucine, Chg = cyclohexylglycine, Phg = phenylglycine.

^bSee Table 3 for specific structures.

SCHEME 4a.

^aReagents: (a) H-Leu-Tle-NHMe, MeOH, reflux; (b) 0.15 N HCl, MeOH, reflux.

6a: R = HN-Ile-OMe	15: R = HN-Ile-OMe
12d: R = HN-Ile-Phe-NHMe	16: R = HN-Ile-Phe-NHMe
12g: R = HN-Ile-Leu-NHMe	17: R = HN-Ile-Leu-NHMe
12h: R = HN-Leu-Tle-NHMe	18: R = HN-Leu-Tle-NHMe

SCHEME 5.
Thionation of Cyclobutenediones

SCHEME 6a.

^aReagents: (a) L-Phe-OMe·HCl, KOH, MeOH; (b) Lawesson's reagent, CH₂Cl₂, 25 °C.

TABLE 1

Inhibition of MMP-1 by Compounds 4a–f

compound	R ¹	R ²	IC ₅₀ (mM) ^a
4a	H	-CH ₂ CH(CH ₃) ₂	1.8 ± 0.3
4b	H	-(CH ₂) ₅ CH ₃	1.0 ± 0.2
4c	Me	-(CH ₂) ₅ CH ₃	0.31 ± 0.03
4d	Me	-(CH ₂) ₄ CH ₃	0.31 ± 0.03
4e	cyclohexyl	-(CH ₂) ₅ CH ₃	>10
4f	benzyl	-(CH ₂) ₅ CH ₃	>10

^a All experiments were performed in duplicate.

TABLE 2

Inhibition of MMP-1 by Compounds 6a–g

compound	R ¹	R ³ = side chain of amino acid	IC ₅₀ (μM) ^a
6a	Me	Ile	190 ± 30
6b	-CH(CH ₃) ₂	Ile	210 ± 10
6c	Me	Nleb	280 ± 20
6d	Me	Leu	320 ± 20
6e	Me	Phe	330 ± 20
6f	Me	Trp	380 ± 50
6g	Me	Met	300 ± 50

^a All experiments were performed in duplicate.^b Racemic Nle was used to prepare compound **6c**.

TABLE 3

Inhibition of MMP-1 by Dipeptides 12 a–h^a

compound	R ³ =side chain of amino acid	R ⁴ =side chain of amino acid	IC ₅₀ (μM) ^a
12a	Ile	Tle	500 ± 100
12b	Ile	Chg	300 ± 100
12c	Ile	Tyr(Me)	280 ± 30
12d	Ile	Phe	270 ± 30
12e	Ile	Phg	260 ± 25
12f	Ile	Trp	210 ± 30
12g	Ile	Leu	200 ± 20
12h	Leu	Tle	95 ± 7

^a All experiments performed in duplicate.

TABLE 4

Inhibition of MMP-1 by Thiocarbonyl-Containing Inhibitors

$X = S \text{ or } O$

R	compound number for X = S	IC ₅₀ (μM) ^a	compound number for X = O	IC ₅₀ (μM)
HN-Ile-OMe	15	70 ± 8	6a	190 ± 30
HN-Ile-Phe-NHMe	16	15 ± 1	12d	270 ± 30
HN-Ile-Leu-NHMe	17	170 ± 40	12g	200 ± 20
HN-Leu-Tle-NHMe	18	96 ± 30	12h	95 ± 7

^a All experiments performed in duplicate.