Published in final edited form as:

J Org Chem. 2006 April 14; 71(8): 3285–3286. doi:10.1021/jo052333q.

Rapid Conversion of Hindered Arylsulfonates to Alkyl Chlorides with Retention of Configuration

Salvatore D. Lepore, Anjan K. Bhunia, Deboprosad Mondal, Pamela C. Cohn, and Craig Lefkowitz

Department of Chemistry, Florida Atlantic University, Boca Raton, FL 33431-0991, slepore@fau.edu

Abstract

Arylsulfonates of hindered secondary alcohols are converted to the corresponding alkyl chlorides very rapidly and in good yields in the presence of titanium tetrachloride at low temperatures. These reactions proceed with exclusive retention of configuration.

We are interested in the development of highly efficient leaving groups containing chelating units capable of attracting incoming nucleophiles. We envision that electrophiles containing sulfonate leaving groups could be rendered more reactive by modifying them to contain a metal chelating moiety.¹ In this chelated form, the nucleophilic portion of a salt or Lewis acid is localized near the electrophilic center thus decreasing the entropic barrier relative to intermolecular reactions. The chelated metal would also be expected to stabilize the developing negative charge on the oxygens of the sulfonate leaving group in the transition state.² We have recently demonstrated that metal halides react at greatly accelerated rates with arylsulfonate-based nucleophile assisting leaving groups (NALGs) that contain a polyether unit (including macrocyclic) attached to the aryl ring ortho to the sulphonate.³ In an attempt to further explore the generality of the NALG technology, we discovered a highly efficient method to convert NALG sulfonates of hindered alcohols to the corresponding chlorides using TiCl_4 at low temperatures.

While a related chlorination procedure with alkyl tosylates has been reported,⁴ in this note, we demonstrate, for the first time, the stereospecificity and utility of the sulfonate/ TiCl_4 technique in the synthesis of a variety of alkyl chlorides.⁵

A series of alcohols were reacted with aryl sulfonyl chlorides containing a diethylene glycol moiety to give sulfonate esters **1** (Table 1). In general, the treatment of sulfonate esters **1** with TiCl_4 in methylene chloride at -78°C led to high yields of the corresponding alkyl chlorides. In each case, the reaction was complete in less than 2 minutes. With 1,3-diphenyl-2-propanol, we have observed that its NALG and tosyl esters primarily give the elimination product in the

presence of metal chloride agents.⁶ To our knowledge this chloride has only been prepared by radical chlorodecarboxylative methods.⁷ However, upon treatment of the NALG ester with TiCl_4 , the respective chloride was obtained exclusively in 91% yield (entry 1).

With sulfonates of chiral secondary alcohols (Table 1, entries 2–4), the chloride products were obtained as single diastereomers with complete retention of configuration.⁸ Thus, the NALG sulfonates of highly hindered menthol and isomenthol were both converted to the corresponding chlorides of the same configuration in excellent yields by exposure to TiCl_4 for two minutes. Similarly, the NALG sulfonate of cholesterol (entry 4) was converted to the corresponding chloride almost instantaneously at -78°C . Proton NMR confirms the formation of one stereoisomer with retention of configuration

In the 2-adamantyl system, we observed a 90% conversion to the chloride with no side product arising from rearrangement of the adamantine nucleus (entry 5).⁹ Backside nucleophilic displacement of 2-adamantyl sulfonate is essentially precluded due to steric crowding,¹⁰ thus the chlorination reaction likely proceeds via a front-side $\text{S}_{\text{N}}\text{i}$ -type mechanism. While reactions involving the $\text{S}_{\text{N}}\text{i}$ mechanism are generally mediated by four-centered cyclic transition states,^{11,12} the chlorination of the various sulfonate substrates in Table 1 is one of the first examples of an $\text{S}_{\text{N}}\text{i}$ transition state stabilized by intramolecular chelation which may account for the very rapid conversion rates.³ Neopentyl NALG substrates also reacted very rapidly with TiCl_4 leading to the rearrangement products (Scheme 1).

In conclusion, we believe that the method described in this note offers an important alternative to existing techniques for the synthesis of secondary alkylchlorides with retention of configuration.

Experimental Section

Representative Procedure: preparation of 1,3-diphenylpropyl 2-(2,5-dioxoheptylcarboxy)-1-benzo-sulfonate and conversion to 2-chloro-1,3-diphenylpropane (Table 1, entry 1)

Step 1—A mixture of *o*-sulfobenzoic acid anhydride (1.23 g, 6.6 mmol) and phosphorous pentachloride (3.68 g, 13.2 mmol) was heated at 90°C for 6 h. The oil was allowed to cool, dissolved in ether, and rinsed with ice water to remove unreacted phosphorous pentachloride. The solvent was evaporated *in vacuo* leaving 2.1 g of crude oil. The crude oil (1.5 g, 6.2 mmol) was then dissolved in excess methoxyethoxyethanol (2.4 g, 20 mmol) and heated to 60°C for 20 h. The reaction mixture was purified by flash chromatography by eluting with a hexane/acetone (10% v/v) to yield 2,5-dioxoheptyl 2-(chlorosulfonyl)-benzoate (1.9 g, 95%): IR (neat) 1731 ($\text{C}=\text{O}$), 1353, 1196 (SO_2) cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 8.14 (dd, 1H, $J = 0.9$ and 7.74 Hz), 7.80–7.69 (m, 2H), 4.57–4.54 (m, 2H), 3.86–3.84 (m, 2H), 3.67–3.65 (m, 2H), 3.55–3.53 (m, 2H), 3.36 (s, 3H); ^{13}C NMR (100.75 MHz, CDCl_3) δ 166.0, 141.6, 135.4, 132.4, 131.6, 130.4, 129.2, 72.0, 70.6, 68.7, 65.9, 59.2; HRMS (EI) $[\text{M}+\text{H}]^+$: calcd for $\text{C}_{12}\text{H}_{16}\text{ClO}_6\text{S}$ 323.0356, found 323.0350.

Step 2—To a dichloromethane solution (50 mL) of 2,5-dioxoheptyl 2-(chlorosulfonyl)-benzoate (2.0 g, 6.2 mmol) was added DMAP (0.9 g, 7.4 mmol) under argon. The reaction mixture was cooled to 0°C followed by the addition of 1,3-diphenyl-2-propanol (2.6 g, 12.4 mmol). The resulting mixture was warmed to room temperature and stirred for 15 h. The solvent was evaporated *in vacuo* and the resulting crude oil was then purified by flash chromatography by eluting with hexane/acetone (30% v/v) to give 1,3-diphenylpropyl 2-(2,5-dioxoheptylcarboxy)-1-benzosulfonate as a gummy material (2.7 g, 88 %): IR (neat) 1730 ($\text{C}=\text{O}$), 1357, 1201 (SO_2) cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.51–7.43 (m 3H), 7.27–7.22 (m, 1H), 7.18–7.09 (m, 10H), 5.12 (p, 1H, $J = 6.25$ Hz), 4.54–4.52 (m, 2H), 3.82–3.79 (m, 2H), 3.63–3.61 (m, 2H), 3.54–3.51 (m, 2H), 3.35 (s, 3H), 3.01–2.91 (m, 4H); ^{13}C NMR (100.75 MHz,

CDCl_3) δ 167.2, 136.3, 134.3, 133.0, 130.8, 129.8, 129.1, 129.0, 128.6, 126.9, 86.3, 72.0, 70.6, 68.9, 65.5, 59.2, 40.7. HRMS (EI) $[\text{M}+\text{H}]^+$: calcd for $\text{C}_{27}\text{H}_{31}\text{O}_7\text{S}$ 499.1790, found 499.1813.

Step 3—To a dichloromethane solution (10 mL) of 1,3-diphenylpropyl 2-(2,5-dioxoheptylcarboxy)-1-benzosulfonate (200 mg, 0.46 mmol) cooled to -78°C was added TiCl_4 (100 mg, 0.91 mmol). The resulting mixture was stirred for 1 min, quenched with water (50 mL) and diluted with additional dichloromethane (50 mL). The organic phase was collected and passed through a silica gel plug followed by solvent evaporation *in vacuo* to give pure 2-chloro-1,3-diphenylpropane (98 mg, 0.42 mmol, 91 %).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank the National Institute of Mental Health (66963-01) and the Florida Center of Excellence in Marine Biotechnology (contribution P200412) for financial support. An acknowledgment is also made to the National Science Foundation Division of Undergraduate Education (0311369) for the 400 MHz NMR used in this study.

References

1. (a) Cacciapaglia R, Di Stefano S, Mandolini L. *J Org Chem* 2002;67:521. [PubMed: 11798326] b Cacciapaglia R, Mandolini L. *Pure and App Chem* 1993;65:533. c Alfimov MV, Gromov SP, Fedorov YuV, Fedorova OA, Vedernikov AI, Churakov AV, Kuz'mina LG, Howard JAK, Bossmann S, Braun A, Woerner M, Sears DF Jr, Saltiel J. *J Am Chem Soc* 1999;121:4992.
2. Gobbi A, Landini D, Maia A, Secci D. *J Org Chem* 1995;60:5954.
3. Lepore SD, Bhunia AK, Cohn PC. *J Org Chem* 2005;70:8117. [PubMed: 16277337]
4. Vizgert RV, Polyakova EV, Chervinski A. *Yu Zhurnal Obshchei Khimi* 1984;54:905.
5. Other complementary chlorination techniques have also proved successful with intransigent substrates. (a) Yasuda M, Yamasaki S, Onishi Y, Baba A. *J Amer Chem Soc* 2004;126:7186. [PubMed: 15186150] b Kozikowski AP, Lee J. *Tetrahedron Lett* 1988;29:3053.
6. Lepore SD, Bhunia AK, Cohn PC. unpublished results
7. (a) Okada K, Okamoto K, Oda M. *J Chem Soc Chem Comm* 1989;21:1636. b Barton DHR, Crich D, Motherwell WB. *Tetrahedron* 1985;41:3901.
8. The stereochemistry of all chloride products was established by comparison of their NMR spectra with those previously reported. For the chloride products in Table 1 see the following references: entry 1 (ref. 7), entries 2–4 (ref. 5b), entries 5 and 6 (ref. 5a).
9. Sinnott ML, Storesund HJ, Whiting MC. *Chem Commun* 1969:1000.
10. Kozikowski AP, Lee J. *Tet Lett* 1988;29:3053. and references cited therein
11. Lee CC, Clayton JW, Lee DG, Finlayson AJ. *Tetrahedron* 1962;18:1395. b Lewis ES, Herndon WC, Duffy DC. *J Am Chem Soc* 1961;83:1959.
12. Moss RA, Fu X, Tian J, Sauers R, Wipf P. *Org Lett* 2005;7:1371. [PubMed: 15787509]

Scheme 1.

Reaction of neopentyl NALG sulfonates with TiCl_4 resulting in rearrangement product.

Table 1Reaction of various sulphonate esters with TiCl_4 leading to alkyl chlorides

entry	ROH	% yield of chloride ^a
1		91 (2)
2		88 (2)
3		93 (2)
4		85 (1)
5		90 (1)
6		66 (1) ^b

^a Isolated yields. All chloride products have been previously reported^b Elimination product also observed (21%, 1.5:1 *E/Z*).