Published in final edited form as:

J Mol Biol. 2008 August 29; 381(2): 256–260. doi:10.1016/j.jmb.2008.06.032.

Millisecond Time-Resolved Changes Occurring in Ca^{2+} -regulated Myosin Filaments upon Relaxation

Fa-Qing Zhao and Roger Craig¹

Department of Cell Biology, University of Massachusetts Medical School, Worcester, MA 01655, USA

Summary

Contraction of many muscles is activated in part by the binding of Ca^{2+} to, or phosphorylation of, the myosin heads on the surface of the thick filaments. In relaxed muscle, the myosin heads are helically ordered and undergo minimal interaction with actin. On Ca^{2+} binding or phosphorylation, the head array becomes disordered, reflecting breakage of the head-head and other interactions that underlie the ordered structure. Loosening of the heads from the filament surface enables them to interact with actin filaments, bringing about contraction. On relaxation the heads return to their ordered positions on the filament backbone. In scallop striated adductor muscle, the disordering that takes place on Ca^{2+} binding occurs on the millisecond timescale, suggesting that it is a key element of muscle activation. Here we have studied the reverse process. Using time-resolved negative staining electron microscopy we show that the rate of re-ordering on removal of Ca^{2+} also occurs on the same physiological timescale. Direct observation of images, together with analysis of their Fourier transforms, shows that activated heads regain their axial ordering within 20 msec and become ordered in their final helical positions within 50 msec. This rapid reordering suggests that re-formation of the ordered structure, and the head-head and other interactions that underlie it, is a critical element of the relaxation process.

Keywords

myosin filament; regulation; muscle; relaxation; electron microscopy

Muscle contraction is generated by cyclic interaction of myosin heads on the thick filaments with actin subunits in the thin filaments. This interaction, and hence contraction, is regulated by Ca^{2+} , via the troponin-tropomyosin complex on the thin filaments (actin-linked regulation) or the myosin light chains on the thick filaments (myosin-linked regulation).^{1–6} The regulatory movements of tropomyosin in actin-linked regulation are well understood. Electron microscopy (EM) reveals that tropomyosin blocks myosin-binding sites on actin in relaxed muscle, and moves so that these sites become exposed upon activation.^{7, 8} Time-resolved x-ray diffraction studies of contracting muscle demonstrate that movement of tropomyosin from its blocking position on activation is rapid (~ msec timescale) and occurs before myosin heads bind to actin.⁹ This supports the view that tropomyosin movement, in response to calcium binding to troponin, is physiologically relevant and a prerequisite for head attachment.

¹Corresponding author, Phone: (508) 856 2474, Fax: (508) 856 6361, Email: roger.craig@umassmed.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The structural basis of myosin-linked regulation, which in many muscles occurs together with the troponin-tropomyosin system, is less well understood. Two types of myosin-linked regulation have been identified. In one, initiation of contraction requires binding of Ca^{2+} to the essential light chains, whereas the other requires phosphorylation of the regulatory light chains by Ca^{2+} -calmodulin activation of myosin light chain kinase.^{2, 4, 10, 11} In relaxed muscles the heads of regulated myosin filaments are helically ordered on the filament surface.^{12–17} This ordered structure involves intramolecular head-head and head-tail interactions together with intermolecular interactions between myosin heads of axially neighboring molecules along the helical tracks.¹⁸ These multiple weak interactions are thought to switch myosin off by inhibiting ATP product release and actin-binding.^{18, 19}

Electron microscopy of both negatively stained and frozen-hydrated specimens shows that activation of Ca^{2+} - and phosphorylation-regulated thick filaments causes the helically ordered array of myosin heads to become disordered, reflecting a loosening of myosin head interactions.^{16, 20–23} However, the slow timescale of these experiments (seconds to minutes) does not reveal whether these changes are rapid enough to be physiologically relevant. Time-resolved x-ray studies of myosin activation (comparable to those on tropomyosin⁹), which would test the timescale and therefore physiological relevance of disordering, have not been reported. We developed an alternative, electron microscopic approach – millisecond time-resolved negative staining²⁴ – to study the time course of disordering in Ca^{2+} -regulated thick filaments (from the scallop¹⁶). We showed that uranyl acetate, routinely used as a negative stain, fixes macromolecular structure in less than 10 msec, and can therefore be used to capture structural transients on the millisecond timescale. By placing stain and activating solution in series in a pipette (with an air gap separating the two), relaxed filaments on an EM grid can be briefly activated (10–70 msec) and then immediately stained (by rapidly flushing the grid with the contents of the pipette), thus arresting the activated structure at known msec time points for EM observation.^{16, 24} Using this procedure, we showed that Ca^{2+} -induced disordering of scallop filaments is fully developed within 30 msec,¹⁶ well within the 90 msec time for development of peak twitch tension.²⁵ As with tropomyosin movement,⁹ this timescale suggests that disordering is a key requirement of the physiological activation process. Disordering appears to result from breaking of the head-head and head-tail interactions that underlie the relaxed state.^{23, 26} The speed of disordering suggests that breaking of these interactions is also rapid, limited mainly by the kinetics of binding of Ca^{2+} to the heads.^{16, 27}

While these experiments provide insights into the time course of structural changes underlying myosin activation, we know little about the kinetics of the reverse process, when muscle relaxes. Our negative staining and cryo-EM studies have shown that removal of Ca^{2+} leads to the reappearance of helical order,^{16, 20} but as with activation, the slow time scale of these experiments leaves in doubt the physiological significance of this reversal. It is possible, for example, that re-ordering might occur by a slow ‘annealing’ process following dissociation of Ca^{2+} , and not be a requirement for switching off activity.

Here, we have tested the hypothesis that re-ordering occurs on the physiological timescale. We have done this using a variation on the time-resolved negative staining technique. By including a relaxing solution in the pipette, between the activating solution and the stain, filaments were first activated, then relaxed again for predetermined msec time periods prior to rapid fixing with stain.

Figures 1a and b show relaxed and activated control filaments, respectively. Filaments flushed with relaxing rinse for a range of times (30–70 msec) were helically ordered, as judged both by eye and by the presence of layer lines in their Fourier transforms (Fig. 1a). This demonstrates that the rapid flushing procedure used in the time-resolved negative staining technique does not cause disorder, for example by mechanical disruption.¹⁶ The layer lines reveal the helical

parameters of the scallop filament: the repeat of the main (7,3)^{12, 14} helical tracks is ~48 nm, while the axial spacing between neighboring ‘crowns’ of heads is ~14.5 nm (Fig. 1a).^{12, 14, 15} The layer line at 11.2 nm indicates the presence of mass oriented along the (7/13) helical tracks,^{12, 14} while the meridional reflection at 9.8 nm reflects a small perturbation at alternating 14.5 nm levels of heads.^{12, 14}

When filaments were flushed with activating rinse, they rapidly became disordered, with their heads spread away from the filament backbone, reaching maximal disorder within 30 msec (Fig. 1b).¹⁶ A weak meridional reflection at 14.5 nm remaining in the Fourier transform may represent vestiges of axial head ordering, or possibly visualization of the myosin axial periodicity in the filament backbone; similar residual reflections at 14.5 nm are seen in other species of filament with disordered heads (17 and unpublished data). Longer activation times (50–70 msec) showed similar disorder, confirming our earlier findings.¹⁶ We therefore used activation times from 30–70 msec for the reversal experiments.

To determine the rate of structural change following removal of Ca^{2+} , filaments were first activated (30–70 msec), then flushed with relaxing rinse (20–80 msec), and finally rapidly fixed with uranyl acetate. Within 20 msec, the heads had returned close to the filament backbone (Fig. 1c), forming a compact but still helically disordered structure, comparable to that seen at the earliest (10 msec) stage of activation.¹⁶ Signs of axial re-ordering on to the 14.5 nm repeat were visible at this stage, as judged from the appearance of the Fourier transform (Fig. 1c). After 30 msec relaxation, the 48 nm layer line reappeared, demonstrating reordering of the heads onto the main (7,3) helical tracks^{12, 14} (Fig. 1d). At 50 msec, additional ordering became apparent, with the reappearance of the reflections at ~11.2 nm and ~9.8 nm (Fig. 1e), similar to the relaxed controls (Fig. 1a). Longer activation and relaxation times (hundreds of milliseconds) were also studied, and gave similar results (data not shown).

Rapid removal of Ca^{2+} from briefly activated filaments simulates the process of relaxation following a twitch in live muscle. The reordering we observe occurs on the millisecond time scale, reaching a maximum well within the 100 msec taken for half relaxation from peak tension in intact muscle²⁵ (while our experiments were carried out at a temperature 10° C above the tension experiments, rapid reordering would be expected even at the lower temperature, assuming a Q10 of ~2 (Ref. 28). We conclude that re-ordering of the heads is a key physiological component of the relaxation process, rather than the result of slow annealing. The rate of re-ordering would be limited by the rate of Ca^{2+} loss from the Ca^{2+} -specific binding sites²⁹ on the essential light chains of the myosin heads. In the absence of nucleotide, this occurs with a half-time of ~13 msec.^{27, 30} Measurements in the presence of nucleotide have not been reported.

Intramolecular interaction between heads is a characteristic feature of the off-state of phosphorylation-regulated myosin molecules.^{19, 31, 32} These interactions create an asymmetric structure in which the activity of both heads is switched off, by the blocking of sites involved in actin-binding on one head and in ATP hydrolysis on the other. Similar head-head interactions are also present in the off-state of native, helically ordered myosin filaments (tarantula) regulated by phosphorylation.¹⁸ In native filaments, additional interactions, between the heads of axially neighboring myosin molecules along the helical tracks, help to stabilize the off-state.¹⁸ Similar head-head interactions have been demonstrated in Ca^{2+} -regulated (scallop33) myosin molecules and we have recently shown their presence in native scallop myosin filaments.³⁴ The rapidity with which these interactions are reestablished on Ca^{2+} removal from activated filaments (implied by the rapid helical re-ordering that we observe) suggests that their re-formation is integral to the relaxation process of Ca^{2+} -regulated filaments. Our results suggest that this occurs in stages. The initial (within 20 msec) return of the heads to the filament backbone, forming a ‘compact’ structure with a degree of axial

ordering (Fig. 1c), may result from re-formation of the intramolecular head-head interactions. Following this, the heads may reestablish the main intermolecular interactions between adjacent axial levels along the 48 nm tracks, thus re-forming the (7,3) helices.^{12, 14} The subsequent reappearance of the 11.2 and 9.8 nm reflections implies re-formation of the (7,13) helical tracks and the axial perturbation.¹⁴ Three-dimensional reconstruction of scallop filaments³⁴ suggests that the 11.2 nm reflection is due largely to the mass of the interacting motor domains lying along the (7,13) helical tracks. The return of this reflection might thus represent the final docking of the interacting head motor domains back onto the filament, possibly by binding to the associated S2.^{18, 31–33}

Effective relaxation is critical to normal muscle function. Re-ordering of heads near the filament backbone could contribute to the efficiency of this process by rapidly sequestering them away from the thin filaments (in addition to switching off their activity by the head-head and other interactions). This would be expected to minimize resistance to stretch that occurs when relaxed muscles return to their rest length. It may also reduce the work done by active fibers in shortening a muscle at low load in which unstimulated fibers are shortened passively. It could thus play an important role in the energetics and mechanics of relaxation.¹⁷

Acknowledgments

We thank Drs. Maria Elena Zoghbi and John Woodhead for advice regarding averaging of Fourier transforms, Drs. Woodhead, HyunSuk Jung and Raúl Padrón for comments on the manuscript, and one of the referees for valuable suggestions regarding interpretation of our data. This work was supported in part by NIH grant AR34711. Electron microscopy was carried out in the Core Electron Microscopy Facility of the University of Massachusetts Medical School, supported in part by Diabetes Endocrinology Research Center grant DK32520.

References

1. Ebashi S, Endo M, Ohtsuki I. Control of muscle contraction. *Quart. Rev. Biophys* 1969;2:351–384.
2. Kendrick-Jones J, Lehman W, Szent-Györgyi AG. Regulation in molluscan muscles. *J. Mol. Biol* 1970;54:313–326. [PubMed: 4250215]
3. Lehman W, Szent-Györgyi AG. Regulation of muscular contraction. Distribution of actin control and myosin control in the animal kingdom. *J. Gen. Physiol* 1975;66:1–30. [PubMed: 125778]
4. Szent-Györgyi AG, Kalabokis VN, Perreault-Micale CL. Regulation by molluscan myosins. *Mol. Cell Biochem* 1999;190:55–62. [PubMed: 10098969]
5. Gordon AM, Homsher E, Regnier M. Regulation of contraction in striated muscle. *Physiol. Rev* 2000;80:853–924. [PubMed: 10747208]
6. Perry, SV. Activation of the contractile mechanism by calcium. In: Engel, AG.; Franzini-Armstrong, C., editors. *Myology*. New York, NY: McGraw-Hill; 2004. p. 281–306.
7. Lehman W, Craig R, Vibert P. Ca^{2+} -induced tropomyosin movement in *Limulus* thin filaments revealed by three-dimensional reconstruction. *Nature* 1994;368:65–67. [PubMed: 8107884]
8. Vibert P, Craig R, Lehman W. Steric-model for activation of muscle thin filaments. *J. Mol. Biol* 1997;266:8–14. [PubMed: 9054965]
9. Kress M, Huxley HE, Faruqi AR, Hendrix J. Structural changes during activation of frog muscle studied by time-resolved X-ray diffraction. *J. Mol. Biol* 1986;188:325–342. [PubMed: 3735425]
10. Sellers JR. Phosphorylation-dependent regulation of *Limulus* myosin. *J. Biol. Chem* 1981;256:9274–9278. [PubMed: 6114959]
11. Sellers, JR. *Myosins*. New York: Oxford University Press; 1999.
12. Wray JS, Vibert PJ, Cohen C. Diversity of cross-bridge configurations in invertebrate muscles. *Nature* 1975;257:561–564. [PubMed: 1165781]
13. Stewart M, Kensler RW, Levine RJ. Structure of *Limulus* telson muscle thick filaments. *J. Mol. Biol* 1981;153:781–790. [PubMed: 7338924]
14. Vibert P, Craig R. Electron microscopy and image analysis of myosin filaments from scallop striated muscle. *J. Mol. Biol* 1983;165:303–320. [PubMed: 6682452]

15. Vibert P. Helical reconstruction of frozen-hydrated scallop myosin filaments. *J. Mol. Biol* 1992;223:661–671. [PubMed: 1542113]
16. Zhao FQ, Craig R. Ca^{2+} causes release of myosin heads from the thick filament surface on the milliseconds time scale. *J. Mol. Biol* 2003;327:145–158. [PubMed: 12614614]
17. Zoghbi ME, Woodhead JL, Craig R, Padrón R. Helical order in tarantula thick filaments requires the "closed" conformation of the myosin head. *J. Mol. Biol* 2004;342:1223–1236. [PubMed: 15351647]
18. Woodhead JL, Zhao FQ, Craig R, Egelman EH, Alamo L, Padrón R. Atomic model of a myosin filament in the relaxed state. *Nature* 2005;436:1195–1199. [PubMed: 16121187]
19. Wendt T, Taylor D, Trybus KM, Taylor K. Three-dimensional image reconstruction of dephosphorylated smooth muscle heavy meromyosin reveals asymmetry in the interaction between myosin heads and placement of subfragment 2. *Proc. Natl. Acad. Sci. U. S. A* 2001;98:4361–4366. [PubMed: 11287639]
20. Vibert P, Craig R. Structural changes that occur in scallop myosin filaments upon activation. *J. Cell Biol* 1985;101:830–837. [PubMed: 4040918]
21. Craig R, Padrón R, Kendrick-Jones J. Structural changes accompanying phosphorylation of tarantula muscle myosin filaments. *J. Cell Biol* 1987;105:1319–1327. [PubMed: 2958483]
22. Levine RJ, Chantler PD, Kensler RW, Woodhead JL. Effects of phosphorylation by myosin light chain kinase on the structure of *Limulus* thick filaments. *J. Cell Biol* 1991;113:563–572. [PubMed: 2016336]
23. Stafford WF, Jacobsen MP, Woodhead J, Craig R, O'Neill-Hennessey E, Szent-Györgyi AG. Calcium-dependent structural changes in scallop heavy meromyosin. *J. Mol. Biol* 2001;307:137–147. [PubMed: 11243809]
24. Zhao FQ, Craig R. Capturing time-resolved changes in molecular structure by negative staining. *J. Struct. Biol* 2003;141:43–52. [PubMed: 12576019]
25. Rall JA. Mechanics and energetics of contraction in striated muscle of the sea scallop, *Placopecten magellanicus*. *J. Physiol. (Lond)* 1981;321:287–295. [PubMed: 6978395]
26. Azzu V, Yadin D, Patel H, Fraternali F, Chantler PD, Molloy JE. Calcium regulates scallop muscle by changing myosin flexibility. *Eur. Biophys. J* 2006;35:302–312. [PubMed: 16404592]
27. Jackson AP, Bagshaw CR. Transient-kinetic studies of the adenosine triphosphatase activity of scallop heavy meromyosin. *Biochem. J* 1988;251:515–526. [PubMed: 2969725]
28. Hou TT, Johnson JD, Rall JA. Effect of temperature on relaxation rate and Ca^{2+} , Mg^{2+} dissociation rates from parvalbumin of frog muscle fibres. *J. Physiol* 1992;449:399–410. [PubMed: 1522515]
29. Xie X, Harrison DH, Schlichting I, Sweet RM, Kalabokis VN, Szent-Györgyi AG, Cohen C. Structure of the regulatory domain of scallop myosin at 2.8 Å resolution. *Nature* 1994;368:306–312. [PubMed: 8127365]
30. Ankrett RJ, Walmsley AR, Bagshaw CR. Kinetic analysis of regulated myosin ATPase activity using single and limited turnover assays. *J. Cell Sci. Suppl* 1991;14:1–5. [PubMed: 1832160]
31. Burgess SA, Yu S, Walker ML, Hawkins RJ, Chalovich JM, Knight PJ. Structures of smooth muscle myosin and heavy meromyosin in the folded, shutdown state. *J. Mol. Biol* 2007;372:1165–1178. [PubMed: 17707861]
32. Jung HS, Craig R. Head-head and head-tail interaction: A general mechanism for switching off myosin II activity in cells. *Mol. Biol. Cell*. 2008In press
33. Jung HS, Burgess SA, Billington N, Colegrave M, Patel H, Chalovich JM, Chantler PD. Conservation of the regulated structure of folded myosin 2 in species separated by at least 600 million years of independent evolution. *Proc. Natl. Acad. Sci. U. S. A* 2008;105:6022–6026. [PubMed: 18413616]
34. Zhao FQ, Woodhead JL, Craig R. Head-head interaction characterizes the relaxed state of scallop and *Limulus* muscle myosin filaments. *Biophysical Society Meeting Abstracts. Biophys. J*. 2008Abstract, 630-Pos
35. Perrin DD, Sayce IG. Computer calculation of equilibrium concentrations in mixtures of metal ions and complexing species. *Talanta* 1967;14:833–842.

Figure 1. Time-course of head re-ordering on thick filaments following Ca^{2+} removal

Each panel shows a field of negatively stained scallop thick filaments, an oscilloscope trace (left inset) showing time in activating and/or relaxing rinse, and an averaged Fourier transform of 20 representative filaments at right (numbers indicate spacing of main ~ 48 nm helical repeat (7,3) helices, axial spacing of head crowns (~ 14.5 nm), 11.2 nm spacing of the (7,13) helices, and 9.8 nm reflection coming from head perturbation (cf.12, 14). (a) **relaxed control**: grid rinsed for 70 msec with relaxing rinse, then stained; (b) **activated control**: 30 msec activating rinse then stained; (c) **20 msec reversal**: 30 msec activating rinse, 20 msec reversing rinse, then stained; (d) **30 msec reversal**: 60 msec activating rinse, 30 msec reversing rinse, then stained; (e) **50 msec reversal**: 60 msec activating rinse, 50 msec reversing rinse, then stained. Scale bar = 200 nm. **Methods**: Scallop (*Placopecten magellanicus*) thick filaments were prepared from skinned striated adductor muscle and observed by time-resolved negative staining using solutions and techniques described previously.^{16, 24} Relaxing rinse was 100 mM NaAc, 3

mM MgCl₂, 0.2 mM EGTA, 1 mM MgATP, 2 mM imidazole, 1 mM NaN₃ (pH 7.0). Activating rinse was the same, with CaCl₂ added to 0.4 mM total concentration (pCa ~ 4.0³⁵). Reversing rinse was relaxing rinse containing total 4 mM EGTA. Experiments were carried out at ~ 21° C. Time-resolved relaxation following activation was achieved by a variation of the technique previously described.^{16, 24} Three solutions (instead of two) were placed in the pipette tip, separated by air gaps. The first activated the filaments for 30–70 msec, producing similar disordering in all cases; the second was reversing rinse, which was applied for the next 20, 30, 50, 70 or 80 msec following activation to obtain ‘snapshots’ of the relaxation process. Finally, the structure present following these different relaxation periods was ‘instantaneously’ arrested by the third solution, 1% (w/v) uranyl acetate, which also served as a negative stain.²⁴ Most experiments were carried out by first making 3–5 grids each of the relaxed and activated controls and then 3–5 grids for the reversal. To minimize possible variation between grids, we also used the following sequence: one grid relaxed control, one grid activated control then one grid reversal, the whole sequence being repeated three times. The results were the same with both approaches. Control experiments showed that the changes in order were not due to the minor changes in pH (due to proton liberation on binding of Ca²⁺ to EGTA) or ionic strength occurring as the filaments were flushed with the different solutions. Grids were examined at 80 kV in a Philips CM 10 electron microscope (FEI, Hillsboro, OR). Conventional dose micrographs were recorded on Kodak 4489 electron microscope film. Fourier transforms of 20 representative filaments in each case were computed and averaged using ImageJ (v1.34s, NIH, Bethesda, MD). All averaged transforms shown were brightness- and contrast-adjusted to show a clear signal; the adjustments for each condition were identical so that they could be objectively compared.