Published in final edited form as:

Inorg Chem. 2007 December 24; 46(26): 10968–10980. doi:10.1021/ic700555y.

Adventures in Bioinorganic Chemistry

Thomas G. Spiro

Chemistry Department, Princeton University, Princeton, N.J. 08544

Prologue

Preparing an acceptance talk for the ACS Award for Distinguished Service in the Advancement of Inorganic Chemistry provided me opportunity to reflect on the human side of my professional journey. Subsequently I accepted the Editor's kind invitation to publish these reflections, in hopes that they will amuse my colleagues, and that they may even hold some meaning for those starting out on their careers.

From Chemistry Set to Scandinavia

I liked chemistry from the moment my parents bought me a chemistry set when I was twelve. I had a great time dissolving and mixing the metal salts, watching the resulting colors and smelling the smells. My little sister was terrified that I would blow up the house, but somehow we survived. This introduction to chemistry was reinforced by Mr. Nicklin, my enthusiastic high school chemistry teacher. So when I enrolled at UCLA, I majored in chemistry, though I did cast about in social science courses. I also considered a career in science journalism, after writing for the college paper, the UCLA Daily Bruin, and serving as its managing editor. However, the chemistry faculty were very supportive, and I decided to continue for the Ph.D.

My favorite undergrad courses had been in analytical chemistry. So, when I arrived at MIT for graduate work, in 1956, I joined the analytical division, headed by David Hume and Lockhart (Buck) Rogers. Hume was a great thesis advisor. He left me to do pretty much what I wanted, but was always available with thoughtful advice.

Research in the Hume lab focused on the measurement of metal complex formation constants. I chose mixed halide complexes of mercury^{1, 2} for my thesis. Outside the lab, I took key courses in vibrational spectroscopy from Dick Lord, and in group theory from Al Cotton, then a young turk on the faculty. Cotton's course gave me my first taste of the hot new field of ligand field theory. When I finished my thesis, I won a Fulbright scholarship to study in Denmark. Hume and Cotton helped get me a spot with Carl Ballhausen, who had just returned from Harvard to become the youngest full professor in the Danish academic world.

In 1960 I set sail for Copenhagen with my bride of a year, Helen Handin (one still crossed the Atlantic by ship in those days). When Ballhausen showed me to my desk, my new office mate, also just arrived from the U.S., extended his hand and said, 'Hi, I'm Harry Gray'. And so began a life-long friendship.

Harry and I read Ballhausen's classic book, 'Ligand Field Theory' in manuscript, and Harry went on to apply the theory to a host of interesting compounds. But, after making a trigonal Ni(II) complex and analyzing its spectrum,³ I returned to my roots in complexation constants. Following a year working for Standard Oil in Southern California, where I could be near my cancer-stricken father, I returned to Scandinavia, this time to the Stockholm lab of Lars Gunnar Sillen, the capital of complex formation measurements at the time. There I determined hydrolysis constants for Tl(III) chloride complexes.⁴

Princeton and Raman

From Stockholm, I landed a job at Princeton. Helen and I moved there in 1963, and our sons Peter and Michael were born soon after. I was hired as an instructor in analytical chemistry. Princeton had a distinguished analytical tradition, established by Howell Furman and Clark Bricker, but lost interest in analytical chemistry in the 60's, as did many other schools. Luckily for me, I could by then pass as an inorganic chemist.

The Tl(III) chloride complexes I had been studying in Stockholm had irregular stability constants, and I was curious about their structures. I remembered from Lord's course at MIT that Raman spectroscopy was a useful structural tool in aqueous solution. Donald Hornig, my department chair, had just gone to Washington as Lyndon Johnson's science advisor, and left behind a Cary 81 Raman spectrometer, in the care of his graduate student, Steve Kittleberger. The Cary 81 was the pre-laser Cadillac of Raman spectroscopy, and occupied a 5-foot high metal cabinet (Figure 1). Steve showed me how to load my Tl(III) chloride solution into a horn-shaped tube, slip it horizontally into the coils of the Hg-arc lamp, and then light the lamp with a Tesla coil. His jaw dropped, as a beautiful spectrum came rolling out of the strip-chart recorder (Figure 1).⁵ I had inadvertently chosen one of the strongest Raman scatterers, while he had been toiling away at one of the weakest: solid HF.

This experience hooked me on Raman spectroscopy. My first grad students took advantage of the Hornig instrument to explore a number of metal complexes, including the polynuclear hydroxy complexes of Bi(III),⁶ Pb(II)⁷ and Me₃ Pt(IV),⁸ with which I had become familiar in Stockholm. Their spectra gave evidence of interaction among the metal ions, leading us to explore the vibrational spectroscopy of metal-metal bonds in a series of metal cluster compounds.⁹⁻¹³ Raman spectroscopy was well suited to this application because of the low frequencies and large polarizability derivatives of the metal-metal stretching vibrations.¹⁴

Iron Balls and Environmental Chemistry

My stay in Stockholm had been financed by a NIH post-doctoral fellowship, which I had won with an application that argued the need for understanding the nature of metal complexes in biological systems. My arguments at the time were naïve, but this topic did in fact become the main theme of my research career. To find out more, I attended the Gordon Conference on 'Metals in Biology' in 1964, in New Hampshire. On the first afternoon, I went for a swim in Squam Lake, and a big guy pulled himself up onto the raft beside me, stuck out his hand and said 'Hi, I'm Paul Saltman'. Thus began another lifelong friendship.

This was only the second 'Metals in Biology' conference and coherence was not its strong point. Paul characterized it as the 'EPR spectroscopists' talking with the 'pig feeders' i.e. nutritionists trying to make sense of the roles of trace metals in the diet. Paul was one of the latter. Upon discovering that I knew something about metal complexes, he thrust a paper in my hand, which showed that ferric salts in the presence of fructose were efficiently absorbed through the intestines. At the time only ferrous salts were thought to be taken up.

Paul wanted me to tell him what the ferric fructose complex was. Fresh from my stay in Stockholm, I advised him to measure the formation curve via potentiometry. He looked doubtful, but said 'If you're so smart come to my lab and do it yourself.' I accepted the invitation, and next summer I set off for the Saltman lab at USC (which provided a nice opportunity to introduce 8-month-old Peter to both sets of grandparents in Los Angeles). We set up potentiometers, prepared electrodes and solutions, adjusted the pH and waited for the potentials to settle down. But they never did. They drifted for hours, days. We had rediscovered ferric hydroxide colloid formation! The same chocolate brown solutions were formed if we left out the fructose, and just neutralized ferric salts with bicarbonate. What made these colloids

interesting, however, was that electron microscopy revealed a fairly uniform distribution of spherical particles, 70 Å across (Figure 2)¹⁵ This was the same size as the hydroxy-iron core of the iron storage protein, ferritin.

Could this be coincidence? We suggested that ferritin might form by templated assembly of the protein subunits around a naturally formed hydroxy-iron core. This hypothesis is almost certainly incorrect. The protein shell is very stable, and one can load and unload iron into it, most readily via oxidation and reduction. Still, our work stimulated a good deal of research on what Harry dubbed 'Saltman's balls.' The physiological mechanisms of iron uptake and release are still being worked out.

I spent several enjoyable summers in the Saltman lab, first at USC and later at UCSD, working on ferric hydroxide colloid stabilization.¹⁶ Recently, my interest in metal oxides has been revived by a fruitful environmental science collaboration on the remarkable propensity of many bacteria to coat themselves with nanoparticulate MnO₂, produced by Mn(II) oxidation. We are working on the mechanism of this process, which appears to involve a multi-copper oxidase.¹⁷

My interest in environmental science began in politically turbulent 1970, when my department chair asked if I could perhaps teach a course that was 'relevant'. The first Earth Day materialized, and issues like DDT, global warming and threats to the ozone layer began to seep into the news. I advertised a course in environmental chemistry, and invited the 20 or so enrollees to accompany me to the library and research the background of these subjects. The course attracted the interest of Bill Stigliani, then a postdoc in the department, who volunteered to help teach it. Bill went on to a career in environmental science, and he and I eventually co-authored a textbook, *Chemistry of the Environment*.¹⁸ I also became involved with environmental studies university-wide, and helped establish the Princeton Environmental Institute, which now includes the Center for Environmental Bioinorganic Chemistry.

Romance of Raman

Soon after arriving in Princeton I obtained a NIH grant to study biologically interesting complexes using Raman spectroscopy. The grant included \$20K to purchase a new Raman light source, a 20 mW He-Ne laser that was being marketed as an attachment to the Cary 81. Fortunately, faculty colleague George Leroi persuaded me that this was not a good buy, and that we should pool our resources to build a proper laser-Raman spectrometer, using the recently published design of Sergio Porto. This design, soon marketed by the Spex Corp., quickly swept the field, making the Cary 81 obsolete.

George and his students put together a spectrometer that worked well. The Porto design, taking advantage of the laser's directional property, allowed us to examine colored samples without losing all the light to absorption. This capability, and the availability of several visible wavelengths from the Ar⁺/Kr⁺ mixed gas laser that we bought, let us explore the resonance Raman (RR) effect, which amplifies the Raman cross section via coupling of vibrational and electronic transitions. This effect promised to greatly extend the utility of Raman spectroscopy, which normally required concentrated solutions because of low cross sections. There had been very few experimental studies of resonance enhancement, because of the pre-laser difficulties associated with colored samples. We felt like explorers on a new continent. Not only could we obtain RR spectra, we could study colored metallo-proteins because the spectrum was dominated by vibrational bands of the chromophore; there was no interference from the myriad vibrations of the protein itself, which were un-enhanced.

The quest was led by an early group of talented post-docs. One of the first was, Jim Nestor, a master experimentalist who really got our instruments working, then trained the rest of the

group in their proper use. He built an early CARS (coherent anti-Stokes Raman spectroscopy) instrument from scratch, publishing landmark papers on the application of resonance CARS.^{19, 20} The appeal of CARS was that the directional character of its signal eliminates interference from fluorescence, which is the great bane of Raman spectroscopy. The catch was that the CARS lineshapes are severely distorted (Figure 3), because of interference from an electronic background signal. Spectral interpretation required a complex analysis.²⁰ We abandoned the technique as too unwieldy, although others have persevered, and the advent of femtosecond techniques has given great impetus to CARS.

We were joined by Paul Stein, a talented theorist. Paul guided us through the interpretation of interesting phenomena that we were discovering, such as resonance de-enhancement,²¹ and antisymmetric Raman scattering.²²⁻²⁴

Vinny Mizkowski joined us from Harry's lab. Vinny had an incredible nose for inorganic spectroscopy. He correctly deduced that the intense blue color of 'blue' copper proteins, and the complex resonance Raman spectrum they produced, require a trigonal coordination geometry, with an equatorial cysteinate ligand, and weak axial ligation (Figure 4).²⁵ This work preceded the first crystal structure, by Hans Freeman, and the elegant spectroscopic analysis by Harry and by Ed Solomon.

We explored many colored metalloproteins. In addition to the 'blue' copper proteins, we studied the type III di-copper site, combining RR spectroscopy with EXAFS, a technique then just emerging at the Stanford synchrotron.²⁶⁻²⁹

Iron-sulfur proteins became a major focus. Greg Kubas, one of my first postdocs, took a stab at preparing a model compound, which turned out to have the non-biological but very interesting structure shown in Figure 5.³⁰ Later, when Dick Holm showed how to prepare proper models with sulfide and thiolate ligation, we compared their RR spectra with those of the proteins being modeled, to decipher protein-specific effects. Roman Czernuszewicz and Mike Johnson led this effort.³¹⁻³⁴

We studied cobalamins, as models for vitamin-B₁₂-containing enzymes.^{35, 36} In the early work, photolysis from the Raman laser precluded studies of the physiologically relevant alkyl cobalamins, but later we discovered how to circumvent this problem using frozen solutions, leading to mechanistic studies of important enzymes.^{37,38} There was a brief foray as well, into molybdenum enzymes.^{39, 40}

Eventually we went beyond metal centers, and investigated other biological chromophores, including flavins,^{41, 42} NAD,⁴³ and pyridoxal,⁴⁴ using various stratagems to circumvent fluorescence.

In a collaboration with biochemistry colleague Charles Gilvarg, we published the first Raman spectrum of a bacterial spore, which was dominated by the spore coat constituent calcium dipicolinate.⁴⁵ This spectrum has recently become the basis for promising approaches to detecting anthrax spores.

It was obvious that even bigger game lay in the ultraviolet region. Taking up the challenge, Rich Rava and Steve Fodor built an ultraviolet Raman spectrometer, based on a pulsed Nd:YAG laser, with a H₂ Raman-shifter, producing several ultraviolet wavelengths.⁴⁶ The laser was noisy and slow (10 pulses/s, each 10 ns long), but it worked well enough to explore the UVRR spectra of the nucleic acid bases, singly⁴⁷ and in DNA,⁴⁸ the aromatic amino acids, in and out of proteins.^{49, 50}

Work continued on nucleic acids⁵¹⁻⁵³ but proteins and the amide bond became our major UVRR focus. We studied the UVRR signatures of tyrosine,⁵⁴ of histidine,^{55, 56} and of proline.⁵⁷ Peptide studies were extended deep into the ultraviolet, where amide enhancement patterns, and the dependence on conformation could be worked out.⁵⁸⁻⁶⁰ UVRR was also applied to enzyme mechanisms in a steroid isomerase⁶¹ and in thymidylate synthase.⁶²

Porphyrins Proliferate

An Early Surprise: Anomalous Polarization

What really got RR spectroscopy off the ground was the discovery of anomalous polarization by Tom Strekas, which we reported in 1972.⁶³ Among the complexes proposed for study under my first NIH grant, I had included the recently discovered iridium dioxygen complexes, which were the first reversible non-biological O₂ adducts. This was to be Tom's thesis, but he soon found that the complexes decomposed in the laser beam before he could get a spectrum. The NIH proposal had indicated that we would study biological O₂ carriers once we had characterized the 'simpler' iridium complexes, and in desperation I suggested that Tom give hemoglobin (Hb) a try. This experiment too was a failure; no spectrum was forthcoming.

Just then Hb expert Chien Ho dropped by the lab. When we told him of our frustrations, he asked how we prepared our Hb. Tom pointed to a bottle of powdered Hb from Sigma. Chien shook his head, explained it was probably oxidized, and gave us the procedure for purifying Hb from whole blood. In an act of true scientific dedication, Tom went to our local hospital, had them draw a pint of his own blood, and purified his own Hb. When he put it in the laser beam, he was rewarded with a gorgeous RR spectrum, filled with numerous strong peaks.

We had no idea how to interpret the spectrum, having only a vague concept of what the resonance Raman effect was. However, Tom found that the peaks shifted when he de-oxygenated the Hb. So there was clearly chemical information to be deciphered. The big surprise was that some of the peaks had the *wrong* polarization. Every textbook told us that the intensity ratio for scattered light polarized parallel and perpendicular to the incident light was $\frac{3}{4}$ (depolarized) or less (polarized). Instead we had peaks with much higher ratios, most of the signal appearing in perpendicular polarization, cytochrome c showed the same effect (Figure 6). I told Tom he must be mistaken. But he checked the polarizers and stuck to his guns. Moreover, acting on an idea that the anomalously polarized peaks might arise from vibrations having rotational symmetry (A_{2g} in the idealized D_{4h} symmetry of the heme group), Tom worked out the rotational invariants from first principles, and showed that their depolarization ratio should indeed be infinite. These modes have antisymmetric Raman tensors and are normally forbidden, but become allowed in resonance; they are a specifically RR phenomenon. Our paper on this subject⁶³ created a stir, and brought RR spectroscopy a good deal of attention.

Porphyrin Systematics

The heme RR spectrum is very rich, and many students and postdocs were involved in sorting it out, from systematization of chemical effects with model hemes,^{64,65} to our first normal mode analysis,⁶⁶ to assignment of the important Fe-histidine stretching mode in Hb.⁶⁷ Band frequencies were correlated with porphyrin core size,^{68, 69} and we used isotopic substitution to unravel the contributions from vinyl substituents,⁷⁰ and from out-of-plane modes.⁷¹

Assignment of the numerous porphyrin vibrations was a continuing preoccupation. Jim Kincaid, Roman Czernuszewicz and Xiao-yuan Li systematically determined isotope shifts for a series of porphyrins, and Xiao-yuan did a heroic normal mode analysis in order to construct a consistent porphyrin force-field which accounted for all of these data, including the out-of-plane modes.^{72, 73} The force field was later refined further,⁷⁴ and theory at the MNDO/3

level was used to investigate the systematics of porphyrin structure and vibrational frequencies.⁷⁵

Soon after, however, the *ab initio* DFT era dawned, and we found that the normal modes of Ni-porphine⁷⁶ and NiTPP⁷⁷ were better reproduced by DFT than by our painfully refined empirical force-field. RR intensities were also computed⁷⁸ and we explored the important issue of out-of-plane mode enhancements resulting from porphyrin distortion. These modes became important in our subsequent work on the mechanism of metal insertion via porphyrin distortion in ferrochelatases.^{79,80}

Radicals, Excited States and Energy Conversion

Early on we found Martin Gouterman's four-orbital model of the porphyrin electronic structure to provide a satisfying explanation of the porphyrin resonance enhancement pattern,⁶³ and later of the frequency shifts and Jahn-Teller effects in porphyrin radical cation⁸¹ and anion^{82, 83} formation. Oxo-metal porphyrins, relevant to peroxidase intermediates were also characterized, in collaboration with colleague Jay Groves.⁸⁴ Later we studied RR spectra of porphyrin and chlorin excited states,⁸⁵⁻⁸⁷ with an eye to modeling photosynthetic reactions.⁸⁸

This work dovetailed with a developing project on photoelectrochemical solar energy conversion, using electrochemically polymerized films of vinyl-derivatized Ru (bipyridine)₃²⁺,⁸⁹ and of protoporphyrin.⁹⁰ Photochemical efficiencies were low for these films, and we decided to test the efficacy of covalently attached donor groups in extending the lifetime of charge transfer states in Co(II) porphyrins, hoping to generate H₂ via the transiently formed Co(I). However, picosecond spectral studies on these constructs revealed that the hoped-for charge transfer states were short-circuited by low-lying d-d states.⁹¹

Heme Protein Menagerie

We branched out into heme proteins other than Hb. Songzhou Hu carried out complete assignments of the complex RR spectra of cytochrome c⁹² and of myoglobin,⁹³ via isotopic labeling. Peroxidases were studied intensively⁹⁴⁻⁹⁶ especially by Giulietta Smulevich, who made them the base of her subsequent research career.

The peroxidases revealed complex patterns of Fe-CO and CO stretching modes in the carbonyl adducts, and we carried out many studies to understand the systematics of these indicator modes,^{97, 98} demonstrating that the observed variations in many heme proteins and models reflected backbonding changes associated with electrostatic influences in the heme pocket, and not geometric distortion as had been initially supposed. This led us directly into the longstanding controversy over whether proteins control the CO affinity by sterically hindering the upright binding of CO to the heme. Pawel Kozlowski was able to resolve a conflict between the crystallographers, who found modest bending of the FeCO unit, and the IR spectroscopists, who insisted that the FeCO was essentially linear. He showed via DFT calculations that the IR transition dipole was not collinear with the CO bond vector, and that the angle allowed by the IR measurements was actually consistent with the crystal structures (Figure 7).⁹⁹ Nevertheless, the energy required for modest bending was found to be low, and analysis of kinetic and equilibrium data showed that ligand discrimination by the distal histidine in myoglobin is almost entirely due to a stronger H-bond to O₂ than to CO.¹⁰⁰

Our attention has recently shifted to heme sensor proteins, in which ligand binding to heme controls enzyme activity or DNA-binding at another site in the protein. In the case of the NO sensor, soluble guanylyl cyclase, we helped establish that NO binding induces dissociation of the proximal histidine ligand.¹⁰¹ We also work out the mechanism and dynamics of CO

activation of the transcription factor CooA,^{102, 103} and the mechanism of CO inhibition of the heme-containing metabolic enzyme, cystathionine β -synthase.¹⁰⁴

A little heme RR lesson

Some background may help the reader appreciate why heme protein RR spectra are endlessly fascinating, and occasionally even useful.

Nature has done spectroscopists a great favor by choosing heme as a cofactor in myriad proteins. The iron-porphyrin complex absorbs visible light intensely, making heme proteins easy to monitor with absorption spectroscopy. In addition they provide high quality RR spectra, because resonance enhancement scales with the chromophore absorptivity. One can record strong RR signals at heme concentrations of 10 micromolar and below. Lasers are readily available in the 400 nm region, near resonance with the Soret electronic transition that gives rise to the intense absorption. And fluorescence, the bane of Raman spectroscopists, is efficiently quenched by the Fe ion, whose low-lying d-d transitions rapidly depopulate the excited state. Finally, the heme group has high symmetry, making vibrational assignments straightforward, despite the large number of atoms.

Small wonder, then, that the literature is replete with RR spectra of all sorts of heme proteins.

How resonance enhancement works for heme

But what information do these spectra provide?

The answer requires a brief excursion into Raman theory, and heme electronic structure.

When the exciting light is tuned to an electronic transition, certain Raman peaks are enhanced selectively. They arise from vibrational motions which shift the chromophore geometry towards that of the excited state.

Near an electronic resonance the Raman polarizability (proportional to the square root of the intensity) can be expressed as

$$\alpha = A + B \quad (1)$$

where

$$A = \mu_e^2 \sum_v F_v / (\Delta\nu_v + i\Gamma_v) \quad (2)$$

and

$$B = \mu_e \mu'_e \sum_v F'_v / (\Delta\nu_v + i\Gamma_v) \quad (3)$$

In these equations, $\Delta\nu_v$ is the detuning interval, the difference between the laser energy and the energy of a vibrational level, ν , with bandwidth Γ_v , in the electronic excited state. Resonance occurs as $\Delta\nu_v$ approaches zero. μ_e is the dipole moment of the electronic transition, and μ'_e is its derivative along the vibrational coordinate, Q , associated with the Raman band being monitored. For strong transitions, μ_e is much larger than $\mu_e \cdot \mu'_e$, and consequently the A term dominates the Raman spectrum. The B term becomes important in resonance with a weak transition, when μ'_e can become larger than $\mu_e \cdot \mu'_e$. μ'_e depends on mixing of the resonant transition, e , with an allowed transition, s (called vibronic or Hertzberg-Teller coupling),

$$\mu'_e = \mu_s \langle s | dH/dQ | e \rangle / (\nu_s - \nu_e) \quad (4)$$

and becomes large when the two electronic transitions are close in energy.

The vibrational selectivity of resonance enhancement is contained in the factors F_v and F'_v , which are products of vibrational wavefunctions (Franck-Condon integrals) between the ground and excited states. They are

$$F_v = \langle 1|v \rangle \langle v|0 \rangle \quad (5)$$

where 0 and 1 are the wavefunctions of the zeroth and first vibrational levels of the ground state, and v is any vibrational level of the excited state.

$$F'_v = \langle 1|Q|v \rangle \langle v|0 \rangle + \langle 1|v \rangle \langle v|Q|0 \rangle \quad (6)$$

A non-zero value for F_v requires the excited state potential surface to shift along the vibrational coordinate, and this is only possible for modes that carry the full symmetry of the molecule (totally symmetric modes). However, F'_v selects for modes that mix the e and s states, and these have symmetries contained in the direct product of the e and s transition symmetries. Thus resonance with strong electronic transitions is dominated by totally symmetric modes, but resonance with weak transitions can bring out non-totally symmetric modes.

The heme chromophore illustrates these principles beautifully. The visible region electronic transitions are based on π orbital excitations from close-lying HOMO's, a_{2u} and a_{1u} (in the idealized D_{4h} symmetry of the porphyrin ring), to a degenerate pair of LUMO's, e_g^* (see Figure 8). These two pairs of excitations give rise to two degenerate electronic transitions, E_u , which undergo configuration interaction. The transition dipoles add up for the higher transition, and nearly cancel for the lower one; they cancel exactly if the a_{1u} and a_{2u} have the same energy. This is Gouterman's four-orbital model, and it explains the characteristic porphyrin absorption spectrum, with its very strong ~ 400 nm band, called Soret or B, and its much weaker ~ 550 nm band, called α or Q. Some of the cancelled intensity (about 10 %) can be 'stolen' back from the B to the Q transition, via vibronic mixing, producing a ~ 500 nm vibronic side band, called β or Q_1 .

Gouterman's model also explains the intensity pattern of the heme RR spectra, which vary strongly with the laser wavelength. This variation allows a clear analysis of the porphyrin vibrations, despite the great complexity of the molecular structure. Even if we ignore (for the moment) the axial ligands and the peripheral substituents on the heme (counting only the C atoms attached directly to the ring - Figure 8), the heme group still has 37 atoms and 105 normal modes of vibration ($3N-6$). However, only nine of these are totally symmetric (A_{1g}). They are the ones that dominate RR spectra when the laser wavelength is near 400 nm, in resonance with the B electronic transition (Figure 9). Their depolarization ratio (intensity ratio for scattered light polarized perpendicular and parallel to the laser polarization) is $1/8$.

When the laser is tuned to the Q bands, entirely different modes become dominant (Figure 9). They are the vibronically active modes, the same ones that produce the Q_1 absorption intensity. Since the B and Q transitions both have E_u symmetry, the symmetries of the vibronic modes are given by

$$E_u \times E_u = A_{1g} + A_{2g} + B_{1g} + B_{2g} \quad (7)$$

However the A_{1g} vibrations are ineffective in vibronic mixing because of the high molecular symmetry, and the remaining modes dominate the Q-resonant spectra. Depolarization ratios are $3/4$ for B_{1g} and B_{2g} bands but infinite (*inverse* polarization) for A_{2g} bands. Inverse polarization is the unexpected effect discovered by Streckas in our first study of hemoglobin and cytochrome c (although it had been predicted in a footnote by Placzek in 1934!).⁶³ Textbooks had asserted that non-totally symmetric modes all have $3/4$ depolarization ratios, but this is clearly not true for A_{2g} modes. The reason is that they have rotational symmetry (R_z) and *anti*-symmetric polarizability tensors ($xy = -yx$). They literally rotate the plane of polarization, and all the intensity is found in perpendicular scattering. (In actual spectra some

parallel intensity is observed, because the symmetry is not fully D_{4h} . Depolarization ratios are nevertheless $>3/4$, and the bands are called *anomalously* polarized.)

A_{2g} mode activity is purely a resonance phenomenon. These bands disappear off-resonance. The reason is that the individual terms in the summation of eq. (3) change signs as the laser sweeps through $\Delta\nu_v = 0$, and successive terms can add or subtract. Because of tensor anti-symmetry adjacent terms add for A_{2g} modes when the laser is tuned between them, but subtract when the laser is tuned away from them. Thus the intensity cancels off-resonance. The opposite happens for B_{1g} and B_{2g} modes, whose tensors are symmetric ($xy = yx$). Their intensity is minimized between resonances but survives off-resonance. These opposite tendencies permit tuning the laser to selectively enhance A_{2g} or $B_{1g/2g}$ modes, as illustrated in Figure 9.

Because of these symmetry effects, the RR modes of the heme group have been thoroughly catalogued.^{72, 74} Symmetry greatly simplifies the spectra, and allows reliable mode identification. It also hides information, however. Many of the 105 modes do not appear in the RR spectra at all. Fifty-four of them are 'u' modes and are infrared, but not Raman active. They have been assigned from IR spectra of metalloporphyrins, but IR spectroscopy is severely limited for heme proteins by the strong IR absorption of water and of interfering protein modes. There are also out-of-plane modes (some 'u' and some 'g'), which are not resonance-enhanced, because they cannot couple to the resonant electronic transitions, which are polarized in the heme plane. However some 'u' modes, both in- and out-of-plane, do appear in heme RR spectra due to symmetry lowering effects.^{71-73, 106} These can arise from the intrinsic asymmetry of the porphyrin substituents, asymmetry in the axial ligands, out-of-plane distortions of the porphyrin ring, and electrostatic asymmetry of surrounding residues in proteins.

Porphyrin RR bands report Fe stereoelectronics

Although the heme RR bands arise from vibrations of the porphyrin ring, they nevertheless are sensitive to ligation changes at the central Fe. Early on it was recognized that frequency shifts in several modes were linked to changes in the oxidation and spin-state of the Fe.⁶⁴ The main mechanism for this chemical sensitivity turned out to stretching of the porphyrin bonds to accommodate changes in Fe size, and its displacement from the center of the ring.^{68, 69} The porphyrin ligand field strength is poised so that the Fe(II) or Fe(III) spin state is tuned by the axial ligands. In heme proteins, one of the axial ligands is usually imidazole (histidine side chain), although thiolate (cysteine), thioether (methionine) and amine (lysine or the protein N-terminus) are also encountered. If the sixth ligand is strong-field (cyanide, thioether or -for Fe (II)-CO, NO and O_2), the Fe is low-spin and the d electrons (5 and 6 for Fe(III) and Fe(II)) are all in π -bonding orbitals (t_{2g}). However, if the sixth ligand is weak (H_2O , F^-) or absent, the Fe is high-spin, and one electron occupies each of the σ -anti-bonding orbitals, lengthening the bonds, and expanding the porphyrin ring. If the sixth ligand is absent, the Fe is displaced toward the fifth ligand, partly relaxing the ring expansion.

The ring-size dependence of several porphyrin bands has been calibrated with metalloporphyrins of known structure,^{68, 69} and the data for heme proteins fall quite predictably on these correlations. Thus the positions of these 'spin marker' bands are routinely used to monitor the ligation status of the heme group.

The strongest band in Soret-resonant RR spectra, designated as ν_4 , was early recognized as a marker of the oxidation state, having characteristic narrow ranges for Fe(III) and Fe(II), regardless of spin-state.⁶⁴ The reasons for this behavior are complex, involving counterbalancing influences of ring size, and of π bonding. For CO, NO and O_2 adducts of Fe (II), the ν_4 position is close to that of Fe(III), because substantial charge is transferred to these π -acid ligands via back-bonding. Because ν_4 is intense, it serves as a useful monitor of ligation for these adducts.

Axial ligands report protein influences

Since the axial ligand bonds are perpendicular to the heme ring, their vibrations are not normally detectable in RR spectra because the resonant electronic transitions are polarized in the plane of the ring. However, there are important exceptions.

Because of back-bonding the vibrations of bound XO ($X = C, N, O$), are resonance enhanced. The ν^* orbitals on XO compete with those on the porphyrin ring (e_g^*) for back-donation of the Fe d_π electrons. Excitation of the porphyrin $\pi-\pi^*$ transitions weakens porphyrin but strengthens XO back-bonding. Consequently there is a shift of the excited state potential along Fe-X and X-O stretching coordinates.

The RR bands associated with Fe-X and X-O stretching have proved to be valuable probes of protein interactions with the heme, especially for CO adducts.¹⁰⁷ The extent of backdonation is sensitive to the electrostatic influence of polar residues in the heme pocket. Positive polarity increases backbonding, while negative polarity decreases it, with anti-correlated changes in the positions of ν_{FeX} and ν_{XO} ; increased backbonding strengthens the Fe-X bonds and weakens the X-O bonds. A series of myoglobin variants with distal residue replacements have provided a standard back-bonding correlation for CO adducts,¹⁰⁷ which has been much used in assessing distal influences in other proteins.

NO adducts have seemed not to follow this pattern,¹⁰⁸⁻¹¹⁰ but recent computational modeling has revealed that the complexity apparent in the NO data results from the ambidextrous nature of H-bonding to the bound NO.¹¹¹ Because Fe(II)NO is bent, with a partially filled non-bonding (sp^2) orbital on N, H-bonds can be stably formed with either N or O, despite the greater charge on O. $H\cdots O$ bonds produce increase backbonding, with anti-correlation of Fe-N and N-O bond strengths, but $H\cdots N$ bonds weaken both Fe-N and N-O bonds by drawing more electron density into the sp^2 orbital. Which end forms the H-bond depends on the geometry of the H-bond donor in the heme pocket, an effect that is apparent in the vibrational data.

Another important RR band is that associated with the bond to the proximal ligand, the only bonding connection between the heme and the protein.⁶⁷ Usually the associated vibration is not resonance-enhanced, but five-coordinate Fe(II) heme is an exception, and Fe-histidine RR bands have been reported for many heme proteins in this state. The displacement of Fe(II) toward the heme plane when this bond is stretched evidently shifts the excited state potential enough to produce appreciable enhancement. The position of this vibration reflects the strength of the Fe-histidine bond, which is variable among heme proteins, and is an important determinant of the heme chemistry.

There are many more strands in the heme protein RR fabric, which can be found in the cited literature. For now we return to the story of the Princeton Raman program.

The Time Dimension

When Woody Woodruff joined the group he had a burning ambition, which he has pursued throughout his research career, namely to add a time dimension to vibrational spectroscopy. He realized that harnessing the structural specificity of RR spectroscopy to kinetic studies would form a powerful combination.

At the time, we had no pulsed lasers to provide time resolution. So Woody constructed a flow system with a dual syringe pump, and aimed the Raman laser downstream of the mixing chamber to produce a spectrum of horseradish peroxidase compound II, the second intermediate of the peroxide reaction.¹¹² Soon we used Jim Nestor's pulsed-dye laser CARS apparatus to show that the Fe atom was displaced from the heme plane within 6 ns after photolysis of

the CO adduct of Hb.¹¹³ Later we pressed the time-resolution of this experiment into the picosecond regime using a rapid flow technique¹¹⁴ and subsequently, the HbCO photoproduct was characterized more fully with the 7 ns Nd:YAG laser that we acquired for UVRR studies.¹¹⁵

Truly time-resolved RR spectroscopy on the nanosecond time scale became possible with the acquisition of a second pulsed laser, which could be delayed electronically. We realized that UV excitation would enable us to study protein dynamics by spatial reference points, in the form of the aromatic residues. Initial attempts to do this on Hb led us astray;¹¹⁶ the signal/noise of our 10 Hz Nd:YAG lasers was just not up to the task.¹¹⁷ The situation improved dramatically when we traded up to 300 Hz XeCl excimer-dye lasers. Kent Rodgers implemented this system, and introduced key sampling improvements.¹¹⁸ The resulting difference spectra were sufficiently resolved (Figure 10) to allow structural interpretation, and subsequent studies produced the first structural model for the allosteric pathway connecting the R and T states.¹¹⁹⁻¹²¹ Complementary studies were carried out with a new time-resolved FTIR spectrometer.^{122,123}

The excimer lasers helped us produce important results, but they were difficult to operate, and broke down frequently. We switched to a solid state system, with Nd:YLF-pumped Ti:sapphire lasers, whose frequency could be shifted into the blue and ultraviolet regions via harmonic generation in mixing crystals. The lasers are stable and operate at 1 kHz repetition rates, producing spectra of superior quality.¹²⁴ We were able to sharpen the structural interpretation of the Hb time course, and found an important new intermediate on the pathway.¹²⁵ We examined site-directed mutants designed to test our pathway model, and came up with exciting new results, implicating dynamic cooperativity at the level of subunit dimers within the tetrameric structure.^{126, 127} Our old friend Hb continues to surprise and delight us.

Our new laser technology provides access to the deep UV, where amide bands of the protein backbone are optimally enhanced.¹²⁸ Bala Balakrishnan proposed that we look at protein folding via laser-induced temperature jumps. The T-jump technique allows access to the early events of protein folding and unfolding, and had been applied with fluorescence and IR monitoring of the relaxations, but UVRR spectroscopy promised multiple structural probes via the several amide and aromatic residue bands. Bala negotiated the design of a novel Nd:YLF-pumped OPO laser, operating at 1 KHz, to match our UV probe laser.¹²⁹ The system worked beautifully, and we have been applying it to a series of peptide and protein folding issues.¹³⁰⁻¹³² The technique is particularly promising as a way to characterize local unfolding events that may play an important physiological role in protein function.

Coda

This brief account has braided the main strands of the research program that I have been privileged to pursue. I have pushed and pulled stories of our many projects into a semblance of a narrative. Research, like life, is seldom straightforward, and my attempt to provide some coherence to the story required leaving out many side strands, some of them quite interesting in their own right, projects launched and, for a variety of reasons, not pursued very far. Likewise, although I have named a few of the players in our research drama, the narrative leaves out many individuals who contributed importantly to the program over the years; some of their work is cited in the references.

A research group has a life of its own, quite independent of the schemes and wishes of its director. I have been blessed with a wonderful collection of students, undergraduate, graduate, and postdoctoral, and of visitors. They have been the lifeblood of the program, and have pitched in together, era after era, to make the lab work and keep morale high. I have watched with pleasure as they pursue independent paths in science and in life. They are my extended family.

I list their names in Figure 11 in thanks and appreciation.

I also thank my Princeton faculty colleagues, and a wide circle of collaborators, for their friendship and support over the years.

Acknowledgements

Colin Weeks, Radhika Rajendran and Gurusamy Balakrishnan helped greatly in the preparation of this article. I am grateful to the NIH Institute of General Medical Sciences for steadfast support throughout my career.

References

1. Spiro TG, Hume DN. *J. Am. Chem. Soc* 1961;83:4305–4310.
2. Spiro TG, Hume DN. *Inorg. Chem* 1963;2:340–345.
3. Spiro TG, Ballhausen CJ. *Acta Chem. Scand* 1961;15:1707.
4. Biederman G, Spiro TG. *Chemica Scripta* 1971;1:155.
5. Spiro TG. *Inorg. Chem* 1965;4:731–738.
6. Maroni VA, Spiro TG. *J. Am. Chem. Soc* 1966;88:1410–1412. [PubMed: 5914522]
7. Maroni VA, Spiro TG. *J. Am. Chem. Soc* 1967;89:45–48.
8. Bulliner PA, Spiro TG. *Inorg. Chem* 1969;8:1023–1025.
9. Quicksall CO, Spiro TG. *Chem. Commun* 1967:839.
10. Bulliner PA, Quicksall Co, Spiro TG. *Inorg. Chem* 1971;10:13–18.
11. Farrell FJ, Spiro TG. *Inorg. Chem* 1971;10:1606–1610.
12. Fontal B, Spiro TG. *Inorg. Chem* 1971;10:9–13.
13. Terzis A, Strekas TC, Spiro TG. *Inorg. Chem* 1971;10:2617–2620.
14. Spiro TG. *Prog. Inorg. Chem* 1970;11:1.
15. Spiro TG, Allerton SE, Renner J, Terzis A, Bils R, Saltman P. *J. Am. Chem. Soc* 1966;88:2721–2726.
16. Spiro TG, Bates G, Saltman P. *J. Am. Chem. Soc* 1967;89:5559–5562.
17. Webb SM, Dick GJ, Bargar JR, Tebo BM. *Proc. Natl. Acad. Sci. U. S. A* 2005;102:5558–5563. [PubMed: 15800042]
18. Spiro, TG.; Stigliani, WM. *Chemistry of the environment*. 2nd ed.. Prentice Hall; 2001.
19. Nestor J, Spiro TG, Klauminzer G. *Proc. Natl. Acad. Sci. U. S. A* 1976;73:3329–3332. [PubMed: 185608]
20. Dutta PK, Nestor JR, Spiro TG. *Proc. Natl. Acad. Sci. U. S. A* 1977;74:4146–4149. [PubMed: 16592440]
21. Stein P, Miskowski V, Woodruff WH, Griffin JP, Werner KG, Gaber BP, Spiro TG. *J. Chem. Phys* 1976;64:2159–2167.
22. Stein P, Brown JM, Spiro TG. *Chem. Phys* 1977;25:237–244.
23. Dallinger RF, Stein P, Spiro TG. *J. Am. Chem. Soc* 1978;100:7865–7870.
24. Stein P, Jensen PW, Spiro TG. *Chem. Phys. Lett* 1981;80:451–454.
25. Miskowski V, Tang SPW, Spiro TG, Shapiro E, Moss TH. *Biochemistry* 1975;14:1244–1250. [PubMed: 804316]
26. Brown JM, Powers L, Kincaid B, Larrabee JA, Spiro TG. *J. Am. Chem. Soc* 1980;102:4210–4216.
27. Larrabee JA, Spiro TG. *J. Am. Chem. Soc* 1980;102:4217–4223.
28. Woolery GL, Powers L, Winkler M, Solomon EI, Spiro TG. *J. Am. Chem. Soc* 1984;106:86–92.
29. Nestor L, Larrabee JA, Woolery G, Reinhammar B, Spiro TG. *Biochemistry* 1984;23:1084–1093.
30. Kubas GT, Spiro TG, Terzis A. *J. Am. Chem. Soc* 1973;95:273–274. [PubMed: 4682896]
31. Johnson MK, Czernuszewicz RS, Spiro TG, Fee JA, Sweeney WV. *J. Am. Chem. Soc* 1983;105:6671–6678.
32. Yachandra VK, Hare J, Gewirth A, Czernuszewicz RS, Kimura T, Holm RH, Spiro TG. *J. Am. Chem. Soc* 1983;105:6462–6468.

33. Han S, Czernuszewicz RS, Spiro TG. *J. Am. Chem. Soc* 1989;111:3496–3504.
34. Han SW, Madden JF, Siegel LM, Spiro TG. *Biochemistry* 1989;28:5477–5485. [PubMed: 2673348]
35. Wozniak WT, Spiro TG. *J. Am. Chem. Soc* 1973;95:3402–3404. [PubMed: 4708830]
36. Salama S, Spiro TG. *J. Raman Spectrosc* 1977;6:57–60.
37. Dong SL, Padmakumar R, Banerjee R, Spiro TG. *J. Am. Chem. Soc* 1999;121:7063–7070.
38. Stich TA, Seravalli J, Venkatesh Rao S, Spiro TG, Ragsdale SW, Brunold TC. *J. Am. Chem. Soc* 2006;128:5010–5020. [PubMed: 16608335]
39. Kilpatrick L, Rajagopalan KV, Hilton J, Bastian NR, Stiefel EI, Pilato RS, Spiro TG. *Biochemistry* 1995;34:3032–3039. [PubMed: 7893715]
40. Pilato RS, Eriksen KA, Greaney MA, Stiefel EI, Goswami S, Kilpatrick L, Spiro TG, Taylor EC, Rheingold AL. *J. Am. Chem. Soc* 1991;113:9372–9374.
41. Dutta PK, Spiro TG. *Biochemistry* 1980;19:1590–1593. [PubMed: 7378367]
42. Copeland RA, Spiro TG. *J. Phys. Chem* 1986;90:6648–6654.
43. Bowman WD, Spiro TG. *J. Raman Spectrosc* 1980;9:369–371.
44. Benecky MJ, Copeland RA, Rava RP, Feldhaus R, Scott RD, Metzler CM, Metzler DE, Spiro TG. *J. Biol. Chem* 1985;260:1671–1678.
45. Woodruff WH, Spiro TG, Gilvarg C. *Biochem. Biophys. Res. Commun* 1974;58:197–203. [PubMed: 4208642]
46. Fodor SPA, Rava RP, Copeland RA, Spiro TG. *J. Raman Spectrosc* 1986;17:471–475.
47. Fodor SPA, Rava RP, Hays TR, Spiro TG. *J. Am. Chem. Soc* 1985;107:1520–1529.
48. Fodor SPA, Spiro TG. *J. Am. Chem. Soc* 1986;108:3198–3205.
49. Rava RP, Spiro TG. *Biochemistry* 1985;24:1861–1865. [PubMed: 3893540]
50. Fodor SPA, Copeland RA, Grygon CA, Spiro TG. *J. Am. Chem. Soc* 1989;111:5509–5518.
51. Grygon CA, Spiro TG. *Biopolymers* 1990;29:707–715.
52. Purrello R, Molina M, Wang Y, Smulevich G, Fossella J, Fresco JR, Spiro TG. *J. Am. Chem. Soc* 1993;115:760–767.
53. Mukerji I, Shiber MC, Fresco JR, Spiro TG. *Nucleic Acids Res* 1996;24:5013–5020. [PubMed: 9016674]
54. Hildebrandt PG, Copeland RA, Spiro TG, Otlewski J, Laskowski M, Prendergast FG. *Biochemistry* 1988;27:5426–5433. [PubMed: 3179264]
55. Caswell DS, Spiro TG. *J. Am. Chem. Soc* 1986;108:6470–6477.
56. Wu Q, Li FB, Wang WX, Hecht MH, Spiro TG. *J. Inorg. Biochem* 2002;88:381–387. [PubMed: 11897354]
57. Caswell DS, Spiro TG. *J. Am. Chem. Soc* 1987;109:2796–2800.
58. Copeland RA, Spiro TG. *Federation Proceedings* 1986;45:1869–1869.
59. Wang Y, Purrello R, Jordan T, Spiro TG. *J. Am. Chem. Soc* 1991;113:6359–6368.
60. Jordan T, Spiro TG. *J. Raman Spectrosc* 1995;26:867–876.
61. Austin JC, Zhao QJ, Jordan T, Talalay P, Mildvan AS, Spiro TG. *Biochemistry* 1995;34:4441–4447. [PubMed: 7703258]
62. Austin JC, Fitzhugh A, Villafranca JE, Spiro TG. *Biochemistry* 1995;34:7678–7685. [PubMed: 7779814]
63. Spiro TG, Strekas TC. *Proc. Natl. Acad. Sci. U. S. A* 1972;69:2622–2626. [PubMed: 4506783]
64. Spiro TG, Burke JM. *J. Am. Chem. Soc* 1976;98:5482–5489. [PubMed: 182734]
65. Woodruff WH, Spiro TG, Yonetani T. *Proc. Natl. Acad. Sci. U. S. A* 1974;71:1065–1069. [PubMed: 4524615]
66. Stein P, Burke JM, Spiro TG. *J. Am. Chem. Soc* 1975;97:2304–2305. [PubMed: 1133415]
67. Kincaid J, Stein P, Spiro TG. *Natl. Acad. Sci. U. S. A* 1979;76:549–552.
68. Choi S, Spiro TG, Langry KC, Smith KM, Budd DL, Lamar GN. *J. Am. Chem. Soc* 1982;104:4345–4351.
69. Parthasarathi N, Hansen C, Yamaguchi S, Spiro TG. *J. Am. Chem. Soc* 1987;109:3865–3871.
70. Choi S, Spiro TG, Langry KC, Smith KM. *J. Am. Chem. Soc* 1982;104:4337–4344.

71. Choi SH, Spiro TG. *J. Am. Chem. Soc* 1983;105:3683–3692.
72. Li XY, Czernuszewicz RS, Kincaid JR, Su YO, Spiro TG. *J. Phys. Chem* 1990;94:31–47.
73. Li XY, Czernuszewicz RS, Kincaid JR, Stein P, Spiro TG. *J. Phys. Chem* 1990;94:47–61.
74. Hu SZ, Mukherjee A, Piffat C, Mak RSW, Li XY, Spiro TG. *Biospectroscopy* 1995;1:395–412.
75. Prendergast K, Spiro TG. *J. Am. Chem. Soc* 1992;114:3793–3801.
76. Kozlowski PM, Rush TS, Jarzecki AA, Zgierski MZ, Chase B, Piffat C, Ye BH, Li XY, Pulay P, Spiro TG. *J. Phys. Chem. A* 1999;103:1357–1366.
77. Rush TS, Kozlowski PM, Piffat CA, Kumble R, Zgierski MZ, Spiro TG. *J. Phys. Chem. B* 2000;104:5020–5034.
78. Kumble R, Rush TS, Blackwood ME, Kozlowski PM, Spiro TG. *J. Phys. Chem. B* 1998;102:7280–7286.
79. Venkatesh Rao S, Yin J, Jarzecki AA, Schultz PG, Spiro TG. *J. Am. Chem. Soc* 2004;126:16361–16367. [PubMed: 15600337]
80. Blackwood ME, Rush TS, Romesberg F, Schultz PG, Spiro TG. *Biochemistry* 1998;37:779–782. [PubMed: 9457047]
81. Czernuszewicz RS, Macor KA, Li XY, Kincaid JR, Spiro TG. *Abstr. Pap. Am. Chem. Soc* 1988;196:200.INOR
82. Lin CY, Hu SZ, Rush T, Spiro TG. *J. Am. Chem. Soc* 1996;118:9452–9453.
83. Prendergast K, Spiro TG. *J. Phys. Chem* 1991;95:9728–9736.
84. Czernuszewicz RS, Su YO, Stern MK, Macor KA, Kim D, Groves JT, Spiro TG. *J. Am. Chem. Soc* 1988;110:4158–4165.
85. Blackwood ME, Kumble R, Spiro TG. *J. Phys. Chem* 1996;100:18037–18041.
86. Kumble R, Loppnow GR, Hu SZ, Mukherjee A, Thompson MA, Spiro TG. *J. Phys. Chem* 1995;99:5809–5816.
87. Vitols SE, Roman JS, Ryan DE, Blackwood ME, Spiro TG. *Inorg. Chem* 1997;36:764–769.
88. Eads DD, Moser C, Blackwood ME, Lin CY, Dutton L, Spiro TG. *Biopolymers* 2000;57:64–76. [PubMed: 10766957]
89. Ghosh PK, Spiro TG. *J. Electrochem. Soc* 1981;128:1281–1287.
90. Macor KA, Su YO, Miller LA, Spiro TG. *Inorg. Chem* 1987;26:2594–2598.
91. Loppnow GR, Melamed D, Hamilton AD, Spiro TG. *J. Phys. Chem* 1993;97:8957–8968.
92. Hu SZ, Morris IK, Singh JP, Smith KM, Spiro TG. *J. Am. Chem. Soc* 1993;115:12446–12458.
93. Hu SZ, Smith KM, Spiro TG. *J. Am. Chem. Soc* 1996;118:12638–12646.
94. Rakshit G, Spiro TG. *Biochemistry* 1974;13:5317–5322. [PubMed: 4433523]
95. Smulevich G, Evangelistakirkup R, English A, Spiro TG. *Biochemistry* 1986;25:4426–4430. [PubMed: 3019391]
96. Evangelistakirkup R, Smulevich G, Spiro TG. *Biochemistry* 1986;25:4420–4425. [PubMed: 3756147]
97. Li XY, Spiro TG. *J. Am. Chem. Soc* 1988;110:6024–6033.
98. Ray GB, Li XY, Ibers JA, Sessler JL, Spiro TG. *J. Am. Chem. Soc* 1994;116:162–176.
99. Spiro TG, Kozlowski PM. *J. Am. Chem. Soc* 1998;120:4524–4525.
100. Spiro TG, Kozlowski PM. *Accounts Chem. Res* 2001;34:137–144.
101. Dierks EA, Hu SZ, Vogel KM, Yu AE, Spiro TG, Burstyn JN. *J. Am. Chem. Soc* 1997;119:7316–7323.
102. Coyle CM, Puranik M, Youn H, Nielsen SB, Williams RD, Kerby RL, Roberts GP, Spiro TG. *J. Biol. Chem* 2003;278:35384–35393. [PubMed: 12796503]
103. Ibrahim M, Kerby RL, Puranik M, Wasbotten IH, Youn H, Roberts GP, Spiro TG. *J. Biol. Chem* 2006;281:29165–29173. [PubMed: 16873369]
104. Puranik M, Weeks CL, Lahaye D, Kabil Ö, Taoka S, Nielsen SB, Groves JT, Banerjee R, Spiro TG. *J. Biol. Chem* 2006;281:13433–13438. [PubMed: 16505479]
105. Gouterman, M. *Optical Spectra and Electronic Structure of Porphyrin and Related Rings*. In: Dolphin, D., editor. *The Porphyrins*. 3. Academic; New York: 1979. p. 1-156. Part A

106. Li XY, Czernuszewicz RS, Kincaid JR, Spiro TG. *J. Am. Chem. Soc* 1989;111:7012–7023.
107. Spiro TG, Wasbotten IH. *J. Inorg. Biochem* 2005;99:34–44. [PubMed: 15598489]
108. Vogel KM, Kozlowski PM, Zgierski MZ, Spiro TG. *J. Am. Chem. Soc* 1999;121:9915–9921.
109. Coyle CM, Vogel KM, Rush TS, Kozlowski PM, Williams R, Spiro TG, Dou Y, Ikeda-Saito M, Olson JS, Zgierski MZ. *Biochemistry* 2003;42:4896–4903. [PubMed: 12718530]
110. Ibrahim M, Xu C, Spiro TG. *J. Am. Chem. Soc* 2006;128:16834–16845. [PubMed: 17177434]
111. Xu C, Spiro TG. 2007submitted for publication
112. Woodruff WH, Spiro TG. *Appl. Spectrosc* 1974;28:576–578.
113. Dallinger RF, Nestor JR, Spiro TG. *J. Am. Chem. Soc* 1978;100:6251–6252.
114. Turner J, Spiro TG, Nagumo M, Nicol MF, Elsayed MA. *J. Am. Chem. Soc* 1980;102:3238–3239.
115. Dasgupta S, Spiro TG. *Biochemistry* 1986;25:5941–5948. [PubMed: 3790496]
116. Dasgupta S, Copeland RA, Spiro TG. *J. Biol. Chem* 1986;261:960–962.
117. Su C, Park YD, Liu GY, Spiro TG. *J. Am. Chem. Soc* 1989;111:3457–3459.
118. Rodgers KR, Su C, Subramaniam S, Spiro TG. *J. Am. Chem. Soc* 1992;114:3697–3709.
119. Jayaraman V, Rodgers KR, Mukerji I, Spiro TG. *Science* 1995;269:1843–1848. [PubMed: 7569921]
120. Hu XH, Spiro TG. *Biochemistry* 1997;36:15701–15712. [PubMed: 9398299]
121. Wang DJ, Spiro TG. *Biochemistry* 1998;37:9940–9951. [PubMed: 9665699]
122. Hu XH, Frei H, Spiro TG. *Biochemistry* 1996;35:13001–13005. [PubMed: 8855934]
123. Chen RP, Spiro TG. *J. Phys. Chem. A* 2002;106:3413–3419.
124. Balakrishnan G, Hu Y, Nielsen SB, Spiro TG. *Appl. Spectrosc* 2005;59:776–781. [PubMed: 16053544]
125. Balakrishnan G, Case MA, Pevsner A, Zhao XJ, Tengroth C, McLendon GL, Spiro TG. *J. Mol. Biol* 2004;340:843–856. [PubMed: 15223325]
126. Balakrishnan G, Tsai CH, Wu Q, Case MA, Pevsner A, McLendon GL, Ho C, Spiro TG. *J. Mol. Biol* 2004;340:857–868. [PubMed: 15223326]
127. Kneipp J, Balakrishnan G, Chen RP, Shen TJ, Sahu SC, Ho NT, Giovannelli JL, Simplaceanu V, Ho C, Spiro TG. *J. Mol. Biol* 2006;356:335–353. [PubMed: 16368110]
128. Huang CY, Balakrishnan G, Spiro TG. *J. Raman Spectrosc* 2006;37:277–282.
129. Balakrishnan G, Hu Y, Spiro TG. *Appl. Spectrosc* 2006;60:347–351. [PubMed: 16613628]
130. Huang CY, Balakrishnan G, Spiro TG. *Biochemistry* 2005;44:15734–15742. [PubMed: 16313176]
131. Ji Ji RD, Balakrishnan G, Hu Y, Spiro TG. *Biochemistry* 2006;45:34–41. [PubMed: 16388578]
132. Balakrishnan G, Hu Y, Case MA, Spiro TG. *J. Phys. Chem. B* 2006;110:19877–19883. [PubMed: 17020373]
133. Spiro, TG.; Czernuszewicz, RS. *Resonance Raman Spectroscopy*. In: Que, L., Jr., editor. *Physical Methods in Bioinorganic Chemistry*. University Science Books; Sausalito: 2000. p. 59-119.
134. Spiro, TG.; Li, Y-Y. *Resonance Raman Spectroscopy of Metalloporphyrins*. In: Spiro, TG., editor. *Biological Applications of Raman Spectroscopy*. 3. Wiley-Interscience; New York: 1988. p. 1-38.

Figure 1. Raman spectrum of 0.9 M TiCl_3 in 1 M HCl and the Carey 81 Raman spectrometer with its Hg arc light source.⁵

Figure 2. 70 Å Ferric hydroxide particles obtained by HCO_3^- addition to $\text{Fe}(\text{NO}_3)_3$ ¹⁵. (The line marks 1 μm).

Figure 3.

(Left) Flavin structure and absorption spectrum. (Right) CARS spectra, at two probe wavelengths, of the flavin co-factor in glucose oxidase, illustrating the complex CARS lineshapes.²⁰ ω_1 is the fixed laser wavelength (nm), ω_2 is the scanned laser wavelength (nm), $\Delta\omega = \omega_1 - \omega_2$ (cm^{-1}) and $\omega_{as} = 2\omega_1 - \omega_2$ is the wavelength of the coherent beam generated in the medium from ω_1 and ω_2 .

Figure 4. RR spectra of three 'blue' Cu proteins, and Mizkowski's model of the active site.¹⁸

Figure 5. Structure of an early FeS model³⁰ (top), and of a Holm-type model for [2Fe-2S] proteins³³ (bottom).

Figure 6. RR spectra of Fe(II) cytochrome c, with 514.5nm excitation, analyzed in parallel and perpendicular polarization.⁶³ Several bands show inverse polarization (ip) whereas the ν_1 band of the SO_4^{2-} internal standard is completely polarized.

Figure 7. DFT Computation of the heme-C $\leftrightarrow$ O transition dipole direction along a minimum energy distortion coordinate (bend + tilt), showing the IR-determined angle to be consistent with modest distortion.¹⁰¹

Figure 8.

Absorption spectrum of NiOEP (OEP=octaethylporphyrin) and its interpretation via Gouterman's four-orbital model.¹⁰⁵ The $e_g \rightarrow a_{2u}, a_{1u}$ orbital excitations (labeled 1 and 2), being of the same symmetry (E_u) and nearly degenerate, interact strongly, the transition dipoles adding up for the intense B transition at 392 nm, and nearly canceling for the weaker Q_0 transition at 552 nm. About 10% of the B band intensity is borrowed by the Q transition producing the Q_1 vibronic side band 516 nm, above Q_0 by $\approx 1300 \text{ cm}^{-1}$, the average frequency of the vibronically effective modes (figure from ref. ¹³³).

Figure 9.

The RR spectra in parallel (||) and perpendicular (⊥) scattering for NiOEP,⁷³ showing the selective enhancement of different modes with different excitation wavelengths. Resonance with the B absorption band (406.7 nm) enhances mainly polarized bands, arising from totally symmetric modes, A_{1g} , although appreciable enhancement is also seen for Jahn-Teller active B_{1g} modes (see ν_{10} and ν_{11}), which are depolarized. Resonance in the Q band region enhances depolarized (B_{1g} , B_{2g}) and anomalously polarized (A_{2g}) modes and are effective in vibronic mixing of the Q and B electronic transitions. Due to interference effects the A_{2g} modes are brought out most strongly via excitation between Q_0 and Q_1 (530.9 nm), while the B_{1g} and B_{2g} modes are brought out more strongly with excitation outside the Q bands (568.2 nm). The

mode labels ν_i refer to porphyrin skeletal modes;¹³⁴ modes assignable to the ethyl substituents are also seen, as indicated (figure from ref. 133).

Figure 10. UVRR spectra of Hb in (a) deoxy (T) and (b) CO (R) forms. The difference spectrum (c) shows tyrosine (Y) and tryptophan (W) signals, which are associated with H-bond changes between the T and R quaternary structures.¹¹⁷

Undergraduates

Adams, David, 1974	LePar, Felice, 1992
Berbit, Zachary, 2001	Li, James, 1974
Bigby, Michael, 1971	Lum Cheong, Ronnie, 1985
Boyd, Henry, 1986	Martin, Lauren, 2001
Carter, Edward, 1977	Mercier, Gustavio, 1982
Cleary, Susan, 1993	Meyne, Michael, 1973
Constantinescu, Andrei, 1987	Munger, John, 1978
Conway, Kelley, 1986	Murphy, Elizabeth, 1978
Daly, Peter, 1976	Myers, Jonathan, 1988
Danar, David, 1978	Narayan, Kartik, 1997
DiGabriele, Anna, 1975	Norwood, Daphne, 1985
Dixon, Frederick, 1971	Packer, Andrew, 1972
Elkus, Barbara, 1971	Peters, Steve, 1975
Friedman, Charles, 1967	Prieto, Roberto, 1989
Galvan, Noe, 1996	Raj, Vishwa, 1999
Gewirth, Andrew, 1981	Sanders, Winifred, 1971
Gilbert, Lewis, 1974	Schuessler-Firenza, Chris, 1996
Griffin, James, 1974	Shu, Aimee, 1996
Haber, Ann, 1978	Starr, Philip, 1983
Hansen, Christine, 1986	Subramanian, Prem, 1989
Ito, James, 1970	Talhah, Baktiar, 1996
Jacobs, Mary, 1974	Tuttle, Paul, 1976
Jack, John, 1965	Waters, Susan, 1985
Kalpaxis, James, 1969	Werner, Kirk, 1973
Kindle, Cecil, 1970	Wolfe, Patrick, 1975
Kish, Stephen, 1974	Worswick, Scott, 2002
Lee, Jane So Chun, 1981	Yager, Paul, 1982
Lee, Joe Yee, 1978	

Graduate Students

Benecky, Michael, 1983	Larrabee, James, 1979	Vitols, Sondra, 1995
Blackwood, Milton, 1997	Lee, Cheng-Cheng, 1975	Vogel, Kathleen, 1998
Bowman, David, 1980	Li, Xiao-Yuan, 1988	Walters, Marc, 1983
Brown, James, 1970	Lin, Ching-Yao, 1997	Wang, Daojing, 1999
Bulliner, Alan, 1970	Liu, Gang-Yu, 1988	Woolery, Geoffrey, 1983
Caswell, Debra, 1986	Macor, Kathleen, 1983	Xu, Changliang -
Chen, Roupian, 2000	Maroni, Victor, 1967	Yachandra, Vital, 1982
Choi-Larrabee, Sunhee, 1982	Melamad, Dan, 1992	Yang, Bing, 1995
Copeland, Robert, 1986	Miller, Lisa, 1986	
Dallinger, Richard, 1978	Mitchell, Melody, 1984	
Dasgupta, Siddharth, 1985	Mukherjee, Arka, 1995	
Dong, Shoulian, 1998	Oyerinde, Oyeyemi -	
Dutta, Prabir, 1978	Payne-Nestor, Lisa, 1984	
Fanelli, James, 1997	Parthasarathi, Niraja, 1987	
Farrell, Francis, 1969	Perno, Joseph, 1987	
Fodor, Stephen, 1980	Piffat, Christine, 1995	
Ghosh, Pushpito, 1980	Prendergast, Kristine, 1990	
Gruber, Suzanne, 1989	Qiu, Di, 1995	
Grygon, Christine, 1989	Quicksall, Carl, 1969	
Han, Sanghwa, 1988	Rakhit, Gopa, 1976	
Hu, Xuehua, 1998	Ray, Gigi, 1992	
Indritz, Doren, 1974	Remsen, Ed, 1980	
Jayaraman, Vasanthi, 1995	Salama, Simon, 1978	
Jordan, Trace, 1994	Sibilia, Sharon, 1997	
Kellett, Richard, 1985	Strekas, Tom, 1973	
Kilpatrick, LaTonya, 1992	Su, Chang, 1990	
Kumble, Ranjit, 1994	Terzis, Aris, 1970	

Postdoctoral Fellows

Austin-Eads, Janina, 1992	Hays, Thomas, 1984	Lobenstein, Eric, 1983	Smulevich, Giulietta, 1985
Balakrishnan, Gurusamy -	Heibel, George, 1996	Loppnow, Glen, 1993	Stein, Paul, 1979
Burke, Michael, 1977	Hildebrandt, Peter, 1986	Maslowsky, Ed, 1973	Stong, John, 1980
Cartling, Bo, 1984	Hu, Songzhou, 1996	McGlashen, Michael, 1994	Su, Oliver, 1986
Chan, Marie, 1979	Hu, Ying, 2004	Miskowski, Vincent, 1974	Sztainbuch, Isaac, 1993
Coppey, Maite, 1983	Huang, Cheng-Yen, 2005	Molina, Marina, 1990	Tengroth, Charbel, 2000
Coyle, Candace, 2000	Huang, Ta-Zan, 1974	Mukerji, Ishita, 1994	Turner, James, 1980
Coyne, Lelia, 1973	Huang, Yu-San, 2006	Nelson, Harold, 1981	Topp, William, 1974
Crisanti, Mark, 1983	Ibrahim, Mohammed -	Nestor, James, 1980	Tran, Dat, 2000
Czernuszewicz, Roman, 1989	Jarzecki, Andrzej, 2000	Nielsen, Steen, 2002	Valentine, Joan, 1972
Dick, Lisa, 1998	Jauffmann, Judith, 1985	Park, Young, 1988	Vargek Maria, 1998
Eads, Daniel, 1992	Jiji, Renee, 2004	Puranik, Mrinalini, 2004	Venkatesh Rao, Swarnalatha, 2004
Evangelista-Kirkup, Ruby, 1985	Johnson, Michael, 1984	Purello, Roberto, 1991	Wang, Yang, 1988
Fleissner, Gerhard, 1997	Kim, Dongho, 1985	Ramsden, Jeremy, 1986	Wasbotten, Ingar, 2004
Fontal, Bernardo, 1971	Kincaid, James, 1979	Rava, Richard, 1984	Weeks, Colin -
Francia-Zuckerman, Marie, 1972	Kneipp, Janina, 2004	Reed, Robert, 1990	Williams, Robert, 2000
Gay, Roger, 1973	Kozlowski, Pawel, 1999	Rodgers, Kenton, 1993	Woodruff, William, 1974
Georgiou, Savas, 1989	Kubas, Gregory, 1972	Rush, Thomas, 1998	Wozniak, Wayne, 1973
Gosztola, David, 1990	Lancaster, Kristie, 1985	Sanchez, Luis, 1984	Wright, Peter, 1978
Gregoriou, Vasilis, 1994	Leahy, Michael, 1981	Schultz, Richard, 1995	Wu, Qiang, 2000
Gutterman, Diane, 1970	Leheny, Rachel, 1991	Scovell, William, 1972	Yamaguchi, Satoshi, 1983
Hare, Jeffrey, 1979	Li, Fangbiao, 1999	Simpson, William, 1996	Zhao, Xiaojie, 2000

Visiting Scientists

Anderson, James, 1977
Andrews, Rodney, 1984
Anzenbacher, Pavel, 1984
Barrett, Terence, 1979
Cwikel, Dory, 2000
Feitelson, Yehuda, 1985
Gaber, Bruce, 1972
Jensen, Palle Waage, 1977
Lee, J.J., 1980
Mabrouk, Pamela, 1998
Nishimura, Yoshifumi, 1980
Okura, Ichiro, 1986
Ozaki, Yukihiro
Powers, Linda, 1982
Revesz, Agnes, 1966
Tang, Pamela, 1979
Tsuboi, Masimichi, 1988

Figure 11.
Colleagues of Thomas G. Spiro