Published in final edited form as:

J Phys Chem B. 2007 January 18; 111(2): 408–422. doi:10.1021/jp065403l.

Single-Ion Solvation Free Energies and the Normal Hydrogen Electrode Potential in Methanol, Acetonitrile, and Dimethyl Sulfoxide

Casey P. Kelly, Christopher J. Cramer^{*}, and Donald G. Truhlar^{*}

Department of Chemistry and Supercomputing Institute, 207 Pleasant Street SE, University of Minnesota, Minneapolis, Minnesota 55455-0431

Abstract

The division of thermodynamic solvation free energies of electrolytes into ionic constituents is conventionally accomplished by using the single-ion solvation free energy of one reference ion, conventionally the proton, to set the single-ion scales. Thus the determination of the free energy of solvation of the proton in various solvents is a fundamental issue of central importance in solution chemistry. In the present article, relative solvation free energies of ions and ion-solvent clusters in methanol, acetonitrile, and dimethyl sulfoxide (DMSO) have been determined using a combination of experimental and theoretical gas-phase free energies of formation, solution-phase reduction potentials and acid dissociation constants, and gas-phase clustering free energies. Applying the cluster pair approximation to differences between these relative solvation free energies leads to values of −263.5, −260.2, and −273.3 kcal/mol for the absolute solvation free energy of the proton in methanol, acetonitrile, and DMSO, respectively. The final absolute proton solvation free energies are used to assign absolute values for the normal hydrogen electrode potential and the solvation free energies of other single ions in the above solvents.

1. Introduction

For neutral species, experimental solvation free energies have been tabulated for a large number of solutes in both aqueous^{1–8} and nonaqueous^{3,9,10} solvents. Typically, these solvation free energies are determined directly from experimental partition coefficients,¹¹ and thus their uncertainty is relatively low (~0.2 kcal/mol).¹² Determining accurate values for the solvation free energies of ionic solutes is much less straightforward than for neutral solutes. It is generally agreed that the chemical potential of an ion has no “operational meaning”¹³ in classical thermodynamics because the difference in electric potential between two media cannot be measured and therefore has no physical significance.^{14–16} Nevertheless, single-ion solvation free energies are well defined in statistical mechanics, and determining their values is an important step in understanding the structure of solutions. The potential of a medium relative to vacuum is called the potential of the phase. The single-ion solvation free energy in the potential of the phase is called the absolute solvation free energy, and it differs from the real (unmeasurable) solvation free energy by the potential of the phase. The present article is concerned with absolute solvation free energies. The single-ion solvation free energy of the

^{*}cramer@chem.umn.edu, truhlar@umn.edu.

Supporting Information Available: A description of the standard state conventions used in this article, thermochemical conventions for the enthalpy and heat capacity of the electron, enthalpies, entropies, and free energies of formation of monatomic ions, conventional solvation free energies of unclustered and clustered ions, slopes and intercepts for the cluster pair fits, and B97–1/MG3S optimized geometries. This material is available free of charge via the Internet at <http://pubs.acs.org>.

proton is conventionally used to set the scale for all other single-ion solvation free energies. Understanding the partitioning of single ions between different liquid phases is also of great importance in many areas of biology. For example, the electrical signals sent by nerve cells are activated by changes in the resting cell potential that are caused by the movement of various ions across the neuronal membrane.¹⁷

Sums of solvation free energies of neutral pairs of cations and anions are well-defined and can be determined directly through the use of thermochemical cycles involving calorimetric or electrochemical measurements.^{7,18-22} A number of different extrathermodynamic approximations have been used^{16,23-30} to partition the sums of cation and anion solvation free energies into single-ion contributions. A popular method for doing this is based on the model of a structureless, spherical ion in a homogenous continuum dielectric, in which case the solvation free energy depends only on bulk electrostatic interactions with the solvent, which are independent of the sign of the ion's charge.^{31,32} Thus, one assumes that in a given solvent, the solvation free energy of a reference salt composed of a quasispherical cation and a quasispherical anion of about the same size can be divided equally. Once the absolute solvation free energies of these reference ions are known, the absolute solvation free energies of other single ions can then be determined using the appropriate thermochemical cycles. The justification for approximations like this is best if the ions are large so that their surface-to-volume ratio is minimized. Commonly, this approximation is invoked using tetraphenylarsonium tetraphenylborate ($\text{AsPh}_4\text{BPh}_4$, or TATB)³³ as the reference salt, although other reference salts have also been used³⁴⁻⁴⁰ for dividing free energy, enthalpy, and entropy. A number of experimental and theoretical studies have focused their attention on the spectroscopic and thermodynamic properties of the AsPh_4^+ and BPh_4^- ions in various solvents.^{28,41-52} The current general consensus seems to be that charge-specific interactions *do* occur between these two ions and neighboring solvent molecules (thus bringing into question the validity of the TATB approximation), although some experimental^{28,43} and theoretical^{45,52} work suggests just the opposite.

Another approximation that has been used to estimate the solvation free energies of single ions is the cluster pair approximation of Coe, Tuttle, and coworkers.^{30,53,54} In this approximation, the absolute solvation free energy of a reference ion (usually the proton) is determined through comparisons between the stepwise clustering free energies of cation-anion pairs containing different numbers of clustering solvent molecules and their bulk solvation free energies. This approximation has been used in several places to estimate single-ion solvation free energies in water.⁵³⁻⁵⁶ Recently, we applied the cluster pair approximation to a database containing relative solvation free energies of ions, which are well defined even in classical thermodynamics, and accurate gas-phase clustering free energies for ion-water clusters containing up to 6 clustering water molecules.⁵⁶ Using this database, we obtained a value of -266.1 kcal/mol for the absolute aqueous solvation free energy of the proton, which is in good agreement with the recommended⁵⁷ value of -265.9 kcal/mol.

Compared to water, estimates of the absolute solvation free energies of single ions in other solvents¹⁶ have been very limited. Kalidas et al. reviewed a large number of transfer free energies of single ions between water and 17 different nonaqueous solvents.⁵⁸ The majority of the transfer free energies listed in this compilation were determined using the TATB approximation; none of them were determined using the cluster pair approximation. Tuttle and coworkers have used the cluster pair approximation to determine the absolute solvation free energy of the proton in ammonia.⁵⁴ However, based on the limited amount of solution-phase and gas-phase data available to these workers, they could only make a tentative estimate. Besides various extrathermodynamic approximations, theoretical calculations have also been used to estimate the solvation free energies of single ions in the liquid phase.⁵⁹⁻¹¹² However,

like the majority of previous experimental work, most of these studies too have focused solely on water.

In this article, the cluster pair approximation will be used to determine the absolute solvation free energy of the proton in methanol, acetonitrile, and DMSO. For this purpose, relative solvation free energies obtained using experimental gas-phase free energies of formation and solution-phase reduction potentials and acid dissociation constants will be combined with experimental and calculated gas-phase clustering free energies.

In the first part of this article, the standard-state conventions used for gas-phase and solution-phase free energies will be outlined, followed by a description of the thermochemical cycles and experimental and theoretical data used to determine relative solvation free energies. Then, the cluster pair approximation will be briefly described, followed by the results obtained through the application of this approximation to the relative solvation free energies determined in the previous section. The accuracy of the absolute solvation free energies obtained here will then be discussed, followed by a comparison between these solvation free energies and those obtained using other extrathermodynamic assumptions and theoretical calculations.

2. Standard States

In this article, experimental and calculated gas-phase free energies use a standard-state gas-phase pressure of 1 atm. A superscript open circle denotes free energies that use this standard-state pressure. Experimental solution-phase free energies use a standard-state solution-phase concentration of 1 mol/L. A superscript asterisk denotes free energies that use this standard-state concentration. Absolute and conventional solvation free energies use for a standard state an ideal gas at a gas-phase concentration of 1 mol/L dissolving as an ideal solution at a liquid-phase concentration of 1 mol/L. A superscript asterisk denotes solvation free energies that use this standard-state concentration in the gas and solution phases. These standard state conventions, including procedures for converting various experimental quantities tabulated using other standard-state pressures and concentrations to the standard-state pressure and concentration used here, are described in full detail in the Supporting Information.

3. Relative and Conventional Solvation Free Energies of Unclustered Ions

Experimentally, only the sum of the solvation free energies of neutral combinations of cations and anions can be obtained. Consequently, solvation free energies of single ions are often tabulated as *relative* free energies by arbitrarily setting the free energy of solvation of some reference ion equal to zero.¹¹³ The choice of reference ion in a particular solvent is arbitrary, although by convention the proton is usually chosen. For monovalent ions, this results in a set of *conventional* free energies of solvation that for cations are shifted from their absolute values by the unknown value for the absolute solvation free energy of the proton

$$\Delta G_s^{*,\text{con}}(\text{M}^+) = \Delta G_s^*(\text{M}^+) - \Delta G_s^*(\text{H}^+) \quad (1)$$

and conventional free energies of solvation for anions that are shifted by an equal amount in the opposite direction

$$\Delta G_s^{*,\text{con}}(\text{X}^-) = \Delta G_s^*(\text{X}^-) - \Delta G_s^*(\text{H}^+) \quad (2)$$

(Above, M^+ refers generically to *any* monovalent cation, and X^- to *any* monovalent anion. Later in this article, BH^+ will be used to refer to a monovalent cation that behaves as a Brønsted–Lowry acid, and A^- will be used to refer to a monovalent anion that behaves as a Brønsted–Lowry base.) Following the above convention results in the following relationship

$$\Delta G_s^{*,\text{con}}(\text{M}^+) + \Delta G_s^{*,\text{con}}(\text{X}^-) = \Delta G_s^*(\text{M}^+) + \Delta G_s^*(\text{X}^-) \quad (3)$$

Thus, sums of conventional solvation free energies for neutral cation-anion pairs, which can be determined directly through the use of various thermodynamic cycles, are well- defined. The thermochemical cycles that are used in this article to determine conventional solvation free energies are described below.

3.1. Conventional Solvation Free Energies from Reduction Potentials

When the convention described above for the absolute solvation free energy of the proton is followed, the solution-phase free energy change associated with the following half-cell reaction

is equal to zero. (In the above reaction, “g” denotes the gas phase and “S” denotes the liquid solution phase.) Reduction potentials that are tabulated following this convention for the hydrogen electrode are referred to as standard reduction potentials. The half-cell reaction for the reduction of a monovalent metal cation is

where the symbol “cr” denotes the crystalline phase. Combining the half-cell reactions shown in eqs 4 and 5 leads to the following redox reaction

Through the use of thermochemical cycle 1 (illustrated in Scheme 1) and eq 1, the ! conventional solvation free energy of M^+ can be written as

$$\Delta G_{\text{s}}^{*,\text{con}}(\text{M}^+) = \Delta_{\text{f}}G_{298}^0[\text{H}^+(\text{g})] - \Delta_{\text{f}}G_{298}^0[\text{M}^+(\text{g})] + FE_{\text{c}}^* \quad (7)$$

where $\Delta_{\text{f}}G_{298}^0[\text{H}^+(\text{g})]$ and $\Delta_{\text{f}}G_{298}^0[\text{M}^+(\text{g})]$ are the gas-phase free energies of formation of H^+ and M^+ , respectively, at 298 K, F is Faraday's constant, which is equal to 23.061 kcal mol⁻¹ V⁻¹, and E_{c}^* is the standard reduction potential (in volts) of M^+ in the solution phase. (The free energies of formation of H_2 in the gas phase and of the neutral metal atom in the crystalline phase do not appear in eq 7 because these species correspond to the standard states for the elements from which they are composed. Consequently, their gas-phase free energies of formation are defined to equal zero at all gas-phase pressures.) The superscript open circles in $\Delta_{\text{f}}G_{298}^0[\text{H}^+(\text{g})]$ and $\Delta_{\text{f}}G_{298}^0[\text{M}^+(\text{g})]$ denote the use of a standard-state gas-phase pressure of 1 atm. The superscript asterisk and the subscript c in E_{c}^* denote the use of a standard-state solution-phase concentration of 1 mol/L (molarity scale). Often, standard reduction potentials taken from the literature use a standard-state solution-phase concentration of 1 mol solute per kg solvent (molality scale). The procedure that was used to convert standard reduction potentials tabulated on the molality scale to the molarity scale is described in the Supporting Information.

Gas-phase free energies of formation of monatomic ions can be calculated using tabulated values for the enthalpy of formation and entropies computed using the Sackur-Tetrode equation.¹¹⁴ In this work, enthalpies of formation of ions are tabulated following the electron convention¹¹⁵ for the integrated heat capacity and entropy of the electron. When this convention is followed, the electron is treated as a standard chemical element, whereby its integrated heat capacity and entropy can be computed using standard formulae from statistical mechanics. Often, Boltzmann statistics are used to compute the integrated heat capacity and entropy of the electron, although Bartmess¹¹⁶ and several earlier workers¹¹⁷⁻¹²⁰ have pointed out that a more correct treatment of the electron uses Fermi-Dirac, not Boltzmann statistics. (Like the electron, the proton is a fermion, although for 298 K, its integrated heat capacity and entropy obtained using Boltzmann statistics differs very little from the values

obtained using Fermi-Dirac statistics; see Table 3 of reference ¹¹⁶.) It is important to point out that conventional solvation free energies are independent of the values used for the integrated heat capacity and entropy of the electron, as long as the same values are used *consistently*. However, for reactions where the electron appears as a reactant or product (*e.g.*, chemical reactions for the free energies of formation of ionic species) the gas-phase free energy of reaction *does* depend on the values used for the electron integrated heat capacity and entropy. For example, using the integrated electron heat capacity and entropy obtained from Fermi-Dirac statistics ¹¹⁶ leads to a value of 361.7 kcal/mol for the free energy of formation of the gas-phase proton at 298 K and 1 atm. If instead the electron integrated heat capacity and entropy obtained from Boltzmann statistics are used, the free energy of formation of the gas-phase proton is equal to 362.6 kcal/mol at 298 K and 1 atm. In this work, all free energies of formation of gas-phase ions are tabulated using the electron integrated heat capacity and entropy obtained using Fermi-Dirac statistics. Further details regarding the different conventions that are commonly used for treating the thermochemistry of the electron, the equations and literature references that were used to determine the enthalpies, entropies, and free energies of formation of ions used in this article, and the procedure used to convert entropies computed using a standard-state gas-phase pressure of 1 bar to a standard-state gas-phase pressure of 1 atm, are given in the Supporting Information.

Using the gas-phase free energies of formation of ions tabulated as part of this work (see Supporting Information) and experimental ^{121,122} values for the standard reduction potentials of different metal cations in methanol, acetonitrile, and DMSO, conventional solvation free energies have been determined in these solvents by eq 7. These conventional solvation free energies and the auxiliary data used in eq 7 are listed in Table 1. To verify the accuracy of eq 7, we also determined the aqueous conventional solvation free energy of Ag⁺ using an experimental value ¹²³ of 0.7996 V for the standard reduction potential of Ag⁺ in water and the tabulated values for the gas-phase free energy of formation of H⁺ and Ag⁺. Substituting these values into eq 7 gives a value of 147.2 kcal/mol for the aqueous conventional solvation free energy of Ag⁺, which is in perfect agreement with the aqueous conventional solvation free energy listed in the recent compilation by Fawcett. ¹²⁴

Conventional solvation free energies in methanol were also determined using experimental values for the reduction potentials at various silver halide electrodes. The half-cell reaction for the silver halide electrode is

Combining the half-cell reactions shown in eqs 4 and 8 gives the following redox reaction

Through the use of thermochemical cycle 2 (illustrated in Scheme 2) and eq 2, the conventional solvation free energy of the halide anion X⁻ can be written as

$$\Delta G_{\text{s}}^{*,\text{con}} (\text{X}^-) = -\Delta G_{\text{g}}^0 - FE_{\text{c}}^* - 2\Delta G^{0 \rightarrow *} \quad (10)$$

where

$$\Delta G_{\text{g}}^0 = \Delta_{\text{f}} G_{298}^0 [\text{X}^- (\text{g})] + \Delta_{\text{f}} G_{298}^0 [\text{H}^+ (\text{g})] - \Delta_{\text{f}} G_{298}^0 [\text{AgX (cr)}] \quad (11)$$

The term $\Delta G^{0 \rightarrow *}$ appearing in eq 10 is the free energy change associated with moving a ! solute from a standard-state gas-phase concentration of 1 atm to a standard-state solution-!phase concentration of 1 mol/L. (This free energy change has also been referred to as the “compression” work of the gas ¹¹⁰ or the liberation free energy ¹²⁵) At 298 K, $\Delta G^{0 \rightarrow *}$ is equal to 1.9 kcal/mol. Full details of this free energy change are given in the Supporting Information.

(Note that the term $\Delta G^{0 \rightarrow *}$ does not appear in eq 7 because in this equation the number of moles of reactants that move from the gas phase to solution cancels out the number of moles of products that move from the gas phase to solution.)

Using tabulated values for the gas-phase free energies of formation of AgCl,¹²⁶ AgBr,¹²⁶ and AgI¹²⁶ (adjusted to a standard-state gas-phase pressure of 1 atm), as well as the gas-phase free energies of formation of H⁺, Cl⁻, Br⁻, and I⁻ tabulated in this work and experimental values¹²⁷ for the standard reduction potentials at the AgCl, AgBr, and AgI electrodes, conventional solvation free energies of Cl⁻, Br⁻, and I⁻ in methanol were determined using eqs 10 and 11. These conventional solvation free energies and the auxiliary data used in eqs 10 and 11 are listed in Table 2. To verify the accuracy of eqs 10 and 11, we determined the conventional aqueous solvation free energy of Cl⁻ using tabulated values for the gas-phase free energy of formation of AgCl, H⁺, and Cl⁻, and an experimental value¹²³ of 0.22233 V for the standard reduction potential at the AgCl electrode in water. Substituting these values into eqs 10 and 11 gives a value of -340.1 kcal/mol for the conventional aqueous solvation free energy of Cl⁻. Adjusting the value for the conventional solvation free energy of Cl⁻ listed in the compilation by Fawcett,¹²⁴ which uses a standard-state gas-phase concentration of 1 bar, to a standard-state gas-phase concentration of 1 mol/L (see the Supporting Information for details regarding this standard-state conversion) gives a value of -340.4 kcal/mol, which is in good agreement with the value obtained in this work.

3.2. Conventional Solvation Free Energies from Solubility Products

Experimental values for the equilibrium constant associated with the following chemical reaction

have been measured for a number of salts in both water and in nonaqueous solvents (this equilibrium constant is usually referred to as the solubility product, or K_{sp}). Through the use of thermochemical cycle 3 (illustrated in Scheme 3), the sum of the solvation free energies for the neutral cation–anion pair M^+X^- can be written as

$$\Delta G_{\text{S}}^* (\text{M}^+) + \Delta G_{\text{S}}^* (\text{X}^-) = 2.303RT \text{p}K_{\text{sp}} + \Delta G_{\text{lat}}^0 [\text{MX (cr)}] - 2\Delta G^{0 \rightarrow *} \quad (13)$$

where $\text{p}K_{\text{sp}}$ and $\Delta G_{\text{lat}}^0 [\text{MX (cr)}]$ are the negative common logarithm of the solubility product and the lattice free energy of the crystalline salt, respectively. The lattice free energy is given by

$$\Delta G_{\text{lat}}^0 [\text{MX (cr)}] = \Delta_{\text{f}}G_{298}^0 [\text{MX (cr)}] - \Delta_{\text{f}}G_{298}^0 [\text{M}^+ (\text{g})] - \Delta_{\text{f}}G_{298}^0 [\text{X}^- (\text{g})] \quad (14)$$

Using tabulated values for the free energies of formation of various salts¹²⁶ (adjusted to a standard-state gas-phase pressure of 1 atm), the gas-phase free energies of formation of ions tabulated in this work, and experimental^{122,128} values for the solubility products of different salts, sums of solvation free energies for different cation–anion pairs in methanol, acetonitrile, and DMSO were determined using eqs 13 and 14. For the majority of the salts considered in this work, experimental values for their solubility products were taken from ref¹²², in which the authors have compiled experimental values taken from their own work and from a number of other experimental and review papers. In cases where more than a single experimental value for a given salt in a given solvent was listed by these authors, we took the average of all the listed values, except in cases where an experimental value was designated “uncertain” by these authors, in which case it was not used to calculate the average. More recently, Labban and Marcus¹²⁹ measured the solubilities of KCl and KBr in several nonaqueous solvents, including acetonitrile and DMSO. The solubility products of KCl and KBr computed using Labban and Marcus' data are both within 0.1 $\text{p}K_{\text{sp}}$ units of the average $\text{p}K_{\text{sp}}$ values computed using the data from ref¹²². (Reference 122 does not contain data for KCl or KBr in DMSO.) For KBr in

DMSO, the pK_{sp} value used in this work (0.6) is the average of three different experimental values: 0.5,¹³⁰ 0.5,¹³¹ and 0.7.¹²⁹ (The pK_{sp} value taken from ref¹³¹ was extrapolated to 298 K using pK_{sp} values measured at five other temperatures by these workers.) The sums of solvation free energies obtained using eqs 13 and 14 and the auxiliary data used in these equations are listed in Table 3.

To verify the accuracy of eqs 13 and 14, we determined the aqueous solvation free energy of the Ag^+Cl^- cation–anion pair using tabulated values for the free energy of formation of $AgCl$, Ag^+ , and Cl^- , and an experimental value¹²³ of 9.75 for the pK_{sp} of $AgCl$ in water. Substituting these values into eqs 13 and 14 gives a value of -193.4 kcal/mol for the aqueous conventional solvation free energy of the Ag^+Cl^- cation–anion pair. Taking the sum of the conventional solvation free energies of Ag^+ and Cl^- in water from the compilation by Fawcett¹²⁴ gives a value of -193.2 kcal/mol, which is in good agreement with the value obtained in this work.

Using the sums of solvation free energies in Table 3, conventional solvation free energies of Cl^- , Br^- , and I^- in methanol, acetonitrile, and DMSO were determined by subtracting from the solvation free energies of different cation–anion pairs the conventional solvation free energies of the corresponding cations from Table 1. (For example, the conventional solvation free energy of I^- in acetonitrile was determined by subtracting from the solvation free energy of the Ag^+I^- cation–anion pair the conventional solvation free energy of Ag^+ from Table 1.) Thus, all of the conventional solvation free energies of anions that are listed in Table 3 are anchored to the data in Table 1. In DMSO, conventional solvation free energies for Na^+ and K^+ are not listed in Table 1. Instead, these conventional solvation free energies were determined using the data in Tables 1 and 3, as described below. The conventional solvation free energy of Na^+ in DMSO was determined by subtracting from the solvation free energy of the Na^+Cl^- cation–anion pair the average value of the conventional solvation free energy for Cl^- obtained using the Ag^+Cl^- cation–anion pair and the Tl^+Cl^- cation–anion pair (these two conventional solvation free energies are listed in the final column of Table 3 and are within 0.4 kcal/mol of one another). Two values for the conventional solvation free energy of K^+ in DMSO were determined in a similar fashion using the K^+Cl^- and K^+Br^- cation–anion pairs and the conventional solvation free energies of Cl^- and Br^- . Thus, the conventional solvation free energy of Na^+ in DMSO is anchored to the data in Table 3 as well as to the conventional solvation free energies of Ag^+ and Tl^+ from Table 1. Similarly, the conventional solvation free energy of K^+ in DMSO is anchored to the Table 3 as well as to the conventional solvation free energy of Ag^+ from Table 1.

3.3. Conventional Solvation Free Energies from Acid Dissociation Constants

Shown in thermochemical cycle 4 (illustrated in Scheme 4)¹³² are the chemical reactions for the dissociation of a cationic Brønsted–Lowry acid (BH^+) in the gas phase and in the solution phase. Through the use of thermochemical cycle 4 and eq 1, the conventional solvation free energy of BH^+ can be written as

$$\Delta G_s^{*,con}(BH^+) = \Delta G_g^0(BH^+) + \Delta G_s^*(B) - 2.303RTpK_a(BH^+) + \Delta G^{0 \rightarrow *}$$
(15)

where $\Delta G_g^0(BH^+)$ is the gas-phase acidity of BH^+ , which is equal to $G^0(B) + G^0(H^+) - G^0(BH^+)$, $\Delta G_s^*(B)$ is the solvation free energy of the neutral species B , and $pK_a(BH^+)$ is the negative common logarithm of the solution-phase acid dissociation constant of BH^+ . Shown in thermochemical cycle 5 (illustrated in Scheme 5)¹³² are the chemical reactions for the dissociation of a neutral Brønsted–Lowry acid (AH) in the gas phase and in the solution phase. Through the use of thermochemical cycle 5 and eq 2, the conventional solvation free energy of A^- can be written as

$$\Delta G_s^{*,con}(A^-) = -\Delta G_g^0(AH) + \Delta G_s^*(AH) + 2.303RTpK_a(AH) - \Delta G^{0 \rightarrow *}$$
(16)

where $\Delta G_g^0(\text{AH})$ is the gas-phase acidity of AH, which is equal to $G^0(\text{A}^- + \text{H}^+) - G^0(\text{AH})$, $\Delta G_s^*(\text{AH})$ is the solvation free energy of the neutral species AH, and $\text{p}K_a(\text{AH})$ is the negative common logarithm of the solution-phase acid dissociation constant of AH.

Listed in Table 4 are experimental $\text{p}K_a$ values of neutral and cationic acids in methanol, acetonitrile, and DMSO.¹³³⁻¹⁵⁶ An experimental value for the solvation free energy is available¹⁰ for ethanol in DMSO; for all remaining neutral species, the solvation free energies in methanol, acetonitrile, and DMSO were calculated at the SM5.43R^{12,157}/*mPW1PW*^{158/6} $-31+G(\text{d,p})$ ¹⁵⁹ level of electronic structure theory, using gas-phase geometries optimized at the *mPW1PW/MIDI*¹⁶⁰⁻¹⁶² level of electronic structure theory (note that the *mPW1PW* functional is also called *mPW0*, *mPW1PW91*, and *MPW25*). All SM5.43R calculations were performed using a locally modified version¹⁶³ of the *Gaussian 03*¹⁶⁴ electronic structure package. Using the experimental $\text{p}K_a$ data in Table 4, experimental values for the gas-phase acidity,¹⁶⁵ and experimental and calculated values for the solvation free energies of neutral species, conventional solvation free energies were determined using eqs 15 and 16 for all of the acids in Table 4. These conventional solvation free energies and the auxiliary data used in eqs 15 and 16 are listed in Table 5.

3.4. Comparison Between Conventional Solvation Free Energies Obtained Using Different Methods

With the exception of eqs 15 and 16, which have appeared in several of our previous papers,^{7,56,111} the accuracy of all of the equations that were used in this article to determine conventional solvation free energies in this article were verified by comparing conventional aqueous solvation free energies determined using these equations to those listed in the compilation by Fawcett.¹²⁴ In all cases, the conventional aqueous solvation free energies determined here are in good agreement with those reported by Fawcett. This is encouraging, because it demonstrates not only the correctness of the equations used here but also, more importantly, the accuracy of the experimental gas-phase and aqueous-phase data that were used to test the equations. For all of the polyatomic ions considered in this article, the conventional solvation free energies were determined using only one of the equations described above (either eq 15 or 16), so that a comparison between these conventional solvation free energies, which might provide some insight into the relative accuracy of the experimental $\text{p}K_a$ data that were used in this article, is not possible. Equations 10 and 13 have both been used to determine the conventional solvation free energies of Cl^- and Br^- , and I^- in at least one of the nonaqueous solvents considered in this work. Thus, the conventional solvation free energies of these anions can be used to examine the relative accuracy of the remaining solution-phase experimental data used in this article. An additional check on the relative accuracy of the data in Table 3 (and thus Table 1) can be made because in all three solvents, two or more values for the conventional solvation free energies of Cl^- and Br^- have been determined in each solvent using experimental solubility products of different salts. For example, for Br^- in methanol, eq 13 has been used to make four independent determinations of its conventional solvation free energy based on experimental values for the solubility products of KBr, RbBr, AgBr, and TlBr and the conventional solvation free energies of corresponding cations. The largest difference between any of the four values listed in Table 3 for the conventional solvation free energy of Br^- in methanol is 0.7 kcal/mol. The largest discrepancy between any of the conventional solvation free energies listed in Table 3 for a given anion in a given solvent is 4.4 kcal/mol, which is the difference between the conventional solvation free energy of Br^- in acetonitrile determined using the solubility product of AgBr and the conventional solvation free energy of Ag^+ , and the value obtained using the solubility product of CsBr and the conventional solvation free energy of Cs^+ . Besides this difference, all of the remaining conventional solvation free energies listed in Table 3 for a given anion in a given solvent are within 2.3 kcal/mol of one another.

Next, we can compare the relative accuracy of the solution-phase data used in eq 10 to that used in eq 13. Using eq 10 leads to values of -334.8 , -329.0 , and -321.6 kcal/mol for the conventional solvation free energies of Cl^- , Br^- , and I^- , respectively, in methanol. From Table 3, the average values for the conventional solvation free energies of Cl^- , Br^- , and I^- are -335.7 , -329.4 , and -321.7 kcal/mol, respectively. The differences between these two sets of conventional solvation free energies are 0.9, 0.4, and 0.1 kcal/mol for Cl^- , Br^- , and I^- , respectively.

4. Conventional Solvation Free Energies of Clustered Ions

Conventional solvation free energies of clustered ions can be defined in the same way as unclustered ions, that is

$$\Delta G_s^{*,\text{con}}[(\text{S})_n\text{M}^+] = \Delta G_s^*[(\text{S})_n\text{M}^+] - \Delta G_s^*(\text{H}^+) \quad (17)$$

$$\Delta G_s^{*,\text{con}}[(\text{S})_n\text{X}^-] = \Delta G_s^*[(\text{S})_n\text{X}^-] + \Delta G_s^*(\text{H}^+) \quad (18)$$

where n is the number of clustering solvent (S) molecules. Through the use of thermochemical cycle 6 (illustrated in Scheme 6)¹³² the conventional solvation free energy of clustered cations and anions can be written in terms of the conventional solvation free energy of their analogous unclustered ions according to

$$\Delta G_s^{*,\text{con}}[(\text{S})_n\text{M}^\pm] = \Delta G_s^{*,\text{con}}(\text{M}^\pm) - \Delta G_{[n]}^0(\text{M}^\pm) + n\Delta G_{\text{self}}^*(\text{S}) + n\Delta G^{0 \rightarrow *} \quad (19)$$

where $\text{M}^\pm$ is the same as M^+ , X^- , BH^+ and A^- , $\Delta G_{\text{self}}^*(\text{S})$ is the free energy associated with moving the solute from the gas phase into its own pure liquid phase (often, we refer to this free energy as a “self solvation free energy”¹⁶⁶) and $\Delta G_{[n]}^0(\text{M}^\pm)$ is the sum of the stepwise clustering free energies of $\text{M}^\pm$ with n solvent (S) molecules in the gas-phase

$$\Delta G_{[n]}^0(\text{M}^\pm) = \sum_{i=1}^n \Delta G_{i-1,i}^0(\text{M}^\pm) \quad (20)$$

where

$$\Delta G_{i-1,i}^0 = G^0[(\text{S})_i\text{M}^\pm] - G^0[(\text{S})_{i-1}\text{M}^\pm] - G^0(\text{S}) \quad (21)$$

Note that the concentration of the pure liquid does not appear in eq 19 because we use as a standard state an ideal dilute solution, for which the activity of the pure liquid is very nearly equal to unity.¹⁶⁷ In this standard-state, the free energy associated with the following reaction

which we have also used in previous work^{7,56,111} and is shown in the bottom leg of thermochemical cycle 6, is equal to zero for any value of n .¹⁶⁸

Free energies of self solvation can be determined using the following equation¹⁶⁶

$$\Delta G_{\text{self}}^* = -2.303RT \log \left(\frac{M_{\text{liq}}/M^0}{P_{\text{vapor}}/P^0} \right) \quad (23)$$

where M^0 is equal to 1 mol/L, P^0 is the pressure (24.45 atm) of an ideal gas at 1 molar concentration and 298 K, and M_{liq} is the molarity of the solute in its pure liquid form, which can be calculated from the density of the solute in its pure liquid state and its molecular weight

$$M_{\text{liq}} = \rho_{\text{liq}} \text{MW}_{\text{liq}} \quad (24)$$

Using eqs 23 and 24, we determined the free energies of self solvation of methanol, acetonitrile, and DMSO. These solvation free energies and the auxiliary data used in eqs 23 and 24 are listed in Table 6.

Experimental values are available for the gas-phase free energies associated with attaching one or more methanol, acetonitrile, or DMSO molecules to many of the monatomic ions in Tables 1-3, as well as to some of the polyatomic ions listed in Table 5.¹⁶⁹ These experimental gas-phase free energies are listed in Table 7, for up to $n = 3$ in some cases. For some of the polyatomic ions in Table 5, we used eq 21 to calculate the gas-phase free energy associated with attaching a single solvent molecule to the ion. These calculated gas-phase clustering free energies, which are also listed in Table 7, are described below.

Theoretical gas-phase clustering free energies were calculated at the B97-1¹⁷⁰/MG3S¹⁷¹ level of theory using *Gaussian 03*.¹⁶⁴ (Because the MG3S basis set is not available in the distributed version of *Gaussian 03*, the “gen” keyword, which allows *Gaussian 03* to read an external, user-defined basis set, was used for all B97-1/MG3S calculations. The MG3S basis set can be downloaded via the internet at <http://comp.chem.umn.edu/basissets/>.) All gas-phase clustering free energies were calculated using a single gas-phase geometry for each solvent molecule, each unclustered ion, and each clustered ion (full geometry optimizations were carried out for all of the clustered ions considered in this work); that is, we did not average over multiple conformations. Gas-phase free energies were calculated following the harmonic-oscillator rigid-rotor approximation.¹⁷² When the rigid-rotor approximation is followed for computing the entropy, a contribution equal to $RT \ln \sigma$, where σ is the symmetry number of the molecule, is added to the calculated gas-phase free energy. (For example, the symmetry number for the ammonium cation, which belongs to the T_d point group, is equal to 12, so 1.47 kcal/mol is added to the gas-phase free energy; when symmetry is enforced in the *Gaussian 03* program, this is done automatically.) Because for many of the molecules considered in this article the above contribution to the gas-phase free energy is not negligible, we were careful to ensure that symmetry was enforced during all gas-phase calculations (when doing so led to a lowest-free-energy structure). The optimized gas-phase geometries for methanol and DMSO belong to the C_s point group; the optimized gas-phase geometry for acetonitrile belongs to the C_{3v} point group. For the majority of the clustered ions, conformational analyses were required in order to identify the global minimum. For this, we used the same level of theory as above. With the exception of the global minimum for the ammonium-acetonitrile cluster, which belongs to the C_{3v} point group, the geometries corresponding to the global minima for the remaining clustered ions belong to the C_1 or C_s point groups. The symmetry point groups and Cartesian coordinates for all of the clustered ions for which B97-1/MG3S geometries were calculated are listed in the Supporting Information.

Using eqs 19 and 20, the free energies of self solvation listed in Table 6, the experimental and calculated gas-phase clustering free energies in Table 7, and the conventional solvation free energies of unclustered ions from Tables 1, 2, 3, and 5, we determined conventional solvation free energies for clustered ions containing up to, in some cases, three methanol, acetonitrile, or DMSO molecules. For the halide anions, in cases where more than one value for the conventional solvation free energy of the unclustered ion is available, we used the average value from Tables 2 and 3. Tables containing all of the conventional solvation free energies of unclustered ions that were used in eq 19, as well as the resulting conventional solvation free energies for all of the clustered ions, are included in the Supporting Information.

5. Cluster Pair Approximation for Determining Absolute Solvation Free Energies of Single Ions

The cluster pair approximation described by Coe, Tuttle, and coworkers^{30,53,54} is based on the approximation that the difference between the absolute solvation free energy of a positive and negative cluster ion goes to zero as the cluster size becomes infinite, that is¹⁷³

$$\lim_{n \rightarrow \infty} \left\{ \Delta G_s^* [(S)_n X^-] - \Delta G_s^* [(S)_n M^+] \right\} = 0 \quad (24)$$

Using eqs 17-19, the above equation can be written as¹⁷⁴

$$\frac{1}{2} \left[\Delta G_s^{*,\text{con}} (X^-) - \Delta G_s^{*,\text{con}} (M^+) + \Delta G_{[n]}^0 (M^+) - \Delta G_{[n]}^0 (X^-) \right] = \frac{1}{2} \left[\Delta G_{[n+1,\infty]}^0 (M^+) - \Delta G_{[n+1,\infty]}^0 (X^-) \right] + \Delta G_s^* (H^+) \quad (25)$$

The left-hand-side (lhs) of the above equation is an approximation to the proton's solvation free energy, based on conventional solvation free energies of unclustered ions and the gas-phase free energy associated with attaching n solvent molecules to the cation and to the anion. On the right-hand-side (rhs) of the above equation are the absolute solvation free energy of the proton and the gas-phase free energy associated with attaching an infinite number of solvent molecules to an anion and to a cation already containing n solvent molecules. All three of the quantities on the rhs of eq 25 are unknown. Thus, the goal of eq 25, which is also referred to as the cluster pair approximation, is to identify a cation-anion pair for which the term in brackets on the rhs of eq 25 equals zero, in which case the lhs of eq 25 is equal to the proton's true aqueous solvation free energy (note that this ideal cation-anion pair need not actually exist). It has been shown^{30,53,54} that an effective way to do this is by plotting the lhs of eq 25 against half the difference between the conventional solvation free energy of X^- and M^+ for different values of n , giving n straight lines that share a common intersection point at the true value for the solvation free energy of the proton. This is the approach that will be used in this article.

6. Absolute Solvation Free Energy of the Proton and Normal Hydrogen Electrode Potential in Different Solvents

6.1. Methanol

Shown in Figure 1 is a plot of half the difference between conventional solvation free energies of anions and of cations for cluster ions containing up to three methanol molecules against half the difference between conventional solvation free energies of anions and of cations containing no methanol molecules, for different numbers of clustering methanol molecules. The differential conventional solvation free energies in this plot were computed using the conventional solvation free energies in Tables 1, 2, 3, and 5 and the gas-phase clustering free energies in Table 6. All of the conventional solvation free energies that were used to determine the differential solvation free energies shown in Figure 1 are listed in Table S2 in the Supporting Information. (Tables S3 and S4 list the conventional solvation free energies that were used to determine the differential solvation free energies plotted in Figures 2 and 3.) The straight lines in Figure 1 are best-fit lines for different numbers of clustering methanol molecules. The line for $n = 0$ (no clustering methanol molecules) is the ideal line ($y = x$). The ordinate of the intersection point between each line gives an approximate value for the absolute solvation free energy of the proton, so in all, 6 individual determinations can be made from the data shown in Figure 1 (The 4 straight lines give 6 unique intersection points between different values of n .) To determine the best value of the absolute solvation free energy of the proton, we used the same method of statistical analysis as in our earlier work.⁵⁶ For this, the average intersection ordinate of the i th straight line with all others (Y_i) is given by

$$Y_i = \frac{1}{3} \sum_{j \neq i=0}^3 y_{ij} \quad (27)$$

where

$$y_{ij} = (m_i b_j - m_j b_i) / (m_i - m_j) \quad (28)$$

In the above equation, y_{ij} is the ordinate of the intersection point between the i th and j th straight line, and m_i and b_i are the slope and intercept, respectively, of the i th straight line. We take the best value for the absolute solvation free energy of the proton to equal the unweighted average of the four Y_i values determined using eqs 27 and 28. For the data in Figure 1, this average equals -263.5 kcal/mol. The standard deviation of the 6 values of y_{ij} from the average value of y_{ij} is 2.0 kcal/mol. The individual slopes, intercepts, Y_i and y_{ij} values obtained from the data shown in Figure 1, as well as from the data shown in Figures 2 and 3 (described below), are included as Supporting Information. Note that we could have also used weighted values of y_{ij} according to the probable errors of each of the straight lines i and j . Indeed, the standard deviations of the 3 straight lines in Figure 1 do increase with n (number of clustering methanol molecules). However, this type of analysis would not take into account that the ordinate of the intersections between the lines with larger slopes (low n) are more prone to error because small changes in their slopes will cause the correspondingly largest variations in the value of the ordinate.

Through the use of thermochemical cycle 7 (illustrated in Scheme 7), the absolute solvation free energy can be used to set the normal hydrogen electrode potential according to

$$E_{\text{NHE}}^* = - \frac{\Delta_f G_{298}^0(\text{H}^+) + \Delta G_s^*(\text{H}^+) + \Delta G^{0 \rightarrow *}}{F} \quad (26)$$

where $\Delta_f G_{298}^0(\text{H}^+)$ is the gas-phase free energy of formation of the proton. (The free energy of formation of $\text{H}_2(\text{g})$ and of the electron do not appear in eq 26 because they are equal to zero.) As was pointed out above, the value for the free energy of formation of the proton depends on whether Boltzmann or Fermi-Dirac statistics are used to compute the integrated heat capacity and entropy of the electron. In this article, we have used the electron integrated heat capacity and entropy reported by Bartmess,¹¹⁶ which were obtained using Fermi-Dirac statistics. When these values for the electron integrated heat capacity and entropy are used, the free energy of formation of the gas-phase proton equals 361.7 kcal/mol at 298 K and 1 atm, which when substituted into eq 26 gives a value of -4.34 V for the normal hydrogen electrode potential in methanol. (Using the electron integrated heat capacity and entropy obtained from Boltzmann statistics leads to a value for the normal hydrogen electrode potential 0.04 V more negative than the value above).

6.2. Acetonitrile

Applying eqs 27 and 28 to the data in Figure 2 gives an average value of -260.2 kcal/mol for the absolute solvation free energy of the proton in acetonitrile. The standard deviation of the 6 values of y_{ij} from the average value of y_{ij} is 1.4 kcal/mol. In our initial fit, we discovered that for $n = 1$, the differences between the conventional solvation free energies of anions and of anilinium (9 data points) were systematically more positive than the differences between the conventional solvation free energies of anions and of the remaining cations by ~ 3 kcal/mol. The experimental value for the gas-phase free energy associated with attaching a single acetonitrile molecule to anilinium is -12.0 kcal/mol.^{169,175} We also calculated this free energy at the B97-1/MG3S level of theory, which leads to a value of -14.4 kcal/mol. When the calculated instead of the experimental value for the gas-phase clustering free energy is used to determine the conventional solvation free energy of the anilinium-acetonitrile cluster, the

differences between the conventional solvation free energies of clustered anions and of the anilinium-acetonitrile cluster fall into line with the remaining data points. Because of this, we used the calculated instead of the experimental value for this clustering free energy in our final fit, which is the one shown in Figure 2.

Substituting the value obtained above for the absolute solvation free energy of the proton in acetonitrile (-260.2 kcal/mol) and the value for the free energy of formation of the gas-phase proton obtained using Fermi-Dirac statistics for the integrated heat capacity and entropy of the electron into eq 26 gives a value of -4.48 V for the normal hydrogen electrode potential in acetonitrile.

6.3. DMSO

Applying eqs 27 and 28 to all of the data in Figure 3 gives an average value of -273.7 kcal/mol for the absolute solvation free energy of the proton in DMSO. The standard deviation of the 6 values of y_{ij} from the average value of y_{ij} is 1.8 kcal/mol. For $n = 1$, the differences between the conventional solvation free energies of anions and of pyridinium, which are shown in the lower left-hand corner of Figure 3, tend to fall on a different line than the differences between the conventional solvation free energies of anions and of the remaining cations. Because of this, we removed all of the differences between the conventional solvation free energies of anions and of pyridinium, and then repeated the $n = 1$ fit using the remaining data. The resulting best-fit line is shown as a solid line in Figure 3.

Experimental gas-phase clustering free energies for ion-DMSO clusters containing more than a single DMSO molecule are scarcer than for the other solvents considered in this work. Experimental clustering free energies for $n > 1$ are only available for 4 of the ions in Table 6 (Br^- , Cl^- , I^- , and K^+). The best-fit lines for $n = 2$ and $n = 3$, albeit based on the minimum amount of data points required to determine a statistically meaningful best-fit line, suggest that the best-fit line for $n = 1$ obtained after removing pyridinium (solid $n = 1$ line) is more accurate than the best-fit line obtained using all of the data (dashed $n = 1$ line). Applying eqs 27 and 28 to the slopes and intercepts of the solid $n = 1$ line and the $n = 2$ and $n = 3$ lines in Figure 3 gives an average value of -273.3 kcal/mol for the absolute solvation free energy of the proton in DMSO. The standard deviation of the 6 values of y_{ij} from the average value of y_{ij} is 0.6 kcal/mol.

Substituting the final value obtained above for the absolute solvation free energy of the proton in DMSO (-273.3 kcal/mol) and the value for the free energy of formation of the gas-phase proton obtained using Fermi-Dirac statistics for the integrated heat capacity and entropy of the electron into eq 26 gives a value of -3.92 V for the normal hydrogen electrode potential in DMSO.

7. Accuracy of the Absolute Solvation Free Energies Obtained in This Work

In our previous work where we applied the cluster pair approximation to aqueous solution,⁵⁶ we estimated that the uncertainty associated with the absolute aqueous solvation free energy of the proton is no less than 2 kcal/mol. Based on the uncertainties associated with the experimental and calculated data used in this work, we estimate that uncertainties associated with the absolute solvation free energy of the proton in methanol and acetonitrile are 2 – 3 kcal/mol. For DMSO, where only a limited amount of gas-phase clustering data for $n > 1$ is available, the estimated uncertainty for the absolute solvation free energy of the proton is 3 – 4 kcal/mol.

Because many of the gas-phase clustering free energies for singly clustered polyatomic ions that were used in this work were obtained theoretically (all gas-phase clustering free energies for clusters containing more than a single solvent molecule are experimental), an important

issue is what effect the inclusion of these theoretical data has on the overall accuracy of the results obtained here. In previous work where we determined the absolute aqueous solvation free energy of the proton using the cluster pair approximation,⁵⁶ we showed that augmenting experimental gas-phase clustering data with theoretical data had little effect on the accuracy of the predictions made there. The same level of theory that was used in that work, B97-1/MG3S, was also used here because it has been shown in several places¹⁷⁶⁻¹⁷⁹ to perform well for nonbonded interactions in the gas phase. To further demonstrate the accuracy of this level of theory, we calculated gas-phase clustering free energies for all of the polyatomic ions in Table 7 for which experimental $n = 1$ clustering free energies are available (total of 8 ion-solvent clusters). Excluding the difference between the experimental and calculated value for the anilinium-acetonitrile cluster (the experimental value for this free energy may be in error, see above), the mean unsigned error between the calculated and experimental clustering free energies for the remaining 7 ion-solvent clusters is only 0.26 kcal/mol. The maximum difference is 1.82 kcal/mol, which is the difference between the calculated and experimental gas-phase clustering free energy of the acetate-methanol cluster.

Besides gas-phase clustering free energies for singly clustered ions, calculated values for the solvation free energies of neutral species were used in eqs 15 and 16 to determine conventional solvation free energies. In this work, we used the SM5.43R continuum solvation model to calculate solvation free energies of neutral species. The accuracy of this model has been previously demonstrated in several places,^{12,157} where it has been shown that for typical neutral organic compounds, the solvation free energies calculated using this model are accurate to within ~ 0.5 kcal/mol. Thus, the accuracy of the results obtained here has most likely not been compromised by the inclusion of calculated solvation free energies of neutral species. On the contrary the consistency of the theoretical and experimental data gives us further confidence in the soundness of our approach.

Most of the experimental pK_{sp} values used in this work are sufficiently large so as to safely assume that the solvation free energies of the corresponding salts are fairly representative of the ion pairs in infinitely dilute solution. A notable exception is the experimental pK_{sp} of KBr in DMSO (equal to 0.6), which we used to determine the conventional solvation free energy of K^+ in DMSO. In this case, the ionic strength of the saturated solution is equal to 1 mol/L, whereas the concentration of pure DMSO is 14 mol/L. Thus, for this electrolytic solution the above assumption of a dilute solution does not apply. However, we also determined the conventional solvation free energy of K^+ in DMSO using an experimental value (3.1) for the pK_{sp} of KCl. The conventional solvation free energies of K^+ from these two pK_{sp} values are within 0.3 kcal/mol of one another, suggesting the above concern is not a serious issue.

Finally, it should be noted that in this work, we assume all experimental pK_a values correspond to the equilibrium constant between the neutral and ionic species in their completely dissociated states. Ion pairing is generally negligible in solvents with dielectric constants above 40, whereas its importance depends on the structure of the solvent and the electrolyte for solvents with dielectric constants of 15–20.¹⁸⁰ The solvents considered in this paper have dielectric constants of 32.66 (methanol), 35.94 (acetonitrile), and 46.45 (DMSO), and we will assume that ion pairing is negligible.

8. Previous Estimates of the Absolute Solvation Free Energy of the Proton in Methanol, Acetonitrile, and DMSO

Kalidas and coworkers⁵⁸ have reported values for the transfer free energies of single ions between water and 17 different organic solvent systems, including pure methanol, acetonitrile, and DMSO. Based on a careful evaluation of the available literature data, these authors were able to list recommended values for the transfer free energies of the proton between water and

the above solvents. All of these recommended transfer free energies were obtained using the TATB approximation. The values recommended by these workers for the transfer free energy of the proton between water and methanol, acetonitrile, and DMSO are 2.1, 10.7, and -4.6 kcal/mol, respectively. These transfer free energies, as well as the transfer free energies discussed below, are listed in Table 8. (It is worth noting that the value above for the transfer free energy of the proton between water and DMSO has been used by Pliego and Riveros to determine the absolute solvation free energies of 30 other monovalent ions in DMSO.¹⁸¹)

Combining Tissandier et al.'s value for the absolute aqueous solvation free energy of the proton (-265.9 kcal/mol), with the values for the absolute solvation free energy of the proton obtained here lead to transfer free energies of 2.4, 5.7, and -7.4 kcal/mol for methanol, acetonitrile, and DMSO, respectively. For methanol, the transfer free energy obtained here using the cluster pair approximation is within 0.3 kcal/mol of the value recommended by Kalidas et al. Considering the uncertainties associated with the absolute solvation free energies obtained here, the agreement between the transfer free energy of the proton in DMSO obtained here and that reported by Kalidas et al. is satisfactory. On the other hand, for acetonitrile the agreement between the value obtained here and that reported by Kalidas et al. is not very good (5.0 kcal/mol difference). One reason for this difference is that errors might be present in the experimental data used by Kalidas et al. to derive this transfer free energy. Indeed, unlike most of the transfer free energies recommended by these workers, which are average values taken from several places, the experimental data used to determine the transfer free energy of the proton between water and acetonitrile was taken from a single source.¹⁸² Furthermore, Kalidas et al. list this value as tentative and state that it “must be considered doubtful at this stage.”

It is important to point out that, like the transfer solvation free energies reported in this work, many of the transfer free energies reported by Kalidas et al. are also based on differences between experimental standard reduction potentials and solubility products in water and in nonaqueous solvents (see ref ¹⁸³). The uncertainties in the relative solvation free energies determined using these data are relatively small, whereas for single ions we assign uncertainties that are much larger (2–4 kcal/mol) based on our estimate of the uncertainty associated with the extrathermodynamic assumption used in this article (i.e. the cluster pair approximation) for splitting well-defined sums of solvation free energies into single-ion values. It is not exactly clear what value for the uncertainty should be assigned to the single-ion transfer free energies reported by Kalidas et al., although these authors state that their values are accurate “within the constraints of the TATB assumption” to ± 0.7 kcal/mol. However, the above differences, as well as previous experimental^{41,42,44,46-48} and theoretical work,⁴⁹⁻⁵¹ suggest that the uncertainty associated with single ion solvation free energies obtained using the TATB assumption is larger than 0.7 kcal/mol (and perhaps larger than any of the uncertainties reported here).

Recently, Westphal and Pliego performed high-level gas-phase calculations on Li^+ and Na^+ clusters containing up to four and five explicit DMSO molecules, respectively.¹⁰⁷ Using these clustered ions and a dielectric continuum model, these workers calculated the absolute solvation free energies of Li^+ and Na^+ using an equation similar to eq 19 shown here. Based on the difference between the calculated absolute solvation free energies of these ions, these workers determined the absolute solvation free energy of the proton in DMSO to be -273.2 kcal/mol. Combining this value with Tissandier's value for the absolute aqueous solvation free energy of the proton leads to a transfer free energy of -7.3 kcal/mol, which is in nearly perfect agreement with the value obtained in this work.

Besides the two extrathermodynamic assumptions discussed above (cluster pair and TATB) single-ion solvation free energies have also been estimated based on measurements of the Volta (outer) potential between two different phases brought into contact with one another.^{18,}

184-193 Unlike the solvation free energies obtained using the cluster pair or TATB approximations, the solvation free energies obtained from these types of experiments contain, in addition to the free energy required to couple the solute to the bulk solvent, a free energy term that depends on the surface potentials of the two phases that are brought into contact with one another. (As discussed in the first paragraph of the present article, the solvation free energies resulting from these types of experiments are usually referred to as real solvation free energies to distinguish them from absolute, or Gibbs, solvation free energies, which do not contain the contribution due to the potential of the phase.) Because it is generally agreed that the surface potential of a bulk liquid cannot be directly measured experimentally, one must resort to extrathermodynamic assumptions or theoretical calculations in order to make direct comparisons between real and absolute solvation free energies of single ions. (The distinction between real and absolute solvation free energies does not need to be made when one considers the transfer for pairs of ions with no net charge because the contribution to the solvation free energy due to the potential of the phase cancels out.) For example, Coetzee¹⁹² assigned tentative values to the surface potential of several nonaqueous solvents by combining transfer free energies obtained from measurements of Volta potential differences and those estimated using the TATB approximation.¹⁹⁴

In this article and in our previous work,^{111,195,196} we have assigned values to the normal hydrogen electrode potential using the absolute solvation free energy of the proton, which does not contain the contribution to the free energy due to the potential of the phase. However, in other work (cf. ref¹⁹¹), real solvation free energies, which do contain this contribution, have been used to assign a value to the normal hydrogen electrode potential. In principle, it is correct to use either real or absolute solvation free energies for assigning a value to the normal hydrogen electrode potential, because, as pointed out by Trasatti,¹⁹⁷ there are several different, but equally valid ways to define the normal hydrogen electrode potential. However, reported values for the normal hydrogen electrode potential are often used in conjunction with thermochemical cycle 7 in order to “back out” the absolute value for the solvation free energy of the proton. Thus, from a practical standpoint, one must be aware of whether this value represents a real or an absolute solvation free energy, since these two quantities can differ significantly. (A recent theoretical estimate of the surface potential of bulk water puts this value at $-12.7 \text{ kcal mol}^{-1} e^{-1}$.¹⁹⁸) The distinction between these two quantities can become especially important when absolute solvation free energies obtained from continuum solvation calculations, which do not include the contribution due to the potential of the phase, are unknowingly combined with solvation free energies that *do* contain this contribution. Because of this, for applications that involve the use of solvation free energies obtained from continuum solvation calculations (e.g., calculating absolute acid dissociation constants via thermochemical cycles 4 and 5), we recommend using the absolute solvation free energies reported in this article.

9. Conclusions

Using a combination of experimental gas-phase free energies of formation and solution-phase reduction potentials and acid dissociation constants, conventional solvation free energies of unclustered ions in methanol, acetonitrile, and DMSO were determined. These conventional solvation free energies were then combined with experimental and calculated gas-phase clustering free energies to determine conventional solvation free energies of ion-solvent clusters containing up to three solvent molecules. The cluster pair approximation was applied to the above data in order to determine the absolute solvation free energy of the proton in methanol, acetonitrile, and DMSO. For methanol, the value obtained here agrees fairly well with an earlier value based on the TATB approximation, whereas the agreement is satisfactory for DMSO and not very good for acetonitrile. The values for the absolute solvation free energy of the proton obtained in this work should be useful as standards against which the absolute

solvation free energies of other single ions can be derived. For example, Table 9 shows the absolute single-ion solvation free energies of the ions considered in this work.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institutes of Health training grant for Neuro-physical-computational Sciences, by the U. S. Army Research Office under the Multidisciplinary Research Program of the University Research Initiative through Grant No. DAAD19-02-1-0176, by the National Science Foundation (Grant Nos. CHE06-10183 and CHE03-49122), and by the Office of Naval Research under Grant No. N 00014-05-01-0538.

References

1. Cabani S, Gianni P, Mollica V, Lepori L. *J. Solution Chem* 1981;10:563.
2. Abraham MH, Whiting GS, Fuchs R, Chambers EJ. *J. Chem. Soc., Perkin Trans. 2* 1990:291.
3. Leo, AJ. Masterfile from MedChem Software. BioByte Corp.; Claremont, CA: 1994.
4. Physical/Chemical Property Database (PHYSPROP). SRC Environmental Science Center; Syracuse, NY: 1994.
5. Yaffe D, Cohen Y, Espinosa G, Arenas A, Giralt F. *J. Chem. Inf. Comput. Sci* 2003;43:85. [PubMed: 12546542]
6. Sander, R. Henry's Law Constants.. In: Linstrom, PJ.; Mallard, WG., editors. NIST Chemistry WebBook, NIST Standard Reference Database Number 69. National Institute of Standards and Technology; Gaithersburg, MD: Jun. 2005
7. Kelly CP, Cramer CJ, Truhlar DG. *J. Chem. Theory Comput* 2005;1:1133.
8. Rizzo RC, Aynechi T, Case DA, Kuntz ID. *J. Chem. Theory Comput* 2006;2:128.
9. Li J, Zhu T, Hawkins GD, Winget P, Liotard DA, Cramer CJ, Truhlar DG. *Theor. Chem. Acc* 1999;103:9.
10. Kelly, CP.; Thompson, JD.; Hawkins, GD.; Chambers, CC.; Giesen, DG.; Winget, P.; Cramer, CJ.; Truhlar, DG. Minnesota Solvation Database version 2.0.1. University of Minnesota; Minneapolis, MN: 2006. p. 55455-0431.
11. Cramer, CJ.; Truhlar, DG. Free Energy Calculations in Rational Drug Design. Reddy, MR.; Erion, MD., editors. Kluwer/Plenum; New York: 2001. p. 63
12. Thompson JD, Cramer CJ, Truhlar DG. *J. Phys. Chem. A* 2004;108:6532.
13. Klotz, IM.; Rosenberg, RM. Chemical Thermodynamics. 5th ed.. Wiley; New York: 1994. p. 459
14. Guggenheim EA. *J. Phys. Chem* 1929;33:842.
15. Guggenheim EA. *J. Phys. Chem* 1930;34:1540.
16. Cox BG. *Annu. Repts. Prog. Chem* 1973;A70:249.
17. Hodgkin AL. *Biol. Rev* 1951;26:339.
18. Izmailov NA. *Russ. J. Phys. Chem* 1960;34:1142.
19. Rosseinsky DR. *Chem. Rev* 1965;65:467.
20. Pliego JR Jr. Riveros JM. *Chem. Phys. Lett* 2000;332:597.
21. Llano J, Eriksson LE. *J. Chem. Phys* 2002;117:10193.
22. Bhattacharyya R, Lahiri SC. *Z. Phys. Chem* 2004;218:515.
23. Conway BE. *Annu. Rev. Phys. Chem* 1966;17:481.
24. Parker AJ. *Chem. Rev* 1969;69:1.
25. Popovych O. *Crit. Rev. Anal. Chem* 1970;7:73.
26. Kolthoff IM. *Pure Appl. Chem* 1971;25:305.
27. Conway BE. *J. Solution Chem* 1978;7:721.
28. Marcus Y. *Pure Appl. Chem* 1986;58:1721.

29. Krestov, GA. Thermodynamics of Solution. Ellis Horwood; New York: 1991.
30. Coe JV. Int. Rev. Phys. Chem 2001;20:33.
31. Born M. Z. Physik 1920;1:45.
32. Cramer, CJ.; Truhlar, DG. Rev. Comp. Chem. Boyd, DB.; Lipkowitz, KB., editors. 6. VCH Publishers; New York: 1995. p. 1
33. Alexander R, Parker AJ. J. Am. Chem. Soc 1967;89:5539.
34. Bernal J, Fowler R. J. Chem. Phys 1933;1:515.
35. Fajans K, Johnson O. J. Am. Chem. Soc 1942;64:688.
36. Vewrey TE. Rec. Trav. Chim 1942;61:127.
37. Mishchenko KP. Russ. J. Phys. Chem 1952;26:1736.
38. Grunwald E, Baughman G, Kohnstam C. J. Am. Chem. Soc 1960;82:5801.
39. Breck WG, Lin J. Trans. Faraday Soc 1965;69:222.
40. Popovych O, Dill AJ. Anal. Chem 1966;38:588.
41. Coetzee JF, Sharpe WR. J. Phys. Chem 1971;75:3141.
42. Jolicoeur C, Dinh The N, Cabana A. Can. J. Chem 1971;49:2008.
43. Alexander R, Parker AJ, Sharpe JH, Waghorne WE. J. Am. Chem. Soc 1972;94:1148.
44. Jolicoeur C, Friedman HL. J. Solution Chem 1974;3:15.
45. Kim JI. J. Phys. Chem 1978;82:191.
46. Farmer RM, Sasaki Y, Popov AI. Aust. J. Chem 1983;36:1785.
47. Shao Y, Stewart AA, Girault HH. J. Chem. Soc. Faraday Trans 1991;87:2593.
48. Stangret J, Kamienska-Piotrowicz E. J. Chem. Soc. Faraday Trans 1997;93:3463.
49. Schurhammer R, Wipff G. J. Mol. Struct. (THEOCHEM) 2000;500:139.
50. Schurhammer R, Wipff G. J. Phys. Chem. A 2000;104:11159.
51. Schurhammer R, Engler E, Wipff G. J. Phys. Chem. B 2001;105:10700.
52. Wachter W, Buchner R, Hefter G. J. Phys. Chem. B 2006;110:5147. [PubMed: 16526759]
53. Tissandier MD, Cowen KA, Feng WY, Gundlach E, Cohen MJ, Earhart AD, Coe JV. J. Phys. Chem. A 1998;102:7787.
54. Tuttle TR Jr. Malaxos S, Coe JV. J. Phys. Chem. A 2002;106:925.
55. Bartels DM, Takahashi K, Cline JA, Marin TW, Jonah CD. J. Phys. Chem. A 2005;109:1299. [PubMed: 16833444]
56. Kelly CP, Cramer CJ, Truhlar DG. J. Phys. Chem. B 2006;110:16066. [PubMed: 16898764]
57. Camaioni DM, Scherdtfeger CA. J. Phys. Chem. A 2005;109:10795. [PubMed: 16863129]
58. Kalidas C, Hefter G, Marcus Y. Chem. Rev 2000;100:819. [PubMed: 11749253]
59. Tunon I, Rinaldi D, Ruiz-Lopez MF, Rivail JL. J. Phys. Chem 1995;99:3798.
60. Florián J, Warshel A. J. Phys. Chem. B 1997;101:5583.
61. Schüürman G, Cossi M, Barone V, Tomasi J. J. Phys. Chem. A 1998;102:6706.
62. da Silva CO, da Silva EC, Nascimento MAC. J. Phys. Chem. A 1999;103:11194.
63. Vogel HJ, Juffer AH. Theor. Chem. Acct 1999;101:159.
64. Chen I-J, MacKerrel AD Jr. Theor. Chem. Acct 2000;103:483.
65. Meijías JA, Lago S. J. Chem. Phys 2000;113:7306.
66. Pliego JR Jr. Riveros JM. J. Phys. Chem. B 2000;104:5155.
67. Silva CO, da Silva EC, Nascimento MAC. J. Phys. Chem. A 2000;104:2402.
68. Topol IA, Tawa GJ, Caldwell RA, Eissenstat MA. J. Phys. Chem. A 2000;104:9619.
69. Jang YH, Sowers LC, Çagin T, Goddard WA III. J. Phys. Chem. A 2001;105:274.
70. Liptak MD, Shields GC. Int. J. Quantum Chem 2001;85:727.
71. Liptak MD, Shields GC. J. Am. Chem. Soc 2001;123:7314. [PubMed: 11472159]
72. Ohno K, Kamiya N, Asakawa N, Inoue Y, Sakurai M. Chem. Phys. Lett 2001;341:387.
73. Toth AM, Liptak MD, Phillips DL, Shields GC. J. Chem. Phys 2001;114:4595.
74. Zhan C-G, Dixon DA. J. Phys. Chem. A 2001;105:11534.

75. Adam K. J. Phys. Chem. A 2002;106:11963.
76. Chipman DM. J. Phys. Chem. A 2002;106:7413.
77. Davies JE, Doltsinis NL, Kirby AJ, Roussev CD, Sprik M. J. Am. Chem. Soc 2002;124:6594. [PubMed: 12047179]
78. Feierberg I, Åqvist J. Theor. Chem. Acct 2002;108:71.
79. Klicic JJ, Friesner RA, Liu S-Y, Guida WC. J. Phys. Chem. A 2002;106:1327.
80. Li H, Hains AW, Everts JE, Robertson AD, Jensen JH. J. Phys. Chem. B 2002;106:1327.
81. Liptak MD, Gross KC, Seybold PG, Feldgus S, Shields GC. J. Am. Chem. Soc 2002;124:6421. [PubMed: 12033873]
82. Lopez X, Schaefer M, Dejaegere A, Karplus M. J. Am. Chem. Soc 2002;124:5010. [PubMed: 11982365]
83. Satchell JF, Smith BJ. Phys. Chem. Chem. Phys 2002;4:4314.
84. Zhan C-G, Dixon DA. J. Phys. Chem. A 2002;106:9737.
85. Asthagiri D, Pratt LR, Ashbaugh HS. J. Chem. Phys 2003;119:2702.
86. Camaioni DM, Dupuis M, Bentley J. J. Phys. Chem. A 2003;107:5778.
87. Grossfield A, Ren P, Ponder JW. J. Am. Chem. Soc 2003;125:15671. [PubMed: 14664617]
88. Jang YH, Goddard WA III, Noyes KT, Sowers LC, Hwang S, Chung DS. J. Phys. Chem. B 2003;107:344.
89. Klamt A, Eckert F, Diedenhofen M, Beck ME. J. Phys. Chem. A 2003;107:9380.
90. Li G, Cui Q. J. Phys. Chem. B 2003;107:14521.
91. Mujika JI, Mercero JM, Lopez X. J. Phys. Chem. A 2003;107:6099.
92. Almerindo GI, Tondo DW, Pliego JR Jr. J. Phys. Chem. A 2004;108:166.
93. Barone V, Improta R, Rega N. Theor. Chem. Acct 2004;111:237.
94. da Silva CO, Barbosa AGH, da Silva ET, da Silva ELL, Nascimento MAC. Theor. Chem. Acct 2004;111:231.
95. Fu Y, Liu L, Li R-Q, Guo Q-X. J. Am. Chem. Soc 2004;126:814. [PubMed: 14733556]
96. Namazian M, Halvani S, Noorbala MR. J. Mol. Struct. (THEOCHEM) 2004;711:13.
97. Quenneville J, Popovic DM, Stucherbrukhov AA. J. Phys. Chem. B 2004;108:18383.
98. Rempe S, Asthagiri D, Pratt LR. Phys. Chem. Chem. Phys 2004;6:1966.
99. Schmidt am Busch M, Knapp EW. ChemPhysChem 2004;5:1513. [PubMed: 15535550]
100. Simonson T, Carlsson J, Case DA. J. Am. Chem. Soc 2004;126:4167. [PubMed: 15053606]
101. Gao D, Svoronos P, Wong PK, Maddalena D, Hwang J, Walker H. J. Phys. Chem. A 2005;109:10776. [PubMed: 16863127]
102. Han J, Deming RL, Tao F-M. J. Phys. Chem. A 2005;109:1159. [PubMed: 16833426]
103. Jensen JH, Li H, Robertson AD, Molina PA. J. Phys. Chem. A 2005;109:6634. [PubMed: 16834015]
104. Kaminski GA. J. Phys. Chem. B 2005;109:5884. [PubMed: 16851640]
105. Popovic DM, Quenneville J, Stucherbrukhov AA. J. Phys. Chem. B 2005;109:3616. [PubMed: 16851400]
106. Takano Y, Houk KN. J. Chem. Theory Comput 2005;1:70.
107. Westpahl E, Pliego JR Jr. J. Chem. Phys 2005;123:074508. [PubMed: 16229602]
108. Eckert F, Klamt A. J. Comput. Chem 2006;27:11. [PubMed: 16235262]
109. Gutowski KE, Dixon DA. J. Phys. Chem. A 2006;110:8840. [PubMed: 16836448]
110. Kastenholz MA, Hünenberger H. J. Chem. Phys 2006;124:224501. [PubMed: 16784292]
111. Kelly CP, Cramer CJ, Truhlar DG. J. Phys. Chem. A 2006;110:2493. [PubMed: 16480309]
112. Lamoureux G, Roux B. J. Phys. Chem. B 2006;110:3308. [PubMed: 16494345]
113. Lewis, GN.; Randall, M.; Pitzer, KS.; Brewer, L. Thermodynamics. 2nd ed.. McGraw-Hill; New York: 1961. p. 399
114. McQuarrie, DA. Statistical Mechanics. University Science Books; Sausalito, California: 2000.
115. Lias SG, Bartness JE, Liebman JF, Holmes JL, Levin RD, Mallard WG. J. Phys. Chem. Ref. Data 1988;16(Suppl 1)

116. Bartmess JE. J. Phys. Chem 1994;98:6420.
117. Toman RC. J. Am. Chem. Soc 1921;43:1592.
118. Lewis GN, Latimer WM. J. Am. Chem. Soc 1922;44:1008.
119. Fermi E. Z. Phys 1926;36:902.
120. Sommerfeld A. Z. Phys 1928;47:1.
121. Strehlow, H. Electrode Potentials in Non-Aqueous Solvents.. In: Lagowski, JJ., editor. The Chemistry of Nonaqueous Solvents. Academic Press; New York: 1966. p. 134
122. Kolthoff IM, Chantooni MKJ. J. Phys. Chem 1972;76:2024.
123. Lide, DR., editor. CRC Handbook of Chemistry and Physics. 87th ed.. CRC Press; Boca Raton, FL: 2006.
124. Fawcett WR. J. Phys. Chem. B 1999;103:11181.
125. Ben-Naim, A. Solvation Thermodynamics. Plenum; New York: 1987.
126. Wagman DD, Evans WH, Parker VB, Schumm RH, Halow I, Bailey SM, Churney KL, Nutall RL. J. Phys. Chem. Ref. Data 1982;11(Suppl 2)
127. Salomon, M. Physical Chemistry of Organic Solvent Systems. Covington, AK.; Dickinson, T., editors. Plenum Press; New York: 1973. p. 190
128. Janz, GJ.; Tomkins, RPT. Nonaqueous Electrolytes Handbook. II. Academic Press; New York: 1973. p. 115
129. Labban AKS, Marcus Y. J. Solution Chem 1991;20:221.
130. Kenttämä J. Suomen Kemistilehti 1960;B33:179.
131. Gopal R, Jha JS. Ind. J. Chem 1974;12:618.
132. In several earlier papers where we used this thermochemical cycle,^{7,56,111} the solvation free energies appearing in the vertical legs were denoted by a superscript asterisk, meaning transfer from a standard-state gas-phase concentration of 1 mol/L to a standard-state liquid-phase concentration of 1 mol/L, and the symbol for the free energy change associated with moving from a standard-state gas-phase pressure of 1 atm to a liquid-phase concentration of 1 mol/L was left out of the diagrams (but not the equations) for simplicity. In this paper, we use an equivalent, and perhaps simpler convention in which the solvation free energies appearing in the vertical legs are denoted by a superscript open circle. The equations resulting from the thermochemical cycles that appear in this article are identical to those equations appearing in our earlier papers.
133. Ogston AG. Trans. Faraday Soc 1936;32:1679.
134. Kolthoff IM, Lingane JJ, Larson W. J. Am. Chem. Soc 1938;60:2512.
135. Shedlovsky T, Kay RL. J. Phys. Chem 1956;60:151.
136. Kolthoff IM, Bruckenstein S, Chantooni MKJ. J. Am. Chem. Soc 1961;83:3927.
137. Julliard J, Dondon M-L. Bull. Soc. Chim. Fr 1963:2535.
138. Coetzee JF, Padmanabhan GR. J. Phys. Chem 1965;69:3193.
139. Kononov OM. Zh. Fiz. Khim 1965;39:693.
140. Julliard J. Bull. Soc. Chim. Fr 1966:1727.
141. Kolthoff IM, Chantooni MKJ. J. Phys. Chem 1966;70:856.
142. Rochester CH. Trans. Faraday Soc 1966;62:355.
143. Kolthoff IM, Chantooni MKJ, Bhowmik S. Anal. Chem 1967;39:315.
144. Kolthoff IM, Chantooni MKJ, Bhowmik S. Anal. Chem 1967;39:1627.
145. Rochester CH. J. Chem. Soc. B 1967:33.
146. Chantooni MKJ, Kolthoff IM. J. Am. Chem. Soc 1968;90:3005.
147. Kolthoff IM, Chantooni MKJ, Bhowmik S. J. Am. Chem. Soc 1968;90:23.
148. Bolton PD, Rochester CH, Rossall B. Trans. Faraday Soc 1970;66:1348.
149. Leung CS, Grunwald E. J. Am. Chem. Soc 1970;74:696.
150. Chantooni MKJ, Kolthoff IM. J. Phys. Chem 1975;79:1176.
151. Kolthoff IM, Chantooni MKJ. J. Am. Chem. Soc 1975;97:1376.
152. Kolthoff IM, Chantooni MKJ. J. Am. Chem. Soc 1976;98:5063.
153. Jasinski T, El-Harakany AA, Halaka FG, Sadek H. Croat. Chem. Acta 1978;51:1.

154. Chantooni MKJ, Kolthoff IM. *Anal. Chem* 1979;51:133.
155. Bordwell FG. *Acc. Chem. Res* 1988;21:456.
156. Rived F, Roses M, Bosch E. *Anal. Chim. Acta* 1998;374:309.
157. Thompson DJ, Cramer CJ, Truhlar DG. *Theor. Chem. Acc* 2005;113:107.
158. Adamo C, Barone V. *J. Chem. Phys* 1998;108:664.
159. Hehre, WJ.; Radom, L.; Schleyer, P. v. R.; Pople, JA. *Ab Initio Molecular Orbital Theory*. Wiley; New York: 1986.
160. Easton RE, Giesen DJ, Welch A, Cramer CJ, Truhlar DG. *Theor. Chim. Acta* 1996;93:281.
161. Li J, Cramer CJ, Truhlar DG. *Theor. Chim. Acta* 1998;99:192.
162. Thompson JD, Winget P, Truhlar DG. *PhysChemComm* 2001;4:72.
163. Chamberlin, AC.; Kelly, CP.; Thompson, JD.; Xidos, JD.; Li, J.; Hawkins, GD.; Winget, P.; Zhu, T.; Rinaldi, D.; Liotard, DA.; Cramer, CJ.; Truhlar, DG.; Frisch, MJ. *MN-GSM*, version 6.0. University of Minnesota; Minneapolis, MN: 2006. p. 55455-0431.
164. Frisch, MJ.; Trucks, GW.; Schlegel, HB.; Scuseria, GE.; Robb, MA.; Cheeseman, JR.; Montgomery, JA., Jr.; Vreven, T.; Kudin, KN.; Burant, JC.; Millam, JM.; Iyengar, SS.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalamani, G.; Rega, N.; Petersson, GA.; Nagatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, JE.; Hratchian, HP.; Cross, JB.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, RE.; Yazyev, O.; Austin, AJ.; Cammi, R.; Pomelli, C.; Ochterski, JW.; Ayala, PY.; Morokuma, K.; Voth, GA.; Salvador, P.; Dannenberg, JJ.; Zakrzewski, VG.; Dapprich, S.; Daniels, AD.; Strain, MC.; Farkas, O.; Malick, DK.; Rabuck, AD.; Raghavachari, K.; Foresman, JB.; Ortiz, JV.; Cui, Q.; Baboul, AG.; Clifford, S.; Cioslowski, J.; Stefanov, BB.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, RL.; Fox, DJ.; Keith, T.; Al-Laham, MA.; Peng, CY.; Nanayakkara, A.; Challacombe, M.; Gill, PMW.; Johnson, B.; Chen, W.; Wong, MW.; Gonzalez, C.; Pople, JA. *Gaussian 03*, revision D.01. Gaussian, Inc.; Wallingford, CT: 2003.
165. Lias, SG.; Bartness, JE.; Liebman, JF.; Holmes, JL.; Levin, RD.; Mallard, WG. *Ion Energetics Data*. In: Linstrom, PJ.; Mallard, WG., editors. *NIST Chemistry WebBook*; NIST Standard Reference Database Number 69. National Institute of Standards and Technology; Gaithersburg MD: Mar. 2003
166. Winget P, Hawkins GD, Cramer CJ, Truhlar DG. *J. Phys. Chem. B* 2000;104:4726.
167. Keepports D. *J. Chem. Educ* 2005;82:999.
168. Lewis, GN.; Randall, M.; Pitzer, KS.; Brewer, L. *Thermodynamics*. 2nd ed.. McGraw-Hill; New York: 1961. p. 272
169. Meot-Ner, MM.; Lias, SG. *Binding Energies Between Ions and Molecules, and the Thermochemistry of Cluster Ions*. In: Linstrom, PJ.; Mallard, WG., editors. *NIST Chemistry WebBook*, NIST Standard Reference Database Number 69. National Institute of Standards and Technology; Gaithersburg, MD: Mar. 2003
170. Hamprecht FA, Cohen AJ, Tozer DJ, Handy NC. *J. Chem. Phys* 1998;109:6264.
171. Fast PL, Sanchez ML, Truhlar DG. *Chem. Phys. Lett* 1999;306:407.
172. Cramer, CJ. *Essentials of Computational Chemistry: Theories and Models*. Wiley & Sons; Chichester, U.K.: 2002. p. 321
173. Klots CE. *J. Phys. Chem* 1981;85:3585.
174. A full description of the steps required in going from eq 23 to eq 24 is given in reference ⁵⁵.
175. Meot-Ner (Mautner) M, El-Shall S. *J. Am. Chem. Soc* 1986;108:4386.
176. Zhao Y, Truhlar DG. *J. Chem. Theory. Comput* 2005;1:415.
177. Dahlke EE, Truhlar DG. *J. Phys. Chem. B* 2005;109:15677. [PubMed: 16852988]
178. Zhao Y, Schultz NE, Truhlar DG. *J. Chem. Theory Comput* 2006;2:364.
179. Johnson ER, DiLabio GA. *Chem. Phys. Lett* 2006;419:333.
180. Reichardt, C. *Solvents and Solvent Effects in Organic Chemistry*. 3rd ed.. Wiley-VCH; Weinheim: 2003. p. 48
181. Pliego JR Jr, Riveros JM. *Phys. Chem. Chem. Phys* 2002;4:1622.

182. The reference given by Kalidas et al. for the experimental transfer free energy of the proton between water and acetonitrile is: Das K, Das AK, Kundu K. *Electrochim. Acta* 1981;26:471. This value was also reported in: Cox BG, Hedwig GR, Parker AJ, Watts DW. *Aust. J. Chem* 1974;27:477.
183. Popovych O. *J. Phys. Chem* 1984;88:4167.
184. Hush NS. *Aust. J. Sci. Res., Ser. A* 1948;1:480.
185. Randles JEB. *Trans. Faraday Soc* 1956;52:1573.
186. Case B, Hush NS, Parsons R, Peover ME. *J. Electroanal. Chem* 1965;10:360.
187. Case B, Parsons R. *Trans. Faraday Soc* 1967;63:1224.
188. Zagórska I, Koczorowski Z. *Rocz. Chem* 1970;44:1559.
189. Parsons R, Rubin BT. *J. Chem. Soc. Faraday Trans* 1974;70:1636.
190. Gomer R, Tryson G. *J. Chem. Phys* 1977;66:4413.
191. Trasatti S. *Pure Appl. Chem* 1986;58:955.
192. Coetzee JF, Dollard WJ, Istone WK. *J. Solution Chem* 1991;20:957.
193. Parfenyuk VI, Chankina TI. *Russ. J. Phys. Chem* 2001;38:326.
194. Marcus Y. *Pure Appl. Chem* 1983;55:977.
195. Lewis A, Bumpus JA, Truhlar DG, Cramer CJ. *J. Chem. Educ* 2004;81:596.
196. Winget P, Cramer CJ, Truhlar DG. *Theor. Chem. Acc* 2004;112:217.
197. Trasatti S. *Electrochimica Acta* 1990;35:269.
198. Sokhan VP, Tildesley DJ. *Mol. Phys* 1997;92:625.

Figure 1.

Half the difference between conventional solvation free energies of anions and of cations in methanol for cluster ions containing up to three methanol molecules plotted vs half the difference between conventional solvation free energies of anions and of cations in methanol containing no methanol molecules. Differences between cluster ions containing one methanol molecule are plotted as circles, two as squares, and three as triangles. The ordinate of the average intersection point between each of the straight lines is an estimate of the value for the absolute solvation free energy of the proton. The asterisk on the ordinate is the absolute solvation free energy that results from using the TATB approximation.

Figure 2.
Same as Figure 1, except for acetonitrile.

Figure 3.

Same as Figures 1 and 2, except for DMSO. The dashed line is the best-fit line computed using differential solvation free energies between all anions and all cations, containing up to a single DMSO molecule (open circles). The corresponding solid line is the best-fit line computed after excluding differential solvation free energies between anions and pyridinium (lower leftmost open circles). The slope and intercept obtained after excluding pyridinium (solid line) was used to obtain the final value of the absolute solvation free energy of the proton in DMSO.

SCHEME 1.
Thermochemical Cycle 1

SCHEME 2.
Thermochemical Cycle 2

SCHEME 3.
Thermochemical Cycle 3

SCHEME 4.
Thermochemical Cycle 4

SCHEME 5.
Thermochemical Cycle 5

SCHEME 6.
Thermochemical Cycle 6

SCHEME 7.
Thermochemical Cycle 7

TABLE 1

Gas-Phase Free Energies of Formation (kcal/mol),^a Standard Reduction Potentials (Volts) and Conventional Solvation Free Energies (kcal/mol) of Metal Ions in Different Solvents

M ⁺	$\Delta_f G_{298}^\circ [M^+ (g)]^d$	E_c^{*b}			$\Delta G_S^{*,con}(M^+)^c$		
		CH ₃ OH ^e	CH ₃ CN	DMSO ^e	CH ₃ OH	CH ₃ CN	DMSO
Na ⁺	136.4	-2.728	-2.87 ^e		162.4	159.1	
K ⁺	114.1	-2.921	-3.16 ^e		180.2	174.7	
Rb ⁺	108.5	-2.912	-3.17, -3.316 ^{ef}		186.0	178.4	
Cs ⁺	101.1		-3.16 ^e			187.8	
Tl ⁺	175.6	-0.379	-0.682 ^f	-0.352	177.3	170.4	178.0
Ag ⁺	232.9	+0.764	+0.23, +0.096 ^{ef}	+0.680, +0.668	146.4	132.4	144.3

^a $\Delta_f H^\circ = 4.184$ kJ.

^b Standard reduction potential for a standard-state solution-phase concentration of 1 mol/L. If more than a single experimental E_c^* value is listed, the average value was used to determine the conventional solvation free energy.

^c Based on the convention that $\Delta G_S^{*,con}(H^+) = 0$.

^d Free energy of formation obtained following the electron convention and Fermi-Dirac statistics for the electron integrated heat capacity and entropy,¹¹⁶ for a standard-state gas-phase pressure of 1 atm.

^e Reference 121.

^f Reference 122.

TABLE 2

Gas-Phase Free Energies of Formation (kcal/mol), Standard Reduction Potentials (Volts), and Conventional Solvation Free Energies (kcal/mol) of Halide Ions in Methanol

X^-	$\Delta_f G_{298}^0[X^-(g)]^a$	$\Delta_f G_{298}^0[AgX(cr)]^b$	E_c^{*c}	$\Delta G_S^{*,con}(X^-)^d$
Cl^-	-56.4	-26.2	-0.0221	-334.8
Br^-	-56.2	-23.2	-0.1507	-329.0
I^-	-52.1	-15.8	-0.32987	-321.6

^aFree energy of formation obtained following the electron convention and Fermi-Dirac statistics for the electron integrated heat capacity and entropy, 116 for a standard-state gas-phase pressure of 1 atm.

^bExperimental values were taken from ref 126, and adjusted to a standard-state gas-phase pressure 1 atm using the procedure described in the Supporting Information.

^cStandard reduction potential for a solution-phase concentration of 1 mol/L. Values were taken from reference 127.

^dBased on the convention that $\Delta G_S^{*,con}(H^+) = 0$.

TABLE 3
Solubility Products and Conventional Solvation Free Energies (kcal/mol) of Metal Halides in Different Solvents

MX	$\Delta G_{\text{int}}^{\text{TMX}}$ (cr)] ^d	pK_{sp}^a			$\Delta G_S^*(M^+) + \Delta G_S^*(X^-)$			$\Delta G_S^*(M^+)^{b,c}$			$\Delta G_S^*(X^-)^c$		
		CH ₃ OH	CH ₃ CN	DMSO	CH ₃ OH	CH ₃ CN	DMSO	CH ₃ OH	CH ₃ CN	DMSO	CH ₃ OH	CH ₃ CN	DMSO
KCl	-155.5	3.1	8.0	3.1	-155.0	-148.5	-155.1	180.2	174.7	180.9 ^e	-335.3	-323.2	
NaCl	-171.8		8.3	2.9		-164.3	-171.6		159.1	164.4 ^f		-323.4	
AgCl	-202.8	13.3	13.2	10.8	-188.5	-188.7	-191.9	146.4	132.4	144.3	-334.8	-321.1	-336.2
TlCl	-163.4	5.6	13.2	6.9	-159.5		-157.8	177.3		178.0	-336.9		-335.8
KBr	-148.9	2.5	5.7	0.6 ^g	-149.3	-145.0	-151.9	180.2	174.7	181.2 ^h	-329.6	-319.7	
RbBr	-143.6	2.7	5.5		-143.7	-140.0		186.0	178.4		-329.7	-318.4	
AgBr	-199.9	15.5	13.6	11.0	-182.7	-185.3	-188.8	146.4	132.4	144.3	-329.0	-317.7	-333.1
CsBr	-138.4		5.8			-134.3			187.8		-322.1		
TlBr	-159.4	8.2	13.0		-152.1			177.3			-329.4		
AgI	-196.7	18.5	14.5	12.1	-175.4	-180.8	-184.1	146.4	132.4	144.3	-321.7	-313.2	-328.4

^a Average value from reference 122, unless indicated otherwise.

^b From Table 1, unless indicated otherwise.

^c Based on the convention that $\Delta G_S^*(\text{con}(\text{H}^+)) = 0$.

^d Gas-phase lattice free energy, calculated using eq 14.

^e Calculated from pK_{sp} (KCl) and average $\Delta G_S^*(\text{con}(\text{Cl}^-))$ value from this table.

^f Calculated from pK_{sp} (NaCl) and average $\Delta G_S^*(\text{con}(\text{Cl}^-))$ value from this table.

^g Average value, see text.

^h Calculated from pK_{sp} (KBr) and $\Delta G_S^*(\text{con}(\text{Br}^-))$ value from this table.

TABLE 4

Acid Dissociation Constants in Different Solvents

ion	neutral	pK_a^a		
		CH ₃ OH	CH ₃ CN	DMSO
NO ₂ ⁻	nitrous acid			7.5
NO ₃ ⁻	nitric acid		8.9 ¹³⁶	
CH ₃ CO ₂ ⁻	acetic acid	9.5, ¹⁴⁰ 9.6, ¹³⁵ 9.6, ¹⁴⁸ 9.6, ¹⁵⁶ 9.7, ¹⁵⁴ 9.8 ¹³⁹	22.3, ¹⁴⁷ 22.3, ¹⁵¹ 22.3 ¹⁵³	12.3
CH ₃ CH ₂ CO ₂ ⁻	propionic acid	9.7, ¹⁴⁰ 9.7 ¹⁵⁶		
NCCH ₂ CO ₂ ⁻	cyanoacetic acid		18.0 ¹⁵²	
CF ₃ CO ₂ ⁻	trifluoroacetic acid		12.7 ¹⁵³	
C ₆ H ₅ CO ₂ ⁻	benzoic acid	9.3, ¹⁴⁰ 9.3, ¹⁵⁶ 9.4, ¹³⁴ 9.4, ¹³⁷ 9.4, ¹⁴⁹ 9.5 ¹³⁹	20.7, ¹⁴² 20.7 ¹⁴¹	11.0
<i>p</i> -FC ₆ H ₄ CO ₂ ⁻	4-fluorobenzoic acid	9.2 ¹⁵⁶		
C ₆ H ₅ O ⁻	phenol	14.3, ¹⁵⁶ 14.4, ¹⁴² 14.5 ¹⁴⁸	26.7, ¹³⁸ 27.2 ¹⁴³	18.0
<i>p</i> -NO ₂ C ₆ H ₄ O ⁻	4-nitrophenol	11.1, ¹⁴⁰ 11.3, ¹⁵⁶ 11.5 ¹⁴⁸		
CH ₃ O ⁻	methanol			29.0
C ₂ H ₅ O ⁻	ethanol			29.8
NH ₄ ⁺	ammonia	10.8 ¹⁵⁶	16.5 ¹³⁸	10.5
CH ₃ NH ₃ ⁺	methylamine	11.0 ¹⁵⁶	18.4 ¹³⁸	11.0
(CH ₃) ₃ NH ⁺	trimethylamine	9.8 ¹⁵⁶		
C ₅ H ₁₀ NH ₂ ⁺	piperidine	11.1 ¹⁵⁶		
pyridineH ⁺	pyridine	5.4, ¹⁴² 5.4 ¹⁵⁶	12.3, ¹³⁸ 12.3 ¹⁴⁶	3.5
C ₆ H ₅ NH ₃ ⁺	aniline	6.1 ¹⁵⁶	10.6, ¹³⁸ 10.7, ¹⁴⁴ 10.7 ¹⁴⁶	3.6

^aFor anions, pK_a of the neutral species. For cations, pK_a of the ionic species. For methanol and acetonitril the literature reference for each experimental value is listed to the right of the value in superscript. For DMSO, a experimental values were taken from reference ¹⁵⁵.

TABLE 5

Absolute (Neutral Species) and Conventional (Ionic Species) Solvation Free Energies of Brønsted-Lowry Acids and Bases (kcal/mol)

ion	neutral	ΔG_g^{ad}	pK_a^a			$\Delta G_S^* \text{ (neutral)}^b$			$\Delta G_S^{*,con} \text{ (ion)}^c$		
			CH ₃ OH	CH ₃ CN	DMSO	CH ₃ OH	CH ₃ CN	DMSO	CH ₃ OH	CH ₃ CN	DMSO
NO ₂ ⁻	nitrous acid	333.7			7.5			-4.2		-329.6	
NO ₃ ⁻	nitric acid	317.8					-5.7			-313.3	
CH ₃ CO ₂ ⁻	acetic acid	341.4			12.3	-7.2	-6.5	-7.4	-337.3	-319.5	-333.9
CH ₃ CH ₂ CO ₂ ⁻	propionic acid	340.4	9.7			-7.1			-336.2		
NCCH ₂ CO ₂ ⁻	cyanoacetic acid	323.7	9.7				-8.5			-309.6	
CF ₃ CO ₂ ⁻	trifluoroacetic acid	316.7					-7.4			-308.7	
C ₆ H ₅ CO ₂ ⁻	benzoic acid	333.0	9.4		11.0	-9.7	-9.4	-10.3	-331.8	-316.1	-330.2
<i>p</i> -FC ₆ H ₄ CO ₂ ⁻	4-fluorobenzoic acid	330.0	9.2			-8.6			-328.0		
C ₆ H ₅ O ⁻	phenol	342.9	14.4		18.0	-8.2	-7.5	-8.1	-333.5	-315.6	-328.4
<i>p</i> -NO ₂ C ₆ H ₄ O ⁻	4-nitrophenol	320.9	11.3			-11.0			-318.5		
CH ₃ O ⁻	methanol	375.0			29.0			-4.8		-342.2	
C ₂ H ₅ O ⁻	ethanol	371.3			29.8			-5.3 ^f		-337.9	
NH ₄ ⁺	ammonia	195.7	10.8	16.5	10.5	-5.0	-4.5	-4.3	177.9	170.7	179.0
CH ₃ NH ₃ ⁺	methylamine	206.6	11.0	18.4	11.0	-4.6	-3.7	-3.6	188.9	179.8	189.9
(CH ₃) ₃ NH ⁺	trimethylamine	219.4	9.8			-5.0			203.0		
C ₃ H ₁₀ NH ₂ ⁺	piperidine	220.0	11.1			-6.5			200.3		
pyridineH ⁺	pyridine	214.7	5.4	12.3	3.5	-6.4	-6.1	-5.6	202.9	193.8	206.3
C ₆ H ₅ NH ₃ ⁺	aniline	203.3	6.1	10.7	3.6	-7.4	-7.3	-7.1	189.5	183.4	193.2

^eThis conventional solvation free energy was not used.

^a Average value from Table 4.

^b Calculated (SM5.43R/*m*PW1PW/6-31+G(d,p)//*m*PW1PW/MIDI¹) value, unless indicated otherwise.

^c Based on the convention that $\Delta G_S^{*,con} \text{ (H}^+) = 0$.

^d For anions, gas-phase acidity of the neutral species. For cations, gas-phase acidity of the ionic species. Experimental data are from reference 169.

^f Reference 10.

TABLE 6

Solvation Free Energies of Different Solutes in Their Pure Liquids

solvent	MW (g/mol)	ρ (g/mL) ^a	P_{vapor} (atm) ^b	ΔG_{self}^* (kcal/mol)
CH ₃ OH	32.04	0.7914	0.166	-4.84
CH ₃ CN	41.05	0.7857	0.120	-4.88
DMSO	78.13	1.0955	0.00798	-7.66

^aReference 123.^bReference 166.

TABLE 7
Gas-Phase Clustering Free Energies^a (kcal/mol) of Ions with Various Solvents

	CH ₃ OH			CH ₃ CN			DMSO		
	(0,1)	(1,2)	(2,3)	(0,1)	(1,2)	(2,3)	(0,1)	(1,2)	(2,3)
Cl ⁻	-10.0	-7.0	-4.7	-9.5	-7.1	-5.0	-12.5	-8.9	-6.0
Br ⁻	-8.4	-6.3	-4.2	-8.7	-6.2	-3.9	-10.9	-7.8	-5.4
I ⁻	-5.9	-4.4	-3.1	-6.6	-4.6	-3.0	-9.2	-6.2	-4.1
NO ₂ ⁻							-8.7		
NO ₃ ⁻				[-7.3]					
CH ₃ CO ₂ ⁻	-10.5			-9.4			[-10.2]		
CH ₃ CH ₂ CO ₂ ⁻	[-8.0]								
NCCH ₂ CO ₂ ⁻				[-7.0]					
CF ₃ CO ₂ ⁻				[-9.0]					
C ₆ H ₅ CO ₂ ⁻	[-7.3]			[-7.6]			[-7.7]		
<i>p</i> -FC ₆ H ₄ CO ₂ ⁻	[-6.9]								
C ₆ H ₅ O ⁻	[-8.1]						[-7.9]		
<i>p</i> -NO ₂ C ₆ H ₄ O ⁻	-5.0			[-7.0]					
CH ₃ O ⁻							[-17.4]		
C ₂ H ₅ O ⁻	-16.7	-14.2	-9.9	-23.6	-18.5	-12.7	[-14.0]	-18.9	-11.7
Na ⁺	-12.5	-7.6	-6.2	-18.0	-13.4	-9.8	-31.0		
K ⁺				-15.3	-11.5	-8.3	-25.8		
Rb ⁺				-13.7	-10.3	-7.2			
Cs ⁺				-20.4	-13.6	-8.4			
NH ₄ ⁺	[-16.1]			[-17.6]			[-29.1]		
CH ₃ NH ₃ ⁺	-11.8						[-23.1]		
(CH ₃) ₃ NH ⁺	-8.8	-4.6	-2.5						
C ₃ H ₁₀ NH ₂ ⁺	[-7.6]								
pyridineH ⁺	[-8.8]			[-13.9]			[-18.3]		
C ₆ H ₅ NH ₃ ⁺	[-9.6]			[-14.4] ^b			[-18.9]		

^a Gas-phase free energy change for the reaction (S)_j—[M[±]+S→(S)]M[±], for a standard-state gas-phase pressure of 1 atm. Experimental free energies are from reference 169. Calculated (B97-1/MG3S) free energies are listed in brackets.

^b An experimental value of -12.0 kcal/mol is reported for this free energy, but was not used in this work (see text).

TABLE 8

Solvation Free Energy of the Proton (kcal/mol) and Normal Hydrogen Electrode Potential (Volts) in Different Solvents

solvent	$\Delta G_s^*(H^+)^{b,b}$	$E_{NHE}^*{}^c$	$G_t^*(H_2O \rightarrow S)^{a,a}$	
			this work	literature
H ₂ O	-265.9 ^d	-4.24	—	—
CH ₃ OH	-263.5	-4.34	2.4	2.1 ^e
CH ₃ CN	-260.2	-4.48	5.7	10.7 ^e
DMSO	-273.3	-3.92	-7.4	-4.6, -7.3 ^{ef}

^a Transfer free energy of the proton from water to the solvent in this row.^b Absolute solvation free energy of the proton obtained using the cluster pair approximation.^c Normal hydrogen electrode potential. These values were obtained using the values for the electron integrated heat capacity and entropy reported by Bartmess,¹¹⁶ who used Fermi-Dirac statistics. Using Boltzmann statistics for the integrated heat capacity and entropy of the electron gives values of -4.28, -4.38, -4.52, and -3.96 V for the normal hydrogen electrode potential in water, methanol, acetonitrile, and DMSO, respectively.^d Reference 53.^e Reference 58.^f Reference 107.

TABLE 9

Absolute Single-Ion Solvation Free Energies ΔG_S^* (in kcal/mol) in Four Solvents^a

ion	H ₂ O ^b	CH ₃ OH ^c	CH ₃ CN ^c	DMSO ^c
H ⁺	-265.9	-263.5	-260.2	-273.3
Na ⁺	-103.2	-101.1	-101.1	-108.9
K ⁺	-86.0	-83.3	-85.5	-92.2
Rb ⁺	-80.6		-81.8	
Cs ⁺	-75.1		-72.4	
Tl ⁺	-87.6	-86.2	-89.8	-95.3
Ag ⁺	-118.7	-117.1	-127.8	-129.0
NH ₄ ⁺	-85.2	-85.6	-89.5	-94.3
CH ₃ NH ₃ ⁺	-76.4	-74.6	-80.4	-83.4
(CH ₃) ₃ NH ⁺	-61.1	-60.5		
C ₅ H ₁₀ NH ₂ ⁺	-64.2	-63.2		
pyridineH ⁺	-61.1	-60.6	-66.4	-67.0
C ₆ H ₅ NH ₃ ⁺	-72.4	-74.0	-76.8	-80.1
Cl ⁻	-74.5	-71.5	-62.4	-62.7
Br ⁻	-68.3	-65.8	-59.3	-59.8
I ⁻	-59.9	-58.2	-53.0	
NO ₂ ⁻				-56.3
NO ₃ ⁻			-53.1	
CH ₃ CO ₂ ⁻	-77.6	-73.8	-59.3	-60.6
CH ₃ CH ₂ CO ₂ ⁻	-76.2	-72.7		
NCCH ₂ CO ₂ ⁻			-49.4	
CF ₃ CO ₂ ⁻	-59.3		-48.5	
C ₆ H ₅ CO ₂ ⁻	-71.2	-68.3	-55.9	-56.9
<i>p</i> -FC ₆ H ₄ CO ₂ ⁻		-64.5		
C ₆ H ₅ O ⁻	-71.9	-70.0	-55.4	-55.1
<i>p</i> -NO ₂ C ₆ H ₄ O ⁻	-57.8	-55.0		
CH ₃ O ⁻	-95.0			-68.9
C ₂ H ₅ O ⁻	-90.7			-64.6
CH ₃ NO ₂ ⁻	-76.5			-61.3
CH ₃ SOCH ₂ ⁻	-67.7			-55.4

dFrom $\Delta G_g^0 = 350.4$ kcal/mol,¹⁶⁵ $\Delta G_S^*(\text{CH}_3\text{NO}_2) = -5.7$ kcal/mol,¹⁰ and $pK_a = 17.2$.¹⁵⁵ eFrom $\Delta G_g^0 = 366.8$ kcal/mol,¹⁶⁵ $\Delta G_{\text{self}}^*(\text{DMSO}) = -7.7$ kcal/mol (from Table 6) and $pK_a = 35$.¹⁵⁵

^a Add 1.9 kcal/mol to obtain ΔG_S^0 .

^b Reference 56.

^c Present work, based on the average $\Delta G_S^{*,\text{con}}$ values of Table 1-3 and 5 and the $\Delta G_S^*(\text{H}^+)$ values in the first row (from Table 8).