Published in final edited form as:

J Med Chem. 2007 December 27; 50(26): 6725–6729. doi:10.1021/jm070895r.

Allosteric Modulation of the Dopamine Receptor by Conformationally Constrained Type VI β -Turn Peptidomimetics of Pro-Leu-Gly-NH₂

Ashish Vartak[†], Kevin Skoblenick[‡], Nancy Thomas[‡], Ram K. Mishra[‡], and Rodney L. Johnson^{*,†}

Department of Medicinal Chemistry, University of Minnesota, 308 Harvard St. SE, Minneapolis, Minnesota 55455-0343 and Department of Psychiatry and Behavioral Neuroscience, McMaster University, 1200 Main St. West, Hamilton, Ontario L8N 2Z5, Canada

Abstract

A peptidomimetic of Pro-Leu-Pro-NH₂, **7**, possessing an indolizidinone type VI β -turn mimic was synthesized via improved high-yielding protocols for the preparation and Cbz-protection of α -allyl proline. Bicyclic peptidomimetic **7** and spirobicyclic peptidomimetic **8** enhanced the binding of [³H] N-propylnorapomorphine to dopamine receptors indicating that a type VI β -turn is a possible bioactive conformation of the homochiral Pro-Leu-Pro-NH₂ and Pro-Pro-Pro-NH₂ analogues of Pro-Leu-Gly-NH₂ at the dopamine receptor allosteric regulatory site.

Introduction

The neuropeptide prolyl-leucyl-glycinamide (**1**, PLG) has been shown to modulate dopaminergic neurotransmission as illustrated by its ability to enhance both amphetamine- and apomorphine-dependent rotational behavior in 6-hydroxydopamine-lesioned rats.¹⁻³ Evidence supports the hypothesis that PLG acts as an allosteric modulator of the dopamine receptor.⁴ PLG and its peptidomimetics have been shown to increase the binding affinity of agonists to the high-affinity state of the dopamine receptor and to shift the ratio of high- and low-affinity states of the dopamine receptor in favor of the G-protein-coupled high-affinity state.^{5,6} The actions of PLG appear to be specific toward dopamine receptors as PLG does not interact with other aminergic receptors such as -adrenergic,^{5,7} GABA-ergic,⁸ or serotonergic receptors.⁹ Furthermore, studies carried out in cell lines transfected with human dopamine receptor subtypes have shown that PLG and a PLG peptidomimetic enhance agonist binding to the D_{2S}, D_{2L}, and D₄ subtypes, whereas the D₁ and D₃ subtypes are unaffected.⁷

* Corresponding author. Phone: +1-612-624-7997. Fax: +1-612-624-0139. E-mail: johns022@umn.edu.

[†]University of Minnesota

[‡]McMaster University

Pro-Leu-Gly-NH₂ Type VI β -Turn Peptidomimetics

Supporting Information Available. Experimental details for the synthesis of compounds **9–14** and **16–22**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Investigations in our lab have centered on the synthesis and evaluation of conformationally constrained analogues of PLG designed to test the hypothesis that the bioactive conformation of PLG at the allosteric site is a type II β -turn conformation.¹⁰ This approach led to the successful development of numerous PLG peptidomimetics;¹¹⁻¹³ one example of which was the highly rigid spiro bicyclic analogue **2**.¹⁴ These peptidomimetics supported the proposition that the bioactive conformation of PLG is a type II β -turn.

In other studies, a set of PLG analogues possessing a more flexible framework was examined in which prolyl residues were substituted for the leucyl and glycyl residues of PLG.^{15, 16} These analogues consisted of Pro-Leu-Pro-NH₂ (**3**), Pro-Leu-D-Pro-NH₂ (**4**), Pro-Pro-Pro-NH₂ (**5**), and Pro-Pro-D-Pro-NH₂ (**6**). Because of the effect that the proline pyrrolidine ring has on the ϕ torsion angle, those PLG analogues that contained a D-prolyl residue in place of the glycyl residue are able to attain a type II β -turn, while those possessing a L-prolyl residue at the C-terminal end are not able to attain such a conformation.

Surprisingly, when these prolyl PLG analogues were evaluated in both *in vitro* and *in vivo* assays for dopamine receptor modulation activity, the homochiral set of analogues were found to be as active or even more active, depending on the assay, than the heterochiral set of analogues.^{15,16} The activity of the homochiral set of prolyl PLG analogues, which could not adopt a type II β -turn, was inconsistent with the hypothesis, strongly supported by the good activity of the highly constrained PLG peptidomimetics like **2**, that the bioactive conformation of PLG is a type II β -turn. To explain this inconsistency, we speculated that the homochiral prolyl PLG analogues are able to adopt a conformation other than a type II β -turn that places the key pharmacophores in the same relative topological space in which these same key pharmacophores exist when PLG assumes a type II β -turn conformation.

In considering potential conformations that the prolyl homochiral peptides could assume, we noted that the presence of a prolyl residue in the *i*+2 position introduces the possibility of a *cis*-amide bond with a resulting type VI β -turn conformation.¹⁷ A comparison of this turn with the type II β -turn reveals several similarities; like the key amide NH₂ group projecting in the same relative space in both turn types (Figure 1). The major topological difference between the two turn conformations is a nearly 109° difference in the projection of the carboxamide carbonyl groups. This suggested to us the possibility that the homochiral peptides **3** and **5** might assume a type VI β -turn when interacting with the PLG allosteric binding site. To test this hypothesis, the design, synthesis and evaluation of analogues of **3** and **5** constrained in a type VI β -turn conformation were undertaken. Although the restriction of the ω dihedral angle to a value of 0° can be achieved in a number of ways, we viewed the indolizidinone approach of Germanas¹⁸ as the most apt, since the 2-carbon bridge between the α -carbons of the *i*+1 and *i*+2 residues of the turn provides minimal steric deviation. Incorporation of this mimic into the prolyl tripeptides **3** and **5** afforded the bicyclic peptidomimetic **7** and the spirobicyclic peptidomimetic **8**, respectively, (Figure 2).

Chemistry

For the synthesis of **7**, we followed Germanas' general route to the indolizidinone framework,¹⁸ but made several improvements and modifications. The key starting material for the indolizidinone synthesis was (*R*)- α -allyl proline (**9**; Scheme 1), of which substantial quantities were required. This amino acid is traditionally prepared through Seebach's 'self-regeneration of chirality' approach where the acid-catalyzed condensation of proline with six equivalents of pivalaldehyde in pentane with continuous removal of water yields a highly moisture-sensitive oxazolidinone.¹⁹ Alkylation of the lithium enolate of this oxazolidinone with allyl bromide yields a stable allylated oxazolidinone, which when hydrolyzed provides **9**. There are several shortcomings to this approach including: the high cost of pivalaldehyde, the large excess of pivalaldehyde required, as well as the duration of the condensation reaction (typically 3–4 days). Modifications made to the condensation and hydrolysis reaction steps of this sequence provided **9** in an overall yields of 70–80 % on a 20 g scale (see Supporting Information).

Cbz-protection of **9** has been a low yielding process, as documented in the past. Repeated additions of Cbz-Cl under Schotten-Baumann conditions over 3 days do produce excellent yields of **9** on a small scale,²⁰ but we found this approach not to be scalable. We previously reported a procedure for the Boc-protection of **9** that involved its solubilization in acetonitrile by forming a salt with tetramethylammonium hydroxide (TMAH).²¹ Application of these same conditions to the Cbz-protection of **9** led to rapid decomposition of Cbz-Cl and a low yield (~20%) of **10**. However, we found that the reaction of the tetramethylammonium salt of **9** with 2 equiv of Cbz-Cl in the presence of one equiv of Et₃N led to the formation of a stable mixed-anhydride, which provided **10** in a 60% yield upon saponification. The reaction could be made quantitative by the addition of an additional equivalent of Cbz-Cl, followed by saponification with aqueous KOH. This quantitative Cbz-protection was reproducible on a scale as large as 15 g and suggests that this protocol may be potentially useful for many such substrates where steric hindrance around the nitrogen prevents efficient Cbz-protection.

Treatment of **10** with diazomethane followed by oxidative cleavage of the terminal olefin of the resulting ester provided aldehyde **11**. The subsequent olefination of **11** required the glycine anion equivalent **12**. Traditionally, this phosphonate is prepared through an Arbuzov reaction between trimethyl phosphite and a Cbz-protected α -bromoglycine ester, followed by replacement of the Cbz group with a Boc group.²² We used an alternative method in which a photochemical electrophilic bromination of Boc-Gly-OEt by NBS²³ was carried out in hexanes instead of the typical hazardous and expensive CCl₄ solvent to generate an α -bromoglycine intermediate that could be converted to **12** through an Arbuzov reaction.

In our hands, attempts at affecting the olefination between the lithium enolate of **12** and aldehyde **11** in THF led to low yields of the desired alkene **13**. Fortunately, a protocol by Zhang, et al.²⁴ that uses a DBU-enolate in CH₂Cl₂ at room-temperature worked well in this case and gave alkene **13** in excellent yield with the *Z*-isomer predominating.

Reduction of the olefin and hydrogenolysis of the Cbz group of **13** was accomplished simultaneously with H₂ over Pd/C. Heating the resulting product for 5–6 h at reflux in MeOH in the presence of sodium metal resulted in complete cyclization to give **14** in a 2:1 ratio of *trans* and *cis* isomers, respectively. After removal of the Boc-group from **14** by TFA, the free amine was reacted with benzaldehyde to form a Schiff base. Initial attempts at alkylating the lithium enolate of the Schiff base with isobutyl bromide in THF at temperatures ranging from –78 °C through to the reflux temperature were met with failure, even though the enolate was found to be quite stable to heat. The use of isobutyl iodide in place of the bromide, solvation of the enolate by HMPA, and heating the reaction to reflux in THF, on the other hand, resulted

in a complete reaction within about 20 min to give the alkylated product **15** as a single diastereoisomer. Breakdown of the Schiff base with aqueous TFA provided bicyclic amine **16**.

We also explored the use of bis-Boc protection in place of the Schiff base for the alkylation reaction (Scheme 2). When the 2:1 *trans*:*cis* diastereoisomeric mixture of lactam **14** was treated with 10 equiv of Boc₂O and DMAP in CH₃CN, a diastereoisomeric mixture of the bis-Boc lactam **17** was obtained in an 88% yield. Surprisingly, the coupling constants of the lactam α -proton in this diastereoisomeric mixture (8.1 and 10.2 Hz for the major isomer and 9.9 and 3.6 Hz for the minor isomer) indicated that the *trans*:*cis* ratio was now 4:9. The precise mechanism for this inversion is not known, although it is known that epimerization does take place when indolizidinones of this relative stereochemistry are exposed to alkoxide. On the basis of the observation that DMAP causes a vigorous bubbling of CO₂ when added to a solution of Boc₂O, we believe that the transient *tert*-butoxide liberated in this process may be responsible for the deprotonation of the α -proton. Reprotonation from the less sterically hindered *si* face leads to the preferential formation of a *cis*-lactam.

Deprotonation of diastereoisomeric mixture **17** by LDA followed by addition of isobutyl iodide led to the formation of lactam **18** instead of the expected alkylated product; a net N→C shift. This product formed even without isobutyl iodide being present. An NOE between the carbamate NH and the methyl ester hydrogens served to support the expected *R* configuration about the newly formed quaternary carbon. In contrast, deprotonation of **17** with KHMDS in the presence of 2 equiv of allyl bromide afforded a single diastereoisomer of the α -allylated bicyclic lactam **19**. An NOE between the *t*-butyl hydrogens of the Boc group and the hydrogens of the methyl ester suggested that the allylation had occurred *trans* to the methyl ester function.

The coupling reaction of **16** with Boc-proline to give **21** was sluggish with reagents that produce relatively stable esters (e.g. HOBt ester of Boc-proline via treatment with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC)), as well as with the mixed anhydride reagent isobutyl chloroformate in either the presence or absence of DMAP. The main byproduct in these reactions was the diketopiperazine **20** (Scheme 3). We observed an identical result during the synthesis of spirobicyclic lactam **8**.²⁵ The coupling reagents bis(2-oxo-3-oxazolidinyl)-phosphinic chloride (BOP-Cl) and 1,3-dicyclohexylcarbodiimide (DCC) were comparable to EDC. On the other hand, the coupling behavior of 2-chloro-1-methylpyridinium iodide (Mukaiyama's reagent, CMPI) was radically different depending on whether DMAP was present. With CMPI alone, the yield of **21** was only ~30% after 12 h. When DMAP was combined with CMPI, an exothermic reaction was observed that yielded **21** in a 60–75% yield within 1–2 h, averting formation of **20**.

The direct amidation of **21** to give the primary carboxamide derivative **23** was not attainable by the following conditions: heating a solution of **21** in methanolic ammonia at 100 °C in a sealed stainless steel vessel, heating a solution of **21** in methanolic ammonia in the presence of either DMAP, NaCN or LiI, and Wienreb's amidation with NH₃ in the presence of an equivalent of trimethylalane. However, **21** could be hydrolyzed quite easily to the acid **22** with LiOH in a mixture of THF and HMPA at reflux. EDC-mediated coupling of the free carboxylic acid **22** with ammonia then afforded carboxamide **23**. Subsequent removal of the Boc-protecting group from **23** gave the target molecule **7** as its HCl salt.

We also attempted to incorporate the primary carboxamide function prior to the coupling stage. While we could form carboxamide **24** upon treating **16** with methanolic ammonia, diketopiperazine **20** also formed. Attempts to couple **24** to Boc-Pro-OH with CMPI failed. Instead diketopiperazine **20** and Boc-prolinamide were isolated suggesting that the primary

amide function of **24** initially gets acylated, thereby forming **25**. Nucleophilic attack of the free amine on the imide carbonyl then ejects Boc-prolinamide and forms diketopiperazine **20**.

In CDCl₃, the primary carboxamide hydrogens of **23** appeared in a typical hydrogen-bonded pattern with chemical shifts of 5.7 and 8.3 ppm. Further support for a hydrogen bond was obtained by the addition of CD₃OD to the NMR sample of **23**; that led to complete exchange of the non-H-bonded, upfield hydrogen, while the H-bonded downfield hydrogen required 30 min for complete exchange. Similar behavior was seen with a solution of **7** in CD₃OD, in which case the downfield carboxamide hydrogen resonance at 7.87 ppm diminishes slower than the upfield carboxamide hydrogen resonance at 7.42 ppm. Such behavior was observed previously with **8** and its Boc-protected precursor.²⁵ The results for **7** and **8** are consistent with a hydrogen bond between the carboxamide NH and the prolyl carbonyl group that one would expect in a type VI β -turn conformation.

Pharmacology

Bicyclic peptidomimetic **7** and spirobicyclic peptidomimetic **8**, the synthesis of which was carried out as reported by us previously,²⁵ were both examined for their ability to potentiate the binding of tritiated N-propylnorapomorphine ([³H]NPA) to bovine striatal membranes.^{5, 26} The results are shown in Figure 3. Compound **7** exhibited a bell-shaped dose response curve that is typical of the allosteric modulation of the dopamine receptor by PLG and its analogues. Statistically significant increases were observed at the following concentrations of **7**: 10 nM, 100 nM, and 1 μ M. The maximum effect for **7** was $19.5 \pm 3.75\%$ at 100 nM. For spirobicyclic **8**, a bell-shaped dose response curve was also seen, but only at a concentration of 10 nM was a significant increase in [³H]NPA binding observed ($11.4 \pm 3.3\%$). At a concentration of 100 nM, PLG peptidomimetic **2** and the triproline analogue **5** enhance NPA binding to dopamine receptors by $38.2 \pm 11.9\%$ and $34.0 \pm 13.1\%$, respectively.

Discussion

The greater activity of bicyclic peptidomimetic **7** compared to the spirobicyclic peptidomimetic **8** may be because the greater rigidity of the spirobicyclic scaffold prevents the optimal interaction of **8** with the allosteric binding site. It also may be that the isobutyl side chain of **7** interacts more effectively with the hydrophobic pocket in the allosteric binding site than does the propylene-bridging unit of **8**. Previous work has shown that PLG analogues and peptidomimetics with a hydrophobic moiety placed in this topological region possess enhanced dopamine receptor modulating activity.²⁶⁻²⁸

The activity of **7** and **8** supports the hypothesis that the homochiral prolyl peptides **3** and **5** may assume a type VI β -turn as their bioactive conformation and that such a conformation is able to initiate a modulatory response because it can present the necessary topological arrangement of important pharmacophore moieties in a manner similar to that of PLG peptidomimetics that assume a type II β -turn bioactive conformation. In this regard, the data indicate that the amide NH₂ group may be a more important pharmacophore group than the carboxamide carbonyl group in activating the D₂ receptor allosteric modulatory site, as the major difference between a type II β -turn conformation and a type VI β -turn conformation is the nearly 109° difference in orientation of the carboxamide carbonyl groups. The binding conformation of the homochiral prolyl PLG analogues **3** and **5** perhaps possess a greater capability of projecting key pharmacophores in the correct topological space when compared to the heterochiral prolyl analogues **4** and **6** that are relatively unconstrained when compared to the active highly rigid type-II β -turn PLG peptidomimetics, such as **2**. Clearly, in terms of allosteric modulation of the dopamine receptor, more than one secondary structure is able to project pharmacophores in an orientation that satisfies the pharmacophore requirements for bioactivity.

Experimental Section

(6S,8aR,2'S)-6-N-(tert-Butoxycarbonyl-2'-pyrrolidinylcarbonyl)amino-6-(2-methylpropyl)-8a-carboxamido-indolizidin-5-one (23)

To the crude carboxylic acid **22** (480 mg, 1.38 mmol) were added CH₂Cl₂ (5 mL), DMF (5 mL) and HATU (279 mg, 1.05 equiv). The mixture was stirred for 15 min. The flask was evacuated (vacuum needle line) for 1 min and excess NH₃ was introduced through a balloon. This resulted in an exothermic reaction and the precipitation of a canary-yellow solid. The mixture was stirred for 24 h, diluted with CH₂Cl₂ (50 mL), and filtered. The filtrate was concentrated under reduced pressure and the resulting residue was mixed with xylenes (50 mL). The solvent was removed and the crude yellow oil that was obtained was dissolved in EtOAc. This solution was loaded onto a plug of silica gel (30 g) with the aid of EtOAc. The plug was washed with ether (300 mL), which removed most of the HMPA, and then eluted with EtOAc (100 mL), EtOAc/MeOH (20:1, 100 mL), EtOAc/MeOH (15:1, 200 mL) and EtOAc/MeOH (10:1, 100 mL). The product was found in the last 80 mL of the eluent. The relevant fractions were concentrated under reduced pressure to yield a white foam that was triturated with Et₂O and then dissolved in EtOAc/CH₂Cl₂/Et₂O (1:1:40). The solution was allowed to stand uncovered to give a white powder that was filtered to give **23** (430 mg, 87 %). [α]_D –43.6 (*c* 1.0, CDCl₃); *R*_f = 0.4 (EtOAc/MeOH, 9:1); ¹H NMR (CDCl₃, 300 MHz, rotamers present about the N–CO₂*t*-Bu bond) δ 8.26 (s, 1H, CONH₂), 7.68 and 6.43 (2s, 1H, CONH), 5.68 (s, 1H, CONH₂); 4.26 (br s, Pro α -CH), 3.73–3.29 (m, 4H, δ -CH₂ and 3-CH₂), 2.56 (dd, *J*₁ = 11.7 Hz, *J*₂ = 6.0 Hz, 1H, β -CH₂), 2.35–1.60 (m, 14H, Pro β -CH₂, 1-CH₂, Pro γ -CH₂, 2-CH₂, 7-CH₂, 8-CH₂, CH₂CH(CH₃)₂), 1.45 (s, 9H, *t*-Bu), 0.97 (dd, *J*₁ = 12 Hz, *J*₂ = 7.2 Hz, 6H, CH(CH₃)₂); ¹³C NMR (CDCl₃, 75 MHz, rotamers present about the N–CO₂*t*-Bu bond) δ 176.2 (CONH₂), 171.2 (CONH), 170.0 (5-C), 80.8 (C(CH₃)₃), 71.0 (8a-C), 60.7, 59.8 (Pro α -C), 58.6 (6-C), 47.4 (Pro δ -C or 3-C), 45.9 (CH₂CH(CH₃)₂), 45.0 (Pro δ -C or 3-C), 38.4 (1-C), 31.2 and 30.0 (Pro β -C), 30.3 (7-C), 29.3 (8-C), 28.1 (*t*-Bu), 27.5 (Pro γ -C), 24.9 (CH₂CH(CH₃)₂), 24.3 (CH₂CH(CH₃)₂), 21.4 (8-C). ESI-HRMS *m/z* 472.2752 (M+Na)⁺, C₂₃H₃₈N₄O₅ + Na⁺ requires 472.2740; Anal. (C₂₃H₃₈N₄O₅) C, H, N.

(6S,8aR)-6-N-(2'-Pyrrolidinylcarbonyl)amino-6-(2-methylpropyl)-8a-carboxamido-indolizidine-5-one Hydrochloride (7•HCl)

To **23** (100 mg, 0.22 mmol) was added 4N HCl in dioxane (5 mL) under a stream of argon. A white solid precipitated from the solution within 15–20 seconds. The suspension was stirred for an extra 15 min and a vacuum line was inserted in the septum to remove excess HCl. The residue was then evaporated under reduced pressure to afford a white gum, which was dissolved in CH₂Cl₂ and the solvent then removed (5 $\times$). The resulting white gum was dissolved in CH₂Cl₂/MeOH (10:1) and this solution was poured into ether, upon which a white precipitate formed. The solvents were decanted and the residue was dried for 2 days under high vacuum to yield 74 mg of **1** as a hygroscopic white solid. *R*_f = 0.7 (EtOH/NH₄OH, 4:1); [α]_D –24.8 (*c* 0.5, CD₃OD); ¹H NMR (300 MHz, CD₃OD, acquired within 5 min of mixing) δ 7.87 and 7.42 (2s, 2H, CONH₂, these resonances disappear within 30 min of mixing), 4.31 (app dd, *J*₁ = 6.2 Hz, *J*₂ = 6.6 Hz, 1H, Pro α -CH), 3.67–3.29 (m, 4H, Pro δ -CH₂ and 3-CH₂), 2.67 (dd, *J*₁ = 5.7 Hz, *J*₂ = 12.0 Hz, 1H, Pro β -CH₂), 2.46–1.66 (m, 14H, Pro β -CH₂, Pro γ -CH₂, 2-CH₂, 1-CH₂, CH₂CH(CH₃)₂, 7-CH₂, 8-CH₂); 0.98 (dd, *J*₁ = 11.4 Hz, *J*₂ = 6.3 Hz, CH(CH₃)₂). ¹³C NMR (CDCl₃, 75 MHz) δ 179.5 (CONH₂), 170.1 (CONH), 173.6 (5-C), 70.1 (8a-C), 61.2 (Pro α -C), 58.6 (6-C), 48.0 (Pro δ -C or 3-C), 44.0 (CH₂CH(CH₃)₂), 43.8 (Pro δ -C or 3-C), 38.1 (1-C), 32.1 (Pro β -C), 31.4 (7-C), 29.3 (8-C), 24.9 (Pro γ -C), 24.5 (CH(CH₃)₂), 24.3 (CH(CH₃)₂), 21.4 (8-C). ESI-HRMS *m/z* 351.2358 (M)⁺, (C₁₈H₃₁N₄O₃)⁺ requires 351.2364. Anal. (C₁₈H₃₁ClN₄O₃•1/3 H₂O) C, H, N.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

This work was supported in part by NIH grant NS20036 to R.L.J. We thank Dr. David Rusterholz for helpful discussions on the proline-pivaldehyde condensation reaction.

References

1. Smith JR, Morgan M. The Effects of PLG on Drug Induced Rotation in Rats. *Gen. Pharmacol* 1982;13:203–207. [PubMed: 6124481]
2. Ott MC, Mishra RK, Johnson RL. Modulation of Dopaminergic Neurotransmission in the 6-Hydroxydopamine Lesioned Rotational Model by Peptidomimetic Analogues of l-Prolyl-l-leucylglycinamide. *Brain. Res* 1996;737:287–291. [PubMed: 8930377]
3. Mishra RK, Marcotte ER, Chugh A, Barlas C, Whan D, Johnson RL. Modulation of Dopamine Receptor Agonist-Induced Rotational Behaviour in 6-OHDA-Lesioned Rats by a Peptidomimetic Analogue of Pro-Leu-Gly-NH₂ (PLG). *Peptides* 1997;18:1209–1217. [PubMed: 9396063]
4. Fisher A, Mann A, Verma V, Thomas N, Mishra RK, Johnson RL. Design and Synthesis of Photoaffinity-Labeling Ligands of the l-Prolyl-l-leucylglycinamide Binding Site Involved in the Allosteric Modulation of the Dopamine Receptor. *J. Med. Chem* 2006;49:307–317. [PubMed: 16392815]
5. Srivastava LK, Bajwa SB, Johnson RL, Mishra RK. Interaction of l-Prolyl-l-Leucyl Glycinamide with Dopamine D₂ Receptor: Evidence for Modulation of Agonist Affinity States in Bovine Striatal Membranes. *J. Neurochem* 1988;50:960–967. [PubMed: 2892892]
6. Mishra RK, Srivastava LK, Johnson RL. Modulation of High-Affinity CNS Dopamine D₂ Receptor by l-Pro-l-Leu-Glycinamide (PLG) Analogue 3(R)-(N-l-Prolylamino)-2-oxo-1-pyrrolidineacetamide. *Prog. Neuro-Psychopharmacol. & Biol. Psychiat* 1990;14:821–827.
7. Verma V, Mann A, Costain W, Pontoriero G, Castellano JM, Skoblenick K, Gupta SK, Pristupa Z, Niznik HB, Johnson RL, Nair VD, Mishra RK. Modulation of Agonist Binding to Human Dopamine Receptor Subtypes by L-Prolyl-L-Leucyl-Glycinamide and a Peptidomimetic Analogue. *J. Pharmacol. Expt. Ther* 2005;315:1228–1236.
8. Miller L, Kastin AJ. MIF-1 and Tyr-MIF-1 Do Not Alter GABA Binding on the GABA_A Receptor. *Brain Res. Bull* 1990;25:917–919. [PubMed: 1981176]
9. Gulati A, Bhargava HN. Effect of Melanotropin Release Inhibiting Factor on Changes by Haloperidol and Centbutindole in Cerebral Cortical 5-Hydroxytryptamine Receptors. *Pharmacology* 1990;41:98–106. [PubMed: 1979878]
10. Yu K-L, Rajakumar G, Srivastava LK, Mishra RK, Johnson RL. Dopamine Receptor Modulation by Conformationally Constrained Analogues of Pro-Leu-Gly-NH₂. *J. Med. Chem* 1988;31:1430–1436. [PubMed: 3385734]
11. Sreenivasan U, Mishra RK, Johnson RL. Synthesis and Dopamine Receptor Modulating Activity of Lactam Conformationally Constrained Analogues of Pro-Leu-Gly-NH₂. *J. Med. Chem* 1993;36:256–263. [PubMed: 8093736]
12. Subasinghe NL, Bontems RJ, McIntee E, Mishra RK, Johnson RL. Bicyclic Thiazolidine Lactam Peptidomimetics of the Dopamine Receptor Modulating Peptide Pro-Leu-Gly-NH₂. *J. Med. Chem* 1993;36:2356–2361. [PubMed: 8103113]
13. Genin MJ, Mishra RK, Johnson RL. Dopamine Receptor Modulation by a Highly Rigid Spiro Bicyclic Peptidomimetic of Pro-Leu-Gly-NH₂. *J. Med. Chem* 1993;36:3481–3483. [PubMed: 7901418]
14. Khalil EM, Ojala WH, Pradhan A, Nair VD, Gleason WB, Mishra RK, Johnson RL. Design, Synthesis, and Dopamine Receptor Modulating Activity of Spiro Bicyclic Peptidomimetics of l-Prolyl-l-leucylglycinamide. *J. Med. Chem* 1999;42:628–637. [PubMed: 10052970]
15. Johnson RL, Rajakumar G, Mishra RK. Dopamine Receptor Modulation by Pro-Leu-Gly-NH₂ Analogues Possessing Cyclic Amino Acid Residues at the C-Terminal Position. *J. Med. Chem* 1986;29:2100–2104. [PubMed: 2876103]

16. Baures PW, Pradhan A, Ojala WH, Gleason WB, Mishra RK, Johnson RL. Synthesis and Dopamine Receptor Modulating Activity of Unsubstituted and Substituted Triproline Analogues of l-Prolyl-l-leucyl-glycinamide. *Bioorg. Med. Chem. Lett* 1999;9:2349–2352. [PubMed: 10476867]
17. a Yao J, Victoria A, Feher B, Fabiola E, Reymond MT, Wright PE, Dyson JH. Stabilization of a Type-VI Turn in a Family of Linear Peptides in Water Solution. *J. Mol. Biol* 1994;243:736–753. [PubMed: 7966293] Richardson, JS.; Richardson, DC. Prediction of Protein Structure and the Principles of Protein Conformation. Fasman, GD., editor. Plenum Press; New York: 1989. p. 48-53.
18. Kim K, Dumas J, Germanas JP. Novel Bicyclic Lactams as XaaPro Type VI β -Turn Mimics: Design, Synthesis and Evaluation. *J. Org. Chem* 1996;61:3138–3144. [PubMed: 11667176]
19. Seebach D, Boes M, Naef R, Schweizer WB. Alkylation of Amino Acids Without Loss of the Optical Activity: Preparation of α -Substituted Proline Derivatives. A Case of Self-reproduction of Chirality. *J. Am. Chem. Soc* 1983;105:5390–5398.
20. Gramberg D, Weber C, Beeli R, Inglis J, Bruns C, Robinson JA. Synthesis of a Type-VI β -turn Peptide Mimetic and Its Incorporation into a High-affinity Somatostatin Receptor Ligand. *Helv. Chim. Acta* 1995;78:1588–1606.
21. Khalil EM, Subasinghe NL, Johnson RL. An Efficient and High Yield Method for the N-tert-Butoxycarbonyl Protection of Sterically Hindered Amino Acids. *Tetrahedron Lett* 1996;37:3441–3444.
22. Ku B, Oh DY. Facile Synthesis of α -Phosphorylated α -Amino Acids. *Tetrahedron Lett* 1998;29:4465–4466.
23. Kober R, Steglich W. Reaction of Acylaminobromomalonates and Acylaminobromoacetates with Trialkyl Phosphates – A Simple Synthesis of Ethyl 2-Amino-2-(diethoxyphosphoryl)acetate. *Lieb. Ann. Chim* 1983;4:599–609.
24. Zhang J, Xiong C, Ying J, Wang W, Hruby VJ. Stereoselective Synthesis of Novel Dipeptide β -Turn Mimetics Targeting Melanocortin Peptide Receptors. *Org. Lett* 2003;5:3115–3118. [PubMed: 12916995]
25. Vartak AP, Johnson RL. Concerted Synthesis of a Spirobicyclic Type-VI β -Turn Mimic of Pro-Pro-Pro-NH₂. *Org. Lett* 2006;8:983–986. [PubMed: 16494490]
26. Dolbeare K, Pontoriero GF, Gupta SK, Mishra RK, Johnson RL. Synthesis and Dopamine Receptor Modulating Activity of 3-Substituted γ -Lactam Peptidomimetics of l-Prolyl-l-leucyl-glycinamide. *J. Med. Chem* 2003;46:727–733. [PubMed: 12593653]
27. Johnson RL, Bontems RJ, Yang KE, Mishra RK. Synthesis and Biological Evaluation of Analogues of Pro-Leu-Gly-NH₂ Modified at the Leucyl Residue. *J. Med. Chem* 1990;33:1828–1832. [PubMed: 1971310]
28. Khalil EM, Pradhan A, Ojala WH, Gleason WB, Mishra RK, Johnson RL. Synthesis and Dopamine Receptor Modulating Activity of Substituted Bicyclic Thiazolidine Lactam Peptidomimetics of l-Prolyl-l-leucyl-glycinamide. *J. Med. Chem* 1999;42:2977–2987. [PubMed: 10425107]

Figure 1. Pro-Leu-Gly-NH₂ (**1**, PLG) in a type-II β -turn conformation (grey) overlaid with Pro-Leu-Pro-NH₂ (**3**) in a type-VI β -turn conformation.

Figure 2. Type-VI β -turn mimics of Pro-Leu-Pro-NH₂ (3) and Pro-Pro-Pro-NH₂ (5), compounds 7 and 8, respectively.

Scheme 1.

Scheme 2.

Scheme 3.

Figure 3.

Enhancement of $[^3\text{H}]$ NPA binding to bovine striatal membranes by **7** and **8**. Data represent the percent increase in specific $[^3\text{H}]$ NPA binding over the control value when the indicated concentration of compound was added directly to the assay buffer. Results are the mean $\pm$ SEM of 3–4 separate experiments carried out in triplicate. The data were analyzed by a one way analysis of variance followed by Dunnett's post-hoc test. *Significantly different ($p < 0.01$) from control value; **significantly different ($p < 0.001$) from control value.