Published in final edited form as:

Org Lett. 2007 November 8; 9(23): 4773–4776. doi:10.1021/ol7020675.

Latrunculin with a Highly Oxidized Thiazolidinone Ring: Structure Assignment and Actin Docking

Safwat A. Ahmed^a, Srinivas Odde^b, Pankaj R. Daga^b, John J. Bowling^{c,d}, Mostafa K. Mesbah^a, Diaa T. Youssef^a, Sherief I. Khalifa^{a,*}, Robert J. Doerksen^{b,c,*}, and Mark T. Hamann^{c,d,*}

Department of Pharmacognosy, Faculty of Pharmacy, Suez Canal University, Ismailia, Egypt and Department of Medicinal Chemistry, National Center for Natural Products Research and Department of Pharmacognosy, School of Pharmacy, The University of Mississippi, MS 38677, USA

Abstract

A new latrunculin, oxalatrunculin B (3), was isolated from Red Sea sponge *Negombata corticata*. Extensive spectroscopic analysis revealed an unprecedented heterocycle in which the rare thiazolidinone ring found in latrunculins was oxidized with three additional oxygens. An actin polymerization inhibition assay agreed with MM-PBSA free energy calculations that 3 binds more weakly than latrunculin B to actin. Significant antifungal and anticancer activity of 3 was found, suggesting an alternate target in addition to actin for latrunculin bioactivity.

Latrunculins A (1) and B (2) (Fig. 1) were first isolated from *Negombata magnifica* (Keller) (formerly *Latrunculia magnifica*), discovered from the Red Sea.¹ Among the characteristic features of these natural products is the presence of a macrocyclic lactone ring of 14 or 16 carbon atoms and a 2-thiazolidinone moiety (Figure 1).^{2,3} *In vitro* experiments revealed that the latrunculins disrupt actin polymerization,⁴ a possible target for cancer.⁵

A recent study has also reported antimigratory and antiangiogenic activity of latrunculins, adding to their possible utility in the control of cancer.⁶ In addition to significant ichthyotoxic and cytotoxic properties, the latrunculins are also antiviral against herpes simplex type 1 virus (HSV-1).⁷

Several new latrunculins including natural, synthetic or semisynthetic analogs have been reported recently.^{6,8} We report here the structure assignment and bioactivity of a unique new

jkhalifa_99@yahoo.com, rjd@olemiss.edu, mthamann@olemiss.edu.

^aDepartment of Pharmacognosy, Suez Canal University.

^bDepartment of Medicinal Chemistry, The University of Mississippi

^cNational Center for Natural Products Research, The University of Mississippi.

^dDepartment of Pharmacognosy, The University of Mississippi.

latrunculin, oxalatrunculin B (**3**) which possesses a novel and highly oxidized 2-thiazolidinone ring. Molecular modeling of **1–3** in the actin monomer active site was utilized as a predictor of biological activity. The Red Sea sponge *N. corticata* was extracted with MeOH/CH₂Cl₂. Silica column chromatography of the extract afforded several fractions, one of which yielded metabolite **3**. HRTOF-MS of **3** displayed a molecular ion peak at $m/z = 442.1551$ [M-H][−] supporting the molecular formula C₂₀H₂₉SO₈N. Comparison of **3** with **2** revealed an increase of 48 mass units. Addition of three oxygen atoms satisfied the formula difference giving a calculated mass of $m/z = 442.1536$ [M-H][−]. As further evidence for this, combustion analysis on **3** showed 53.6% carbon, 6.51% hydrogen, 3.16% nitrogen, 28.6% oxygen and 6.34% sulfur. ¹³C NMR indicated that the carbon skeleton of **3** consisted of 2 methyl carbons, 7 methylenes, 7 methines and 4 quaternary carbon atoms, which is identical to the carbon multiplicities of **2**. The ¹³C NMR chemical shifts for the 14-membered and tetrahydropyran rings were in close agreement with those of **2** revealing that additional oxygenation must occur at the heteroatoms. Significant differences were found in the chemical shifts of the thiazolidinone ring. The C-17 resonance shifted downfield from δ28.5 to δ49.9 indicating the introduction of a neighboring oxygen atom. C-16 shifted slightly upfield so the only possible position for oxidation was at the sulfur atom. C-18 also moved upfield from δ175.3 to δ157.3. The chemical shift of the carbonyl carbon of a 5-membered lactam ring will move upfield around 15ppm when the lactam nitrogen is hydroxylated.^{9,10} It has also been shown that oxygenation of the sulfur atom adjacent to the carbonyl of a 5-membered thiazolidinone will move chemical shifts upfield by ~10 ppm.¹¹ This suggested that two of the three new oxygen atoms are at the S and the last is assigned as the hydroxyl group at N. Further evidence for this was the disappearance of the H-N at δ6.1. In addition the amide IR resonance in **2** is absent in **3** and a new resonance at 1756 cm^{−1} appeared, indicating the presence of a CONOH system.¹¹ The OH band at 3448 cm^{−1} is significantly increased in intensity for **3** indicating the presence of two OH groups in **3** versus one OH group in **2**. The IR showed a strong band at 1051 cm^{−1} indicating the presence of an SO₂ moiety.¹² Inclusion of three oxygen atoms on the thiazolidinone moiety increases its electronegativity and explains the downfield shift of C-15 from δ97.2 to δ104.3. HMBC showed important J^3 correlations of H-16 with C-18, C-14 and C-15; and of H-17 with C-18 and C-15.

We used molecular modeling to calculate the binding affinity of **3** in G-actin. X-ray crystal structures of **1** bound to G-actin, such as 1RDW.pdb,¹³ (Fig. 2) showed **1**'s binding site, but there is no X-ray structure available for other latrunculins bound to actin.

Latrunculins **1** and **2** each docked into 1RDW with a highly similar pose to that of the X-ray structure of **1** in actin, as also found in the docking studies of Fürstner et al.⁸ Oxalatrunculin **3** docked in a similar pose to that of **1** and **2**, but despite the two extra polar functional groups in **3**, the docking scores (Table 2) predicted poor binding of **3** with actin, with lower GOLD and Chem scores and poorer estimated ΔG. Since in docking the receptor is fixed, we proceeded to carry out molecular dynamics (MD) simulations and binding free energy calculations using the MM-PBSA/GBSA method implemented in AMBER 8.0,¹⁴ in water with the entire protein flexible. The major favorable contributions to binding were van der Waals and electrostatic terms (Table 2). In the MD simulations, an H-bond to Glu207 was similar for **1–3** and was persistently maintained. In the different heterocycle of **3**, partial displacement of the macrocyclic ring of **3** away from the active site and smaller macrocycle ring size compared to **1** (Fig. 2) resulted in **3**'s decreased electrostatic and van der Waals energies. **3** showed water-mediated H-bonding of NOH with Asp157, whereas for **1** the NH...Asp157 interaction does not need water-assistance (1RDW). Our theoretically most reliable result, from MM-PBSA,¹⁴ reveals a poorer binding free energy for **3** compared to **1** or **2**, by 20.6 or 6.5 kcal/mol, respectively.

In an assay designed to measure inhibition of actin polymerization, the EC₅₀ of **2** was determined to be 2.56 μ M, which is consistent with previous empirical evidence reporting comparisons with **1**.⁴ A relative comparison of the percent inhibition of actin polymerization by **2** and **3** showed that **3** had significantly less potency. This matched well with the molecular modeling results, in which **3** binds much more weakly than **2** (Fig. 3).

Antifungal activity of **3** against *Saccharomyces cerevisiae* NRRL Y-2034 showed an MIC of 56 μ M vs. 127 μ M for **2**. Cytotoxicity of **3** was evaluated against several cell lines including HepG2, HCT-116 and 1301. **3** showed nonspecific cytotoxicity against solid tumor cells and hematopoietic cancerous cells. **3** was also found to be more potent against hepatocellular carcinoma than **2** or latrunculin T, with MIC of 16.34 μ M for **3**, 19.27 μ M for **2**, and 34.72 μ M for latrunculin T. Considering the evidence presented above that **3** binds more weakly than **2** to actin, the stronger cytotoxic activity of **3** compared to **2** suggests that there is an alternate target to actin for latrunculin bioactivity or the compound is reduced in situ and acts as a prodrug.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors are grateful to the U.S.-Egypt Science and Technology joint fund, project number BIO8-002-013, National Institute of Health (NIAID) 1RO1AI36596, NIH National Center for Research Resources C06 RR-14503-01, and National Science Foundation EPS-0556308 for funding; to the Egyptian Environmental Affairs Agency for facilitating sponge sample collection; to R. van Soest, University of Amsterdam for taxonomic identification of the sponge.

References

- Kelly, M.; Sammaai, T. Family Podospongiidae De Laubenfels. In: Hooper, NA.; Van Soest, RWM., editors. *Systema Porifera: A Guide to Classification of Sponges*. 1. Kluwer Academic/Plenum; New York: 2002. p. 698-699.
- Kashman Y, Groweiss A, Shmueli U. *Tetrahedron Lett* 1980;21:3629–3632.
- Groweiss A, Shmueli U, Kashman Y. *J Org Chem* 1983;48:3512–3516.
- Spector I, Shochet NR, Kashman Y, Groweiss A. *Science* 1983;219:493–495. [PubMed: 6681676]
- Yamazaki D, Kurisu S, Takenawa T. *Cancer Sci* 2005;96:379–86. [PubMed: 16053508]
- El Sayed KA, Youssef DT, Marchetti DJ. *J Nat Prod* 2006;69:219–223. [PubMed: 16499319]
- Hoye TR, Ayyad SN, Eklov BM, Hashish NE, Shier WT, ElSayed KA, Hamann MT. *J Am Chem Soc* 2002;124:7405. [PubMed: 12071749]
- Fürstner A, Kirk D, Fenster MDB, Aïssa C, De Souza D, Nevado C, Tuttle T, Thiel W, Müller O. *Chem Eur J* 2007;13:135.
- Crews, P.; Rodriguez, J.; Jaspars, M. *Organic Structure Analysis*. Houk, KN.; Loudon, GM., editors. Oxford University Press; The Netherlands: 1998. p. 94
- Motorina I, Fowler F, Grierson D. *J Org Chem* 1997;62:2098. [PubMed: 11671514]
- Hanefeld W, Jalili MA. *Liebigs Ann Chem* 1986:1787–1795.
- Pretsch, E.; Bühlmann, P.; Affolter, C. *Structure Determination of Organic Compounds*. Springer; 2000. p. 245-312. Chap 6
- Reutzel R, Yoshioka C, Govindasamy L, Yarmola EG, Agbandje-McKenna M, Bubb MR, McKenna RJ. *Struct Biol* 2004;146:291–301.
- Massova I, Kollman PA. *Perspect Drug Discov Des* 2000;18:113–135. Entropy effects were neglected. See Supporting Information for details

Figure 1.
Latrunculins A (1) and B (2) and oxalatrunculin B (3).

Figure 2. Overlay of **3** (green C, from an MD snapshot), displaced away from the active site compared to **1** (purple C, from 1RDW), in G-actin (cyan C). H-bonding interactions of **3** are shown (yellow dashed lines), including to a crystallographic water (red O).

Figure 3. Relative inhibition (DMSO control) of actin polymerization by latrunculin B (**2**) at the predetermined EC_{50} (μM) compared to various concentrations (μM) of the oxalatrunculin B (**3**) analog. Some inhibition is indicated at the highest concentration tested (15 μM).

Table 1

^{13}C and ^1H COSY and HMBC[§] NMR spectral data[¶]

No.	δ_{C} (2)	δ_{C} (3)	δ_{H} (3)	J (Hz)	HMBC
1	165.6	166.3			
2	118.0	117.9	5.60 (s)		
3	154.7	157.9			
4	35.8	35.9	1.97 (m)		
5	26.9	26.4	3.05 (m)		
6	127.6	128.1	2.15 (m)		
7	135.9	135.2	2.70 (m)		
8	28.9	28.7	5.40 (ddd)	11.2, 11.2, 3.0	
9	31.2	31.0	5.00 (dd)	10.8, 10.8	
10	31.2	34.6	2.70 (m)		
11	62.6	65.2	1.10 (m)		
12	35.4	30.7	1.70 (m)		13
13	68.7	67.5	1.70 (m)		11
14	31.8	34.2	4.50 (m)		
15	97.7	104.3	1.40 (m)		
16	61.8	57.8	1.60 (m)		
17	28.7	49.6	5.20 (m)		11, 15
18	175.3	157.3	2.15 (m)	15.2	15, 16
19	24.1	24.8	2.44 (d)		
20	22.3	21.3	4.00 (m)		14, 15, 18, 17
			3.00 (m)	14.4	15, 16, 18
			3.20 (d)		
			1.96 (s)		
			0.90(d)	6.0	

[¶] 100 MHz;

[#] 400 MHz;

[¶] in MeOD;

[§] H to C.

Table 2
Docking scores and binding energies (ΔE) and free energies (ΔG) for latrunculins **1–3** (energies in kcal/mol).

Molecular Docking			Molecular Dynamics Simulations				
GoI ^a	Chem ^a	ΔG^b	ΔE_{ele}	ΔE_{vw}	ΔG_A	ΔG^c	ΔG^d
1	90.7	36.4	-54.9	-52.9	-6.68	-43.2	-35.5
2	70.0	34.5	-40.6	-40.6	-6.19	-26.6	-21.4
3	58.6	20.3	-36.8	-39.4	-5.92	-22.3	-14.9

^a Docking scores;

^b Estimate from ChemScore;

^c MM-GBSA;

^d MM-PBSA.