Published in final edited form as:

Org Lett. 2006 September 28; 8(20): 4593–4595. doi:10.1021/ol061862t.

Synthesis of a ketone analog of biotin *via* the intramolecular Pauson-Khand reaction

Eric McNeill, Irwin Chen, and Alice Y. Ting

Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Ave., Room 18-496, Cambridge, Massachusetts 02139, USA.

Abstract

We report an improved synthesis of 5-(5-oxohexahydrocyclopenta[c]thiophen-1-yl)pentanoic acid (ketone biotin, **1**) based on the intramolecular Pauson-Khand cyclization. The synthesis proceeds in 8 steps and in 2.7% overall yield from cyclohexene.

Site-specific conjugation of small-molecule probes to proteins of interest in living cells is a powerful technique for investigating protein function, interactions, and trafficking.¹ We recently reported the synthesis of a ketone analog of biotin, “ketone biotin” (**1**; Fig. 1), that is ligated site-specifically to a 15-amino-acid acceptor peptide (AP) by the *E. coli* enzyme biotin ligase (BirA).² Through this ketone handle, AP-tagged cell surface proteins can be labeled with hydrazide- or hydroxylamine-bearing biophysical probes in a two-step process (Fig. 2). The reported synthesis of **1** requires 12 steps from commercially available starting material, and proceeds in 0.66% overall yield, so a more efficient synthesis of **1** is desirable. Herein, we report a more concise synthesis that affords **1** in 8 steps and 2.7% overall yield, a 4-fold improvement over the previous synthesis.

Our approach to ketone biotin (**1**) centers around the intramolecular Pauson-Khand reaction, a [2+2+1] cycloaddition which has proven to be an effective technique for accessing bicyclo-[3,3,0]-oct-5-en-7-ones from 1,6-enynes,³ including allyl propargyl sulfides.^{4,5} We envisioned building the substituted allyl propargyl sulfide **3** from methyl 6-oxohexanoate **4** by addition of acetylene, followed by displacement of the resulting alcohol with allyl mercaptan. After complexation of the alkyne with octacarbonyldicobalt, a Pauson-Khand reaction should give α,β -unsaturated bicyclic ketone **2** (Scheme 1). Reduction of the alkene, followed by cleavage of the methyl ester to the free acid, should afford ketone biotin (**1**).

ating@mit.edu.

Supporting Information Available: Figure S1, experimental procedures and spectral data for new compounds. This material is available free of charge at <http://pubs.acs.org>.

The synthesis of **1** (Scheme 2) proceeded in eight steps. Methyl 6-oxohexanoate (**4**) was synthesized from commercially available cyclohexene in one step in 98% yield according to literature procedure.⁶ Addition of *in situ*-generated lithium trimethylsilylacetylene to **4** under anhydrous conditions led to the formation of the known propargyl alcohol **5**.⁷ Attempts to synthesize only the desired *R* enantiomer of **5** using Carreira addition of trimethylsilylacetylene to **4**⁸ led to almost exclusive formation of aldol self-condensation products, possibly due to the lack of a substituent at the α position of **4**. Removal of the trimethylsilyl protecting group with potassium carbonate afforded **6** in 66% yield from **4**. Following literature precedent,⁴ we next formed the hexacarbonyldicobalt complex of **6** by reaction with octacarbonyldicobalt in dichloromethane. Complexation of the alcohol to $\text{BF}_3 \cdot \text{Et}_2\text{O}$, followed by nucleophilic displacement with allyl mercaptan formed the Pauson-Khand cyclization precursor **7**, which was not isolated due to concerns about air stability.

When run at 70 °C in toluene, the Pauson-Khand reaction was found to proceed in 55% yield in 3 steps from **6**; however, the product formed was a mixture of diastereomers **2a** and **2b** (Fig. 3), with the desired diastereomer **2b** forming only 25% of the total product as determined by ^1H NMR analysis. The cobaltacyclic Pauson-Khand intermediate that places the valeric acid sidechain on the more sterically congested concave face of the bicyclic ring structure is disfavored relative to the intermediate where the sidechain is on the convex face, leading to preferential formation of product **2a** (See Fig. S1, supporting information).⁹ It was found that carrying out the reaction in refluxing toluene rather than at 70 °C gave a 57% total yield of cyclization products, while the proportion of **2b** increased to 35% of the total product formed.

Reducing the double bond of **2** proved more difficult than we had anticipated. Attempts at reduction using Pd/C or PtO_2 failed, even at high catalyst loadings and elevated temperatures and H_2 pressures. In light of this, we turned to copper-mediated 1,4-reduction. A procedure in which **2** was reacted with an *in-situ*-generated copper hydride complex proved to be effective, giving the reduced product **8** cleanly in 73% yield.¹⁰ The diastereomers were separated by preparative HPLC, and the methyl ester was deprotected with lithium hydroxide to afford **1**. The low overall yield for this pair of steps (10%) reflects the fact that only 35% of the diastereomeric mixture was the desired diastereomer.

This synthesis of ketone biotin provides a route to the protein-labeling reagent **1** that is four steps shorter and four-fold higher in yield. Moreover, there is precedent in the literature for intramolecular Pauson-Khand reactions on 1,6-enynes, including those with nitrogen or oxygen groups present between the double and triple bonds.³

This should provide expedient access to oxygenated, nitrogenated, or carbocyclic analogs of **1** (Scheme 3). The previously published synthetic route² cannot access these analogs because it makes use of the reduced acidity of the α carbons of sulfoxides to install the sidechain. Such analogs of ketone biotin should be useful in our continued efforts to probe the active site geometry and chemistry for both *E. coli* BirA and the high affinity biotin-binding protein streptavidin.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

We wish to acknowledge the financial support of NIH Grant R01 GM072670-01, the Sloan Foundation, the Dreyfus Foundation, and MIT. I.C. acknowledges the ACS Division of Organic Chemistry Graduate Fellowship by Wyeth Research. E.M. acknowledges the Novartis Summer Undergraduate Research Opportunities Program award.

References

1. Ting AY, Chen I. Curr. Opin. Biotech 2005;16:35. [PubMed: 15722013]and references cited therein
2. Chen I, Howarth M, Lin W, Ting AY. Nature Methods 2005;2:99. [PubMed: 15782206]
3. Pauson-Khand reaction: a Schore NE, Croudace MC. J. Org. Chem 1981;46:5436. b Pauson PL. Tetrahedron 1985;41:5855. c Geis O, Schmalz H-G. Angew. Chem. Int. Ed 1998;37:911. d Brummond KM, Kent JL. Tetrahedron 2000;56:3263.
4. Stumpf A, Jeong N, Sunghee H. Synlett 1997:205.
5. Castro J, Moyano A, Pericàs MA, Riera A. J. Org. Chem 1998;63:3346.
6. Schreiber SL, Claus RE, Reagan J. Tetrahedron Lett 1982;23:3867.
7. Hess C, Phillips RS. J. Chem. Soc. Perkin Trans. 1 2000:2821.
8. Anand NK, Carreira EM. J. Am. Chem. Soc 2001;123:9687. [PubMed: 11572696]
9. Mukai C, Kim JS, Uchiyama M, Sakamoto S, Hanaoka M. J. Chem. Soc. Perkin Trans. 1 1998:2903.
10. a Tsuda T, Hayashi T, Satomi H, Kawamoto T, Saesuga T. J. Org. Chem 1986;51:537. Dahnke, KR.; Paquette, LA. Organic Syntheses, Coll. 9. Wiley; New York: 1998. p. 396

Figure 1.
Biotin and ketone biotin.

Figure 2. Labeling of cell surface proteins (black rectangle) linked to acceptor peptide (blue rectangle) with ketone biotin and hydrazide-bearing biophysical probes (green circle).²

Scheme 1.
Retrosynthetic analysis of **1**

Scheme 2.
Synthesis of ketone biotin (**1**)

Figure 3.
Major (**2a**) and minor (**2b**) products of Pauson-Khand reaction.

Scheme 3.

Possible use of the Pauson-Khand reaction to synthesize congeners of **1**