VESICAL PAPILLOMA.

By J. SWIFT JOLY, M.D., F.R.C.S.

(*Senior Surgeon to St. Peter's Hospital for Stone.*)

It is never easy to say whether a villous growth of the bladder is innocent or malignant, unless a careful histological examination has been made; and with so many border-line cases, diagnosed more or less by the personal opinion of the surgeon or pathologist, it is dangerous to compare the results of one clinic with those of another. A growth may be considered benign by one observer, and diagnosed as malignant by another. I am therefore confining myself in this paper to cases I have examined and treated, in which the growth appeared to be benign when I first saw the patient.

Ætiology.

In the majority of cases there is no known ætiological factor. The growth originates in a normal bladder, without apparent cause. One may look on them as the result of some irritation, partly on the analogy of the cutaneous warts, and partly from consideration of their clinical history. The objection to this is, firstly that no such irritant has ever been found, except in the so-called aniline tumours of the bladder, and secondly that it is difficult to explain the predilection of these growths for the postro-lateral walls of the bladder. If they were caused by a diffusible irritant, one would expect to find them all over the vesical mucous membrane. Any irritating substance eliminated by the kidney is projected away from the usual site of the growth, while if it is eliminated by the opposite kidney the force of the ureteric efflux is soon dissipated in the bladder, and the irritant would not be carried across the middle line in any quantity. The postro-lateral wall is the most common site for all forms of bladder tumours, and this is probably only an example of the general rule that neoplasms originate in positions where the embryological development is most complicated.

The aniline tumours of the bladder must be counted as a class apart. In 1895 Rehn pointed out that vesical neoplasms were unusually common among the employees of chemical works producing aniline dyes and similar compounds. Although this was at first received with a good deal of scepticism, it has been confirmed by reports from every district in which the chemical industry has been established for some years. Leuenberger states that there are 33 times as many of these growths among the employees in this industry, than are found among an equal number of the general public. There are several peculiarities about these tumours. They do not appear in any locality until the chemical industry has been established for about ten years; the average age of the patients is from 34 to 47 which is distinctly below that of those with spontaneous growths; and they may develop from 10 to 17 years after the patient has left the industry (Oppenheimer). On the other hand, they resemble the ordinary growths clinically. Some start as papillomata which may ultimately become malignant, while others are definitely malignant from the beginning. In every factory there are one or more definite danger spots, and the growths are only found among those working in certain departments, and to a less degree in the neighbouring rooms. Oppenheimer gives an instance of six men working in one room, two remained healthy, two were successfully operated on for bladder growths, while two died of vesical cancer. There is no unanimity as to the nature of the dangerous chemical. Aniline, fuchsin, benzidine, beta-naphthylamine and naphthionic acid have all been incriminated, but no investigator has succeeded in producing bladder tumours in animals by

injections of any of these substances. Leuenberger found that aniline is hydrolised in the body into para-amidophenol, which is eliminated in the urine with ethylsulphuric acid, and he believes that this aromatic compound is the cause of the bladder tumours.⁽¹⁾

It must be obvious that only a minute proportion of the bladder growths observed in this, or in any country, can be put down to industrial causes. Out of 100 patients questioned on this point, I have only come across one where there was any probability of the disease being caused by the nature of his work. This man was employed in the still room of a petrol refinery. In two other cases the bladder growth was apparently secondary to a villous tumour of the renal pelvis, while in the remainder, no ætiological factor could be determined.

Age.—It is very difficult to give any accurate figures as to the age of the patient when the growth first appeared, as there is always a latent period in which there are no symptoms. In most cases this latent period is comparatively short, as a small growth is usually found when the patient is cystoscoped soon after the onset of the hæmaturia. On the other hand, one may find large growths associated with a very short clinical history. One patient reported difficulty in micturition for two months, and one attack of hæmaturia a fortnight before I first saw him, yet the growth was the size of a tangerine orange. Two other cases had large multiple growths where the duration of the symptoms was less than a year. It is also quite common to find recurrent growths on routine cystoscopy in cases where the patient has been absolutely free from symptoms. On the whole, however, one rarely finds a growth larger than a cherry when cystoscopy has been performed shortly after the first attack of hæmaturia. I believe that there are at least two factors which tend to increase the duration of the latent period. The first is the presence of residual urine, and the second is the position of the growth. If the bladder cannot empty itself the growth is protected from injury by a cushion of urine which prevents early bleeding. Growths springing from the anterior wall do not bleed as readily as those on the base, and one often finds large tumours in the former position which are quite symptomless. The following table gives the ages of 84 patients when they were first seen:—

From 20 to 29 years	...	...	...	I
„ 30 to 39 „	...	...	...	5
„ 40 to 49 „	...	...	...	12
„ 50 to 59 „	...	...	...	34
„ 60 to 69 „	...	...	...	21
„ 70 to 79 „	...	...	...	10
Over 90 years	...	...	...	I

The oldest patient was 93. He had two severe attacks of hæmaturia within a month. The growth was a simple papilloma about the size of a cherry.

Sex.—In 50 patients seen in private there were 33 males and 17 females. In this series approximately a third of the cases were females. At St. Peter's Hospital I do not see female out-patients; so the figures from this source are useless for determining the sex incidence.

Number.—Out of 77 untreated cases the growth was single in 52 and multiple in 25.

Site.—Single growths are almost always found on the postero-lateral wall of the bladder, a short distance above and behind the ureteric orifice. Occasionally they spring from the posterior or the upper portions of the lateral walls, but are

rarely found at the apex, unless an operation has been performed previously. If multiple, they tend to surround the trigone, and encroach on the internal meatus, especially on its lateral aspects. It is unusual to find a growth actually on the trigone. Occasionally small growths spring from the edge of the internal meatus, or from the upper portion of the prostatic urethra, but I have only seen them here when there are a large number of other tumours in the bladder itself.

When an operation has been followed by a recurrence, the largest growth is usually found on the supra-pubic scar. These tumours are frequently sessile, and tend to become malignant, although the original growth was innocent and pedunculated. If a recurrence follows cystoscopic fulguration it tends to lie at the site of the original tumour, close to the internal meatus, or in the urethra itself.

Mode of Spread.

Vesical papillomata spread by implantation. This is the only hypothesis which can account for the conditions observed clinically. Secondary growths form around the parent tumour, then on the opposite side of the bladder, and later round the internal meatus. This spreading is observed in cases which have not been treated, or when cystoscopic treatment has failed or been followed by a recurrence. In these conditions the epithelial cells are carried towards the internal meatus by the urinary stream. If a cystotomy has been performed, the tumour cells either become implanted in the wound at the time of the operation, or during convalescence. In these cases the flow of urine is directed towards the wound, and the tumour cells are implanted on the raw area.

Pathology.

These growths are formed of a central stem of fibrous tissue, from which branches are given off in all directions. The individual villi contain a central blood vessel, supported by the thinnest possible layer of fibrous tissue. They are covered by a thick layer of epithelium. If the growth is benign, the epithelial covering resembles the normal transitional epithelium of the bladder wall. It consists of from 4 to 6 layers of cells. The deeper layers are elongated and fusiform, and arranged at right angles to the central stem, while the superficial layers are flattened. The cells of the individual layers are uniform in shape, size, and staining capacity, but their nuclei are often more deeply stained than those of the normal bladder epithelium.

In other cases, the epithelium is much thicker and more luxuriant. It may have as many as from 12 to 15 layers of cells, and these cells are more irregular in size, shape, and in the manner in which they take up the stain. Their nuclei are also large, and often irregular, and mitotic figures are common. When these appearances are found, the growth is usually classed as malignant, although there may as yet be no down-growth of the epithelium through the basement membrane. When the epithelium takes on this luxuriant character, it often spreads as a continuous sheet from one villus to the next, in which case the flattened superficial cells are not found, and the whole epithelium appears to be formed of fusiform cells. True malignant degeneration is indicated by invasion of the submucous tissues, lymphatic spaces, etc., by epithelial cells.

Symptoms.

The symptom which led the patient to come for advice was noted in 100 cases of vesical papilloma. It was hæmaturia in 89 cases, difficulty in micturition in 8, pain in the renal region in one, passing gravel in one, and the fact that the

patient could only pass urine when lying on his side in one case. Three of the cases complaining of hæmaturia were admitted with clot retention. In two of the cases suffering from difficulty of micturition there were no symptoms of growth, the patients had prostatic obstruction and the papilloma was only discovered on routine cystoscopy; the remaining cases had typical attacks of hæmaturia which preceded the obstructive symptoms in five instances. The patient who complained of pain in the loin had a small papilloma over the corresponding ureteric orifice. I could not detect any evidence of renal disease and the pain disappeared as soon as the growth was destroyed. The man who could only pass urine when lying on his side had a very large growth overhanging the internal meatus. This case was remarkable as there was no hæmaturia. The patient who complained of passing gravel had several growths, with small phosphatic masses hanging on to the villi which formed the "gravel" he passed.

The hæmaturia is usually copious, intermittent, and painless. It comes on suddenly without any assignable cause, lasts from a few hours to a week or more, and ceases just as abruptly. It is not affected by exercise, though in some cases it may be relieved by rest in bed. It is painless, except when clots are passed, or when there is clot retention. Occasionally strangury is produced by impaction of a growth in the internal meatus. The interval between two attacks is variable, but as a rule it becomes shorter as the tumour increases in size. From three to six months' freedom from symptoms is not uncommon in the early stages of the disease, but if the growths are large and numerous, one usually finds that some blood is passed almost every day. This type of intermittent hæmaturia is usually the only symptom.

Difficulty in micturition is a late symptom, and indicates that the growths are so close to the internal meatus that they impede the urinary stream. This symptom is usually associated with a variable amount of residual urine. I have noticed as much as 20 ounces, but it is rarely more than six or eight. The patient can empty his bladder completely as soon as the growths are destroyed. Increased frequency of micturition occurs if the bladder becomes infected, or if the growth is so large that it fills the greater part of the organ. In the former case it is accompanied by pain and strangury, and the urine is always blood stained, so that the symptoms resemble those of advanced vesical carcinoma. It is uncommon to find a spontaneous cystitis, it is usually the result of instrumentation. Frequency of micturition due to the extreme size of the tumour is rare. I have only noticed it on two occasions.

Intermittent attacks of retention due to the growth falling like a ball valve on the internal meatus, are very rare. I have only seen one case.

Complications.—The most common complications are:—anæmia, cystitis, pyelitis, or pyonephrosis, hydronephrosis, retention of urine, clot retention, and most dangerous of all, malignant degeneration of the growth.

Course and Evolution.

A vesical papilloma, if left untreated, usually causes the death of the patient, as these tumours are very apt to become malignant. It is a mistake to classify them as benign, they should rather be considered as a pre-cancerous condition. Ewing⁽²⁾ states that 25 per cent. become malignant, but I believe that this is an understatement. Growths may persist for years without showing clinical evidence of malignancy, but I have only seen one case where symptoms persisted for more than 15 years without definite malignancy developing. This case was remarkable, as it may have been one of the rare cases of spontaneous cure.

A lady, aged 59, was sent to me with a diagnosis of vesical papilloma. She stated that 29 years before she was cystoscoped by a well-known surgeon, and told that she had a papilloma of her bladder. Operation was advised, but she refused it. The hæmaturia continued off and on for over 12 years, when she had an extremely severe attack, and passed large masses of "slough". The hæmaturia diminished for a time, but returned again about two years later, and continued off and on for about 8 years, when she had another severe attack, during which more "sloughs" were passed. Two years later she was persuaded to come to me for treatment but on cystoscopy I found her bladder clear, and could not detect any evidence of renal neoplasm. Unfortunately I was not able to obtain any accurate information as to her previous cystoscopy, as the surgeon who made the examination had died in the meantime. The evidence in this case is not conclusive, but I mention it as it is well known that a spontaneous cure does occasionally occur.

Apart from malignant degeneration, a villous growth of the bladder often gives rise to urinary sepsis which proves fatal. A spontaneous cystitis rarely arises except in cases where the growth is definitely malignant, but it is extremely easy to infect the bladder by instruments, and once this happens, the infection persists until the growth is removed or destroyed. Clot retention, or retention of urine, is particularly dangerous, and I have seen these complications give rise to acute urinary infection which soon proved fatal.

If none of these complications occur the patient may become so exsanguinated that he succumbs to some intercurrent disease.

Examination.

One can rarely feel a vesical papilloma. The palpable tumours of the bladder are usually malignant, and even they are rarely palpable while they are still operable. A growth which can be felt is almost always an inoperable carcinoma. I have only once succeeded in palpating a benign papilloma. In this case the growth was firm, was about the size of a tangerine orange, and felt like an indefinite doughy mass, which I took to be a blood clot. A bladder distended with urine or blood clot is naturally palpable, but as a rule physical examination gives very little information. The same applies to examination of the renal regions; it is usually negative.

Cystoscopy is *the* one method of making the diagnosis, and it should never be omitted if a papilloma is suspected. The tumour appears as a more or less spherical mass, which has a shaggy appearance. If the villi are long and slender they wave to and fro in the currents set up in the fluid medium, but if they are short and thick, the tumour looks much more solid, and resembles a cauliflower. Occasionally they are club shaped. When a single process is closely examined, it is usually possible to see the central vessel, which resembles a very fine hair line running down its long axis. This point is important, as the central vessel is only seen in true neoplasms and is never present in inflammatory conditions, such as bullous oedema. If the growth is pedunculated, it is seldom possible to see the stalk, as the lower branches lie on the bladder wall and hide it from view. When the growth is sessile, it resembles a mossy bank, or a broad tuft of grass.

It is easy to determine the presence of a growth, but it is often difficult to determine whether it is benign or malignant. There may be no difference between the cystoscopic appearances of a growth covered with normal epithelium, and one in which the epithelium is thick and atypical, yet the histologist may consider the former to be benign and the latter malignant. Even if a portion of the growth has been removed by a cystoscopic forceps for microscopic examination, it does

not always fix the diagnosis, as the nature of the epithelium may vary in different parts of the growth, and the early malignant changes may be confined to a small portion of it.

For years I used to remove a small portion of growth for microscopic examination, but have now given it up, as the information obtained was not always reliable. The surgeon must decide whether the growth is infiltrating or not, in order to determine the most suitable treatment. If there is no cancerous infiltration, a villous tumour can be destroyed by means of cystoscopic diathermy, no matter what may be the nature of its epithelial covering. If there is a definite infiltration round its base, nothing less than a wide resection of the whole thickness of the bladder wall will confer any lasting benefit on the patient. Fortunately, this point can be determined in most cases during the preliminary cystoscopy. A non-infiltrating tumour is always freely movable. Not only can it be moved from side to side, but it can also be lifted for some distance off the bladder wall. On the other hand, an infiltrating growth is always fixed to the deeper structures, and cannot be moved relatively to the bladder wall. The mobility of a growth can be easily determined by plunging a diathermy electrode into it, and turning on the current for about 30 seconds. This will make the tumour adhere to the electrode, and its mobility can be determined either by slowly withdrawing the electrode into the cystoscope, or else by moving the whole instrument backwards and forwards, and from side to side. If the growth follows the movement of the cystoscope, it is not infiltrating, and may be destroyed by fulguration, but if it does not, the bladder should be opened.

This method is only applicable to pedunculated tumours. If the growth is sessile, one must take the patient's history into account. If there has been no treatment, and especially if no operation has been performed, a large sessile tumour should be considered malignant. All those that I have seen were malignant. If, on the other hand, the growth is a sessile recurrence after removal of a pedunculated tumour, it may be benign. Cystoscopic treatment should be tried, but unless the growth rapidly disappears, the bladder should be opened, and a radical operation should be performed. When multiple pedunculated growths are found they are usually not infiltrating, and can be destroyed by diathermy.

Treatment.

The four methods of treatment extensively employed at present are:—

(1) Excision. (2) Diathermy applied through the open bladder—"open diathermy." (3) Implantation of radium. (4) Diathermy applied through a cystoscope—"cystoscopic diathermy" or "fulguration." There is not sufficient space to describe the technique of these methods, so I must confine myself to considering the indications for and the results obtained from them.

Excision.—Till the end of 1913 excision was the only radical method of dealing with villous tumours of the bladder. It was about this time that the first satisfactory diathermy apparatus, which could be used in conjunction with the cystoscope, was placed on the English market, although Edwin Beer described his method of "fulguration" in 1911. Since then it has been largely displaced by cystoscopic or open diathermy, and I am now convinced that excision is not to be recommended. At first sight, an open operation in which the whole growth is excised, together with the surrounding mucous membrane or even the whole thickness of the bladder wall, would appear to be sufficiently radical to effect a cure. But this is not so. Recurrence is quite common, and has been estimated at 25 to 30 per cent. of all cases.

I believe that it is better to group patients with a single growth apart from those in which more than one is found. In the former class, the risk of recurrence can be minimised by careful technique, but it can never be entirely eliminated. I operated on thirty cases of single growth before the war, and have reviewed them since. There were four recurrences, two of which occurred during the first year, and two were late. The early recurrences were treated, one by excision, and one by diathermy, both patients remaining well since. The late recurrences were remarkable. In one case a growth about the size of a pea was found at the site of the original growth six years after the operation. It was fulgurated, and the patient has since remained well. The second patient was free from symptoms for eight years, then had a slight attack of hæmaturia once a year for the next five years. When I saw him, thirteen years after his operation, he had an enormous malignant growth filling the greater part of his bladder.

Apart from these personal cases, I have seen in the last six years twelve cases of recurrent growth following excision of a single tumour done by other surgeons. In two of them the recurrent tumour was single, and was easily destroyed by fulguration; in the remainder I found multiple growths. In all of these cases the largest tumour was found under the supra-pubic scar. This is the most common site for recurrences after operation. In three of them the anterior growth was malignant. It was excised, but all these patients died of recurrence about a year afterwards. A fourth patient had an enormous tumour with foul cystitis. He was admitted for open diathermy, but died of pyelonephritis a fortnight after the operation. A fifth had a large recurrence in the scar, which I thought was malignant, and excised, but it proved to be benign. This patient, and the remaining five who were treated, by cystoscopic diathermy are still alive, but none of them is cured. If all the visible growths are destroyed, they invariably return with a fresh crop in three months. The best I can do is to keep them free from symptoms by constant cystoscopic treatment, but I think it probable they will all develop cancer sooner or later.

The operative results of multiple growths are most disappointing. I operated on nine cases of multiple papillomata before the war, they all recurred rapidly, and all the patients died in a comparatively short time. In two of them the growth became malignant, the others died from the combined effects of loss of blood and urinary sepsis.

I have only seen one case of multiple bladder growth originally treated by operation that survived for ten years. His history is unusual and worth recording. The patient, aged 38, was first operated on in August, 1918. He then had three growths in his bladder. Operation for recurrence was undertaken in December of the same year, when "several" growths were removed. Cystoscoped in April, 1919, told that there were "several splashes of growth in his bladder," and was advised to come to me for fulguration treatment. Consulted me October, 1920, then had been bleeding for seven months. Enormous masses of growth filling almost the whole bladder, only a small area of the apex and left side being clear. Fulguration treatment commenced, but was very tedious as the patient did not tolerate the cystoscope well. Bladder clear March, 1922. Cystoscoped every three months between this date and January, 1926. During this interval the bladder was usually clear, but five small growths were fulgurated. Nothing was found in January, 1926, and patient was asked to return the following May, but stayed away till September, and only came then because he noticed blood in his urine. I found a very large carcinoma, about two inches in diameter under the supra-pubic scar. It was non-villous. Resection of the anterior half of his bladder. I had to take away the anterior portion of the prostate and prostatic

urethra in order to get clear of the growth. Recovery. Since then numerous recurrences of villous growths have been fulgurated. In 1932 there was a recurrence of the carcinoma on the anterior wall of the bladder. It was excised, but a year later another malignant tumour appeared on the base. It was inoperable when first seen. Patient died early in 1934.

I feel that resection either cures the patient completely, or makes him distinctly worse. If the growths are multiple, it does not, in my opinion, hold out any hope of a cure. When only one growth is found recurrence is common, and usually multiple. Multiple recurrences are liable to become malignant and I have never succeeded in effecting a cure where I found them.

Diathermy applied through the open bladder.—I have only employed this method in a limited number of cases. They can be placed in two groups. In the first, the diathermy apparatus was not powerful enough, and in every case the tumour rapidly recurred. In the second group a powerful apparatus was used, and all these cases on which it was used have remained free from recurrence up to date. I have chiefly employed this method when the growth was unusually large, or when it was necessary to perform some other operation on the bladder at the same time. Resection of a papilloma combined with another operation, *e.g.*, prostatectomy, is in my experience invariably followed by a rapid recurrence.

Implantation of radium.—I have no personal experience of this method, as the radium at St. Peter's Hospital has hitherto been reserved for malignant tumours. It may be applied in two ways. The first is to open the bladder and to implant the radium directly into the growth. The second is to implant radon seeds through an operating cystoscope. It is possible to cause a disappearance of the growth by either method, but I have no information as to the late results.

Diathermy through the cystoscope.—This is, in my opinion, the method of choice for all non-infiltrating papillomata. Any growth that is freely movable can be destroyed by this means; if it is not freely movable it is far safer to assume that it is malignant and treat it accordingly. The objections to the method are:—

- (1) That it takes a considerable time to destroy a growth if it is at all large.
- (2) That it may be followed by a local recurrence.

The advantages are:—

- (1) That no incision is made in the patient's abdominal wall.
- (2) That the patient is not incapacitated for a considerable time. He can carry on his usual work during treatment.
- (3) That it rapidly controls the hæmorrhage.
- (4) That it gives rise to very few complications.
- (5) That it holds out the best prospect of a permanent cure in cases of multiple growths.

Many surgeons recommend it for a small single papilloma, and advise an open operation for large multiple growths. I believe that this is an entirely erroneous doctrine. I have pointed out that excision is a most unsatisfactory operation in the presence of multiple tumours, and the treatment of multiple recurrences after operation is always difficult and uncertain. I can only repeat the statement that if open excision does not cure the patient, it usually leaves him in an incurable condition.

On the other hand, fulguration is difficult under certain conditions. If the growth is situated near one of the ureters, it can be easily reached and treated

through an ordinary catheterising cystoscope. When I commenced fulguration, I soon found that this instrument was unsuitable for attacking growths in many parts of the bladder. A growth near the internal meatus, and one situated at the apex of the bladder cannot be reached by the usual catheterising cystoscope. I spent a considerable amount of time devising means of reaching small tumours no matter where they may be found, and I think it is now possible to destroy all growths no matter what their position. In this connection, I may quote the case of an American physician I saw in 1926.

In 1923 a vesical papilloma was diagnosed, and treated by radium, with the result that a supra-pubic fistula persisted for over a year. In 1924 the sinus was resected, and the growth removed. The following year a small recurrent growth was found on the anterior wall of the bladder, which his New York surgeon could not reach through the cystoscope. His bladder was reopened, and the growth removed. The next year he was in London, and noticed a recurrence of the hæmaturia. He was cystoscoped, and another tumour on the anterior wall was found. He was then sent to me, to see if I could do anything for him. When I saw him he was in an extremely nervous condition, as he thought that he would require an open operation every year of his life. I cystoscoped him, and burnt the growth away in my consulting room. He was so pleased with the result that he immediately purchased my instrument, and he wrote to me about fifteen months later saying that he had got another small recurrence at the same site, which his New York surgeon had no difficulty in destroying with my cystoscope.

The success of the treatment depends largely on the choice of instrument. If the bladder is full of growth, an irrigating cysto-urethroscope is the best. The treatment is carried out under a stream of circulating fluid, which washes away the blood, and which renders distention of the bladder unnecessary. When the growths are partially destroyed, and the bladder capacity restored, the surgeon can use an ordinary catheterising cystoscope. A cysto-urethroscope is necessary if the growth is situated in the posterior urethra, or springs from the edge of the internal meatus. Growths near the ureteric orifices are easily reached through an ordinary catheterising cystoscope. Those near the apex of the bladder are best treated through a semi-retrograde cystoscope, with my pattern electrode carrier. If the growth originates within half an inch from the internal meatus, this instrument, or a complete retrograde cystoscope, is necessary.

As much of the growth as possible should be burnt away at each sitting. A general anæsthetic is unnecessary, and unadvisable. The treatment is painless, unless the surgeon touches the bladder wall. The amount of pain also depends on the nature of the current. One with a low potential has a greater penetrative effect, and is much less painful than one with a higher potential. One should therefore choose an apparatus which only gives a short spark. A growth the size of a cherry can be destroyed in one or two sittings, but it is always necessary to cystoscope the patient about six weeks later to verify the result. Large multiple tumours can be rapidly destroyed. A modern diathermy apparatus is much more efficacious than those originally produced, and I now believe that I can destroy a large tumour in about a third of the time I took in 1914.

In judging the results of diathermy treatment, it is necessary to consider single tumours apart from multiple, just as one did when considering the results of operation. An early recurrence is not uncommon after fulguration for single growths; I have noticed this in six out of twenty-five cases, which were kept under cystoscopic observation for over five years. In all of them the recurrent tumour was at the site of the original and was either due to incomplete

treatment, or to an implantation tumour which was not visible at the time. These recurrences were all fulgurated, and the patients have since remained well. In two cases a recurrence was noticed three years after the original tumour was destroyed; one patient recently consulted me for difficulty in micturition fourteen years after I had destroyed a bladder growth. I found that he was suffering from a prostatic carcinoma, but on cystoscopying him I found a tiny papilloma at the site of his original growth.

If multiple growths are found, one can still obtain good results by diathermy, but one must naturally expect a considerable number of recurrences. Sometimes these cases require very prolonged observation. The following case serves as an illustration. A man of forty-five was sent to me in 1914. His doctor stated that two years previously the patient had been diagnosed as suffering from an inoperable carcinoma of the bladder, but he thought there must have been an error, as the patient's condition was not bad. I found that his bladder was almost completely filled with growths. I fulgurated them, and destroyed all of them in six months. Towards the end of this year he had a small recurrence which was burnt away, and a second recurrence was destroyed in 1915. During the years 1916-18 about six small recurrences were burnt away, while I was on Active Service. In 1919 I destroyed two small growths, and in 1920, one. His bladder then remained clear for over five years, and he was discharged as cured in 1926. In 1928 he returned as he had another attack of hæmaturia, and I found two small tumours which were easily destroyed. Since then he has remained well. An open operation would have been followed by a rapid recurrence in such a case, and I feel that he would not be alive now if the growths had been excised. In these cases recurrence is the rule, but in all that I have treated I have been able to clear the bladder of growth, and keep the patient free from symptoms.

Conclusions.

(1) Recurrences after any form of treatment may be divided into three groups. The first are cases of imperfect removal, and the recurrent tumour is found within a few months at the site of the original growth. The second are implantation tumours. If an open operation has been performed they usually lie at or near the supra-pubic scar; after cystoscopic treatment they are found as a rule near the internal meatus, or in the posterior urethra. The third are the late recurrences which are only found several years after the original growth has been removed. These late recurrences should make one very cautious in giving a good prognosis, as some of them are malignant.

(2) Open operation is followed by 10 to 20 per cent. of recurrences in cases of single growths, and recurrence is practically certain in every case where multiple growths are found. Approximately 10 per cent. of these recurrences are malignant when the patient is first cystoscoped after operation, and others may become malignant later on. I have only been able to effect a "cure" when the recurrence is single. Cystoscopic diathermy is followed by a larger proportion of recurrences (25-30%), but they are amenable to treatment, and a good result may be ultimately expected. I have not seen a malignant recurrence in any case treated entirely by fulguration.

(3) Diathermy is the safest method of treating vesical papillomata, and excision should only be employed in exceptional circumstances.

REFERENCES.

- (1) *Handbuch der Urologie*: Lichtenberg, Voelcker, and Wildbolz. Vol. V., p. 175, et seq.
- (2) Ewing, J., *Neoplastic Diseases*, 3rd ed., 1928, p. 910.