Published in final edited form as:

J Am Chem Soc. 2007 August 29; 129(34): 10310–10311. doi:10.1021/ja072841c.

Resolution of the Spectroscopy vs. Crystallography issue for NO intermediates of Nitrite Reductase from *Rhodobacter sphaeroides*

Somdatta Ghosh[†], Abhishek Dey[†], Oleg M. Usov[‡], Yan Sun[‡], Vladimir M. Grigoryants[‡], Charles P Scholes^{*,‡}, and Edward I. Solomon^{*,†}

Department of Chemistry, Stanford University, Stanford, California 94305, Department of Chemistry, University at Albany, State University of New York, Albany, New York 12222

Copper nitrite reductase (NiR) is a homotrimeric enzyme, containing a T1 copper site, which transfers electrons to the T2 catalytic site, where nitrite is reduced by one electron to nitric oxide ($\text{NO}_2^- + 2\text{H}^+ + \text{e}^- \rightarrow \text{NO} + \text{H}_2\text{O}$).^{1–3} Recently a side-on bound copper nitrosyl complex of NiR has been crystallized by reacting the reduced enzyme with excess NO.⁴ Based on the EPR spectrum of the species generated by the reaction of reduced NiR with saturated NO in solution, it has been assigned as an $\eta^2\text{-NO}^-\text{Cu}^{2+}$ species. However, spectroscopic data on several Cu-nitrosyl model complexes⁵ and an NO adduct formed by reacting reduced NiR with nitrite⁶ and DFT calculations on an $\eta^2\text{-NOCu}$ species⁷ describe these complexes as $\text{Cu}^+\text{NO}^\bullet$. In this study, we define the species generated in solution by reacting reduced NiR with NO, using EPR, MCD and ENDOR spectroscopy and correlate these with that observed by crystallography.

Reduced NiR reacted with saturated NO solution generates the EPR spectrum in Figure 1A red, which is equivalent to the spectrum reported by Tocheva et al.⁴ This species has an EPR signal characteristic of an oxidized T2 Cu ($g_{\parallel} = 2.30$ and $A_{\parallel} = 117$ G). The MCD spectrum (Fig. 1B, red) has low energy d-d bands, also indicative of a T2 cupric complex. Both EPR and MCD spectra show a negligible T1 contribution indicating that this site is reduced (Fig. S1).

Nitrite binds to the T2 site of the resting WT enzyme,^{8,9} but the EPR and, in particular, the MCD features of the nitrite bound T2 site are obscured by the dominant T1 Cu signals. However under these conditions (nitrite bound), the T1 Cu could be selectively reduced with ascorbate. The EPR spectrum of the nitrite bound T2 species (Fig. 1A, blue) is identical to that produced above by reacting reduced NiR with excess NO. The MCD spectrum (Fig. 1B, blue) of the T1 reduced T2-nitrite bound form is also identical to that generated by the reduced enzyme with NO. The T2 EPR spectrum and the MCD d-d bands of the resting WT enzyme are very sensitive to nitrite binding (Fig. S1C, D). Thus the EPR and MCD data strongly indicate that the species generated by the reaction of excess NO with reduced NiR has T1 reduced and T2 oxidized with nitrite bound.

Further N^{14} and H^1 ENDOR data for this species generated by the reaction of reduced NiR with an NO solution (Fig. 1C, D, red) and those for a form with T1+T2 oxidized and nitrite bound (collected at $g=2.266$ where there is no contribution from T1, Fig. 1C, D, blue) have the same features. The nitrogen features (Fig. 1C) from $\text{Cu}^{2+}\text{NO}_2^-$ bound T2 are from its histidines, and these features all show lower ENDOR frequencies and smaller hyperfine couplings due to the change of the T2 axial ligand from aquo to nitrite.¹⁰

*To whom all correspondence should be addressed Edward.Solomon@stanford.edu, cps14@albany.edu.

[†]Department of Chemistry, Stanford University, CA

[‡]Department of Chemistry, State University of New York, Albany

From the above results the reaction of reduced NiR with excess NO gives T1 reduced T2 $\text{Cu}^{2+}\text{-NO}_2^-$. However, the reaction of the reduced enzyme with limited NO has an EPR signal (Fig. 2A, green), characteristic of a $\text{Cu}^+\text{NO}^\bullet$ species.⁶ This species lacks d-d bands in the MCD spectrum (Fig. S3), further confirming that the Cu is reduced. Therefore, reduced NiR forms a $\text{Cu}^+\text{NO}^\bullet$ species at low concentrations of NO.

The reaction of reduced NiR with excess NO involves two one-electron oxidations; reduced T2 $\rightarrow$ oxidized T2, and $\text{NO} \rightarrow \text{NO}_2^-$. A similar reaction has been observed for the coupled binuclear Cu active sites in hemocyanins¹¹ (Scheme S1). Deoxyhemocyanin (2Cu^+) reacts with NO generating N_2O , and a met (2Cu^{2+}) form, which further reacts with NO to generate the half-met ($\text{Cu}^{2+}\text{Cu}^+$) nitrite bound form. N_2O evolution is in fact detected during the reaction of reduced NiR with excess NO, using nitrous oxide reductase ($35 \mu\text{mol}$ N_2O reduced $\text{min}^{-1}\text{mg}^{-1}$ of enzyme), which requires N_2O as its substrate.¹² Reaction of the resting WT NiR (analogous to the met form of hemocyanin) with NO generates the T1 reduced, T2 oxidized form of the enzyme ($g_{\parallel}=2.30$, $A_{\parallel}=150\text{G}$), which converts to the nitrite bound T2 form (Fig. 2B, D $\rightarrow$ F $\rightarrow$ G, Scheme 1), indicating NO is oxidized to nitrite and nitrite binding is slow.

To evaluate the entire reaction mechanism, reduced NiR was reacted with limited NO solutions and intermediates were trapped at different times. We were able to trap the $\text{Cu}^+\text{NO}^\bullet$ form at shorter times, leading to the formation of the T1 reduced, T2 oxidized form, and ultimately generating the T1 reduced, T2 oxidized nitrite bound form (Fig. 2A, Scheme 1 A $\rightarrow$ B $\rightarrow$ F $\rightarrow$ G) at longer times. A reaction mechanism consistent with these results is presented in Scheme 1. Note that in the scheme, intermediate E is not observed because of rapid ET from T2 $\rightarrow$ T1.¹³ The experimental details and kinetic rates are provided in the supporting information.

There is still one issue to resolve on correlating the spectroscopy with crystallography. The Cu-NO species trapped in the crystal structure has a side-on conformation.⁴ DFT calculations indicate that this species is best described as a $\text{Cu}^+\text{-NO}^\bullet$ species.⁷ However an end-on Cu-NO species is also trapped after limited exposure to NO gas (B, in scheme 1) which is equivalent to the species generated by reacting reduced NiR with nitrite ($\eta^1\text{-CuNO}$).⁶ Geometry optimized DFT calculations (B3LYP/6-311g* on Cu, N, O and 6-31g* on C, H)¹⁴ of both side-on (Fig. 3, red) and end-on (Fig. 3, blue) Cu-NO adducts indicate that both forms have a $\text{Cu}^+\text{NO}^\bullet$ electronic configuration and that the end-on structure is more stable by 7 kcal/mol. However the orientation of the 2nd sphere aspartic acid residue (Asp129)^{9,15} appears to play a key role in determining the geometry of the Cu-NO species. In the side-on adduct this residue is perpendicular to the Cu-N-O plane and H-bonds to the O of N- O (Fig. 3 red) while in the end-on structure the Asp129 residue H-bonds to the O of NO and is parallel to the Cu-N-O plane (Fig. 3 blue). Both orientations of the Asp129 residue have been observed in crystal structures.¹⁵ The Asp129 residue exists in the perpendicular orientation in the crystal structures of the oxidized and reduced NiR and H-bonds to the axial ligand. However, to form the more stable end-on $\text{Cu}^+\text{NO}^\bullet$ adduct this aspartate residue has to rotate 90° which can occur in solution but may not be feasible in a crystal.

In summary, the species generated in solution during the reaction of reduced NiR with excess NO is the T1 reduced, T2 oxidized nitrite bound form and not the side-on Cu-NO species observed in the crystal structure. The latter is best described as $\text{Cu}^+\text{NO}^\bullet$ rather than $\text{Cu}^{2+}\text{NO}^-$. The mechanism for the reaction of reduced NiR with excess NO involves its two electron oxidation, generating N_2O . This oxidized form of the enzyme oxidizes NO generating nitrite bound to the oxidized T2 site generated by ET from T2 $\rightarrow$ T1. DFT calculations provide insight into factors which favor the $\eta^2\text{-NO}^\bullet\text{Cu}^+$ complex in the crystal structure.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This research is supported by NIH DK- 31450 (E.I.S), NIH EB00326929 (C.P.S).

References

1. Godden JW, Turley S, Teller C, Adman ET, Liu MY, Payne WJ, LeGall PJ. *Sci* 1991;253:438.
2. Murphy MEP, Turley S, Adman ET. *J Biol Chem* 1997;272:28455. [PubMed: 9353305]
3. Suzuki S, Kataoka K, Yamaguchi K. *Acc Chem Res* 2000;33:728. [PubMed: 11041837]
4. Tocheva EI, Rosell FI, Mauk AG, Murphy ME. *Sci* 2004;304:867.
5. Ruggiero CE, Carrier SM, Antholine WE, Whittaker JW, Cramer CJ, Tolman WB. *J Am Chem Soc* 1993;115:11285.
6. Usov OM, Sun Y, Grigoryants VM, Shapleigh JP, Scholes CP. *J Am Chem Soc* 2006;128:13102. [PubMed: 17017790]
7. Wasbotten IH, Ghosh A. *J Am Chem Soc* 2005;127:15384. [PubMed: 16262398]
8. Strange RW, Murphy LM, Dodd FE, Abraham ZHL, Eady RR, Smith BE, Hasnain SS. *J Mol Biol* 1999;287:1001. [PubMed: 10222206]
9. Zhao Y, Lukoyanov DA, Toropov YV, Wu K, Shapleigh JP, Scholes CP. *Biochem* 2002;41:7464. [PubMed: 12044180]
10. Veselov A, Olesen K, Sienkiewicz A, Shapleigh JP, Scholes CP. *Biochem* 1998;37:6095. [PubMed: 9558348]
11. Solomon, EI. Ch 2. In: Spiro, TJ., editor. *Copper Proteins*. Acad. Press; NY: 1981.
12. There exist micro-organisms with a truncated denitrification chain catalyzing the reduction of NO_2^- to NO .[#] This mechanism suggests that in the presence of excess NO , NiR may also be used to reduce NO to N_2O .[#] Casella S, Shapleigh JP, Toffanin A, Basaglia M. *Biochem Soc Trans* 2006;34:130. [PubMed: 16417501]
13. Farver O, Eady RR, Abraham ZHL, Pecht I. *Febs Lett* 1998;436:239. [PubMed: 9781686]
14. Frisch, MJ., et al., editors. *Gaussian 03, Revision C. 02*. Full Reference in SI
15. Antonyuk SV, Strange RW, Sawers G, Eady RR, Hasnain SS. *Proc Natl Acad Sci* 2005;34:12041. [PubMed: 16093314]

Figure 1. Overlay of reduced NiR with excess NO (red) and nitrite bound NiR (blue) A) EPR B) MCD, C) N^{14} ENDOR, D) H^1 ENDOR, with exchangeable features with 8.5 MHz coupling. Parts A and B have nitrite bound to T2 site with the T1 Cu reduced to remove intense overlaying T1 features. ENDOR data are collected at 1.078T where there is no contribution from T1 Cu.

Figure 2.

EPR spectra A) following the reaction of reduced NiR with sub-stoichiometric NO solution (10:1 enzyme: NO) trapped at 1 second (green) 4 second (red) and 18 seconds (purple). B) following the reaction of the resting oxidized enzyme (black) with NO gas at 4 seconds (red) and 90 seconds (purple). Letters correspond to Scheme 1.

Figure 3. Overlay of the DFT optimized side-on bound NO (red) and end-on bound NO (blue) structures indicating the different orientations of the aspartate residue.

Scheme 1.

Proposed mechanism for the formation of nitrite bound T2 NiR from reduced enzyme with excess NO. Red boxes = species observed.