Antibody Responses to *Cryptosporidium* Antigen in HIV-positive Patients in the Republic of Korea

Sang-Mee Guk¹, Jong-Yil Chai¹, Yung-Oh Shin² and Min Seo^{3,*}

¹Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, and Institute of Endemic Diseases, Seoul National University Medical Research Center, Seoul 110-799, Korea; ²Department of Microbiology, College of Medicine, Kangwon National University, Chuncheon 200-701, Korea; ³Department of Parasitology, College of Medicine, Dankook University, Cheonan 330-714, Korea

Abstract: The diagnosis of cryptosporidiosis has been carried out using coprologic techniques in the Republic of Korea. However, antibody responses to *Cryptosporidium* have rarely been studied. Serum antibodies from HIV-positive/oocyst-positive Korean patients recognized significantly 31 and 27 kDa antigens, and HIV-negative/oocyst-positive individuals clearly reacted to 15/17 kDa antigens. Compared with oocyst-positive cases, 18.7% and 75.8% of sera from HIV-positive patients reacted to 31 and 27 kDa antigens. Only 11.1% of HIV-negative individuals reacted to 15/17 kDa. Based on these findings, serum antibody responses were different between HIV-positive and HIV-negative individuals infected with *Cryptosporidium*, and it is suggested that HIV-positive patients are more frequently exposed to *C. parvum* compared to HIV-negative individuals.

Key words: *Cryptosporidium parvum*, serologic response, antigenic protein, HIV

INTRODUCTION

Cryptosporidium is an intestinal protozoan causing self-limiting diarrhea in immunocompetent individuals, but can be life-threatening in immunocompromised patients [1]. Hence, there is a need to evaluate serologic responses of immunocompromised patients against this pathogen, particularly in HIV-infected individuals. Sorvillo et al. [2] reported that 120 (2.8%) HIV-positive patients were diagnosed as cryptosporidiosis among 4,247 examined, but the overall positive rate for *Cryptosporidium* was 5.3% after follow-ups until death [2]. In India, the positive rate was 9.2% among HIV-positive individuals with diarrhea [3], and oocysts were detected up to 4% of AIDS patients in USA [4]. According to the report of National Institute of Health, a total of 5,155 Koreans were infected with HIV as of January 2008, and the number has been rapidly increasing [5]. There were few reports on opportunistic parasitic infections in the Republic of Korea; oocysts of *Cryptosporidium parvum* or *Isospora belli* were 3.1% of HIV-infected individuals [6], and Guk et al. [7] reported that 10.5% of AIDS patients suffered from *C. parvum* infection. However, the reports on *C. parvum* were insufficient considering its epidemiologic and clinical significances.

The diagnosis of cryptosporidiosis is usually based on exami-

nation of fecal smears with acid-fast stains [8]. However, fecal examinations may underestimate the prevalence of *Cryptosporidium* infection, not only because this infection is not a part of standard parasitological examinations, but because the duration of oocyst shedding can be short and intermittent [9]. Thus, serologic assays may be an alternative way for investigating the epidemiology of *Cryptosporidium* infection. Although detection of specific antibodies should not be necessarily regarded as an active infection, some antigens identified by immunoblot analysis are considered as excellent markers of infection, especially for 15/17 kDa and 27 kDa proteins [10]. The sensitivity and specificity of ELISA using 27 kDa antigen (CP23) were 0.86 and 0.86, respectively, for predicting its infection, and a serological assay using 27 kDa antigen was proved to be more accurate regarding the community levels of *Cryptosporidium* infection [11,12]. By assaying the antibody responses against 27 kDa and 17 kDa antigens, Sandhu et al. [13] emphasized a high risk of HIV-infected population to *Cryptosporidium* infection [13].

In the Republic of Korea, however, epidemiological surveys for *Cryptosporidium* infection using serological assays have never been undertaken either on healthy population or on HIV-infected patients. The present study was performed to investigate serum antibody responses in HIV-positive and HIV-negative individuals compared with the results of oocyst examinations. This report may provide basic data to evaluate the risk of exposure to *Cryptosporidium* infection in each group, and to study the pathogenic

• Received 14 April 2008, accepted after revision 24 May 2008.

* Corresponding author (bbbenji@freechal.com)

proteins of *Cryptosporidium*.

MATERIALS AND METHODS

Collection of sera

Serum specimens were obtained from the following groups;

1) Eight oocyst-confirmed cryptosporidiosis patients living in Ssangbong-ri, Iyang-myon, Jeollanam-do, an endemic area of cryptosporidiosis. Their ages were distributed from 50 to 79 yr, and they discharged *C. parvum* oocysts at routine examinations over 3 times (positive control 1). The samples were collected during 1996-1997.

2) Four HIV-positive patients visiting Seoul National University Hospital during 1995-1998, who were confirmed to be heavily infected with *C. parvum* by microscopic examinations (positive control 2); all of them were males, and 29-34 yr in age range.

3) Ninety-one HIV-positive patients, who were not examined for *C. parvum* infection and whose HIV infections were confirmed by western blotting in Department of AIDS, National Institute of Health, Republic of Korea, from October 1993 to April 1995. The sex ratio and age could not be investigated (control group 1).

4) Twenty-seven healthy individuals not examined for *C. parvum* infection, all of whom were students in their 20s, attending Seoul National University College of Medicine. The sex ratio was not investigated, and the samples were collected in 1997 (control group 2).

5) In each test, human sera uninfected with any parasites were used as a negative control.

Antigen preparation

The oocysts were purified by ether extraction and discontinuous sucrose gradients from the feces of calves, experimentally infected with the Korean isolate of *C. parvum*. The feces were collected daily, mixed with equal volumes of 2.5% potassium dichromate ($K_2Cr_2O_7$), and stored at 4°C. Oocysts were isolated by means of discontinuous sucrose gradients [14]. Briefly, Sheather's solution containing sucrose was diluted with 0.025 M phosphate-buffered saline (PBS) supplemented with 1% Tween 80 to make 1 : 2 and 1 : 4 gradient solutions. After centrifugation, the oocysts were collected from the interface of the 2 gradients. Purified oocysts were washed with PBS and then used for the antigen preparation. Antigens of *C. parvum* were prepared as previously described [15]. In brief, oocysts were disrupted by sonication followed by 5 freeze-thaw cycles. The

lysate was clarified by centrifugation, and the supernatant was used as the crude antigen.

Immunoblotting

The soluble fraction was separated by SDS-PAGE, and electrophoretically transferred onto a polyvinylidene fluoride (PVDF) membrane (Immobilon, Millipore Corp., Bedford, Massachusetts, USA) at 0.5 mA for 1-2 hr. The membranes were blocked in 0.2% skim milk/PBS (pH 7.4) for 2 hr, and incubated with serum samples for 12 hr. The secondary antibody was peroxidase-conjugated goat anti-human IgG (Cappel, Cochranville, Pennsylvania, USA), color developed using a substrate solution (0.6% chloronaphtol, 0.2% diaminobenzidine, 0.02% H_2O_2 /PBS). All examinations were repeated 2-3 times.

RESULTS

The soluble extract of *C. parvum* showed multiple bands ranging 14-148 kDa in 12.5% SDS-PAGE gel. The sera from HIV-positive/oocyst-positive patients reacted with several protein bands, ranging 15/17-128 kDa. The positive rate to either of 62, 31 or 27 kDa band was 100%. The *C. parvum*-negative sera cross-reacted with 128, 124, 114, and 62 kDa bands (Fig. 1).

The 15/17 kDa antigen was recognized by 2 sera (50%). All sera (100%) from HIV-negative/oocyst-positive patients react-

Fig. 1. *Cryptosporidium parvum* antigenic proteins recognized by sera from HIV-positive/*C. parvum* oocyst-positive patients. M, molecular weight (kDa); Ag, oocyst antigen; lanes 1-4, HIV-positive/oocyst-positive patients; lanes 5-7, HIV-positive/oocyst-negative control patients. Arrows point to 27 and 15/17 kDa bands.

ed against 15/17 kDa, and the reaction rates were higher than 60% against 55, 48, and 44 kDa antigens (Fig. 2). The 31 kDa and 27 kDa antigens were recognized only by 3 sera (37.5%), the rate of which was significantly lower compared to the sera of HIV-

Fig. 2. *C. parvum* proteins recognized by sera from HIV-negative/oocyst-positive individuals. M, molecular weight (kDa); lanes 1-4, negative control; lanes 5-12, HIV-negative/oocyst-positive individuals. Arrows point to 55, 44, 31, 27, and 15/17 kDa bands.

positive/oocyst-positive patients. Overall, reaction to 15/17 kDa antigen was the highest (10/12), followed by 44 kDa (8/12), 55, 31, and 27 kDa (all were 7/12) antigens in the positive control groups. In particular, in control group 1, the sera of the patient numbers 1, 84, and 88 strongly reacted with 15/17 kDa antigen (Fig. 3). Compared with the results of positive groups, reactions against 44 kDa (84/118), 27 kDa (73/118), 55 kDa (34/118), and 15/17 kDa (14/118) were observed in control 1 and 2 groups (Figs. 3, 4). Collectively, the sera from HIV-positive/oocyst-positive patients significantly reacted to 27 and 31 kDa, whereas the sera of HIV-negative/oocyst-positive individuals reacted mostly to 15/17 kDa antigens.

The sera of HIV-positive individuals in control group 1 reacted against more proteins than the sera of HIV-negative individuals

Fig. 4. Immunoblot with sera of HIV-negative individuals. Arrows point to 55, 44, 31, 27, and 15/17 kDa bands.

Fig. 3. Immunoblot with sera of HIV-positive patients. Arrows point to 55, 44, 31, 27, and 15/17 kDa bands.

in control group 2. The 27 kDa antigen was the most frequently recognized by the sera of control group 1 (75.8%), whereas the 31 kDa antigen was recognized by 18.7% of the sera. In control group 2, the 31 kDa antigen was not reactive, whereas the 27 kDa was recognized by 14.8% of sera. The sera of 11.1% reacted weakly to the 15/17 kDa antigen.

The 27 and 15/17 kDa antigens were recognized by sera 100% and 50%, respectively, in HIV-positive/oocyst-positive individuals, and 37.5% and 100%, respectively, in HIV-negative/oocyst-positive individuals. Furthermore, the combination of 31 and 27 kDa antigens were recognized 100% by HIV-positive/oocyst-positive sera, and 18.7% in control group 1.

DISCUSSION

The present study is the first to study the serologic responses of humans to *Cryptosporidium* antigens in the Republic of Korea. Although many proteins were identified, a particular emphasis must be focused on 27 and 15/17 kDa antigens, which reacted significantly in oocyst-positive groups and are also known as serologic markers [10-13]. According to Sandhu et al. [13], *Cryptosporidium* infections are supposed to be common among HIV-positive individuals, whereas the absence of overt clinical cryptosporidiosis is due to the presence of intact immune responses among HIV-positive individuals [13]. In the present study, immunoblot results demonstrated that sera from HIV-positive individuals recognized more protein fractions of *C. parvum* than those from HIV-negative individuals. It has been reported that strong serological responses to 27 kDa antigen in immunosuppressed individuals were associated with a reduced risk of developing diarrhea [16]. In the present study, it has been shown that 31 and 27 kDa antigens were significantly reacted in HIV-positive/oocyst-positive group. These results indicate that we could estimate *C. parvum*-infection in HIV-positive patients through the reaction of 27 kDa antigen and the combination of 27 and 31 kDa antigens. In particular, 61.9% of control groups 1 and 2 (*C. parvum*-infection was not examined) reacted with 27 kDa antigen. Only with these results, the real infection rate of *C. parvum* infection cannot be explained. However, in control group 1, the combination of 27 and 31 kDa antigens were recognized by 18.7% and it is suggested that they have a possibility of *C. parvum* infection.

However, serologic assays cannot solely determine *C. parvum* infection. Given there was little agreement between serum ELISA and fecal tests on samples collected from adult cows [17], sero-

logical assays should be used to support fecal examinations. The reported infection rate of randomly selected inhabitants in Seoul was only 0.5% [18], whereas another report identified that the majority (94.6%) of oocyst-positive cases were people over 50 yr of age [19]. Furthermore, Seo et al. [20] reported that *C. parvum* infections were confined to the over 30-yr age group [20]. In the present study, it is considered that HIV-negative/oocyst-positive individuals were repeatedly infected with *C. parvum* in the endemic area. Their sera reacted with mainly 55, 44, and 15/17 kDa antigens, and we focused on the reaction with 15/17 kDa antigen. Since members of control group 2 were young (in their 20s) medical college students, living in Seoul, their *C. parvum* infection rate should have been low. Their sera showed 11.1% reactivity against 15/17 kDa antigens.

In humans infected with *Cryptosporidium*, most oocyst-derived glycoproteins and sporozoite surface proteins, including 31, 27, 15/17 kDa, were recognized by serum antibodies [10]. A study of laboratory-confirmed cryptosporidiosis patients showed that IgG levels against 27 kDa antigen tended to remain above the detection threshold for at least 2 yr after infection, whereas the levels of antibody against 17 kDa antigen decreased rapidly [11]. The higher reactivity against 27 kDa of HIV-positive sera supports this phenomenon. It has been reported that HIV-positive individuals were more likely to have at least one infection during an 8.5-yr follow-up period compared to HIV-negative individuals (59% vs 30%) [16]. Since most infections are acquired from water or food contaminated with infectious oocysts [21], these results may reflect continuous exposure of people to contaminated environments. Although the results of the present study are important, they must be considered in light of methodological limitations, such as limited microscopically confirmed cases of cryptosporidiosis, and no information on their clinical status. Based on the findings of the present study, it is suggested that serum antibody responses were different between HIV-positive and HIV-negative individuals infected with *Cryptosporidium*. In addition, it is possible that HIV-positive individuals may be more frequently exposed to *Cryptosporidium* infection than immunocompetent people in the Republic of Korea.

ACKNOWLEDGEMENTS

This work was funded by Medical Science Research Center of Dankook University Medical Center in 2007. We are also thankful to Dr. M.H. Choi (Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, Seoul,

Korea) for collection of blood samples from HIV+ patients.

REFERENCES

1. Thompson RC, Olson ME, Zhu G, Enomoto S, Abrahamsen MS, Hijjawi NS. *Cryptosporidium* and cryptosporidiosis. *Adv Parasitol* 2005; 59: 77-158.
2. Sorvillo F, Beall G, Turner PA, Beer VL, Kovacs AA, Kraus P, Masters D, Kerndt PR. Seasonality and factors associated with cryptosporidiosis among individuals with HIV infection. *Epidemiol Infect* 1998; 121: 197-204.
3. Morgan UM, Pallant L, Dwyer BW, Forbes DA, Rich G, Thomson RC. Comparison of PRC and microscopy for detection of *Cryptosporidium parvum* in human fecal specimens: clinical trial. *J Clin Microbiol* 1998; 36: 995-998.
4. Soave R. Cryptosporidiosis and isosporiasis in patients with AIDS. *Infect Dis Clin North Am* 1988; 2: 485-493.
5. Korea Center for Disease Control and Prevention, Ministry of Health and Welfare, Republic of Korea. News from Korea Center for Disease Control and Prevention, Ministry of Health and Welfare, Seoul, Korea, 2008.
6. Choi HJ, Park SW, Kim HB, Kim NJ, Shin DH, Oh MD, Jung HC, Song IS, Choe KW. The gastrointestinal infections in Korean patients with human immunodeficiency virus infection. *Korean J Gastroenterol* 1999; 33: 482-488.
7. Guk SM, Seo M, Park YK, Oh MD, Choe KW, Kim JL, Choi MH, Hong ST, Chai JY. Parasitic infections in HIV-infected patients who visited Seoul National University Hospital during the period 1995-2003. *Korean J Parasitol* 2005; 43: 1-5.
8. Ramakrishna BS. Prevalence of intestinal pathogens in HIV patients with diarrhea: implications for treatment. *Indian J Pediatr* 1999; 66: 85-91.
9. Weber R, Bryan RT, Bishop HS, Wahlquist SP, Sullivan JJ, Juranek DD. Threshold of detection of *Cryptosporidium* oocysts in human stool specimens: evidence of low sensitivity of current diagnostic methods. *J Clin Microbiol* 1991; 29: 1323-1327.
10. Moss DM, Chappell CL, Okhuysen PC, DuPont HL, Arrowood MJ, Hightower AW, Lammie PJ. The antibody response to 27, 17, and 15 kDa *Cryptosporidium* antigens following experimental infection in humans. *J Infect Dis* 1998; 178: 827-833.
11. Ong CS, Li SA, Priest JW, Copes R, Khan M, Fyfe MW, Marion SA, Roberts JM, Lammie PJ, Isaac-Renton JL. Enzyme immunoassay of *Cryptosporidium*-specific immunoglobulin G antibodies to assess longitudinal infection trends in six communities in British Columbia, Canada. *Am J Trop Med Hyg* 2005; 73: 288-295.
12. Eisenber JN, Priest JW, Lammie PJ, Colford JM Jr. The serologic response to *Cryptosporidium* in HIV-infected persons: implications for epidemiologic research. *Emerg Infect Dis* 2001; 7: 1004-1009.
13. Sandhu SK, Priest JW, Lammie PJ, Hubbard A, Colford JM Jr, Eisenberg JN. The natural history of antibody responses to *Cryptosporidium* parasites in men at high risk of HIV infection. *J Infect Dis* 2006; 194: 1428-1437.
14. Arrowood MJ, Sterling CR. Isolation of *Cryptosporidium* oocysts and sporozoites using discontinuous sucrose and isopycnic Percoll gradient. *J Parasitol* 1987; 73: 314-319.
15. Moss DM, Lammie PJ. Proliferative responsiveness of lymphocyte from *Cryptosporidium parvum*-exposed mice to two separate antigen fractions from oocysts. *Am J Trop Med Hyg* 1993; 49: 393-401.
16. Frost FJ, Tollestrup K, Craun GF, Fairley CK, Sinclair MI, Kunde TR. Protective immunity associated with a strong serological response to a *Cryptosporidium*-specific antigen group, in HIV-infected individuals. *J Infect Dis* 2005; 192: 618-621.
17. Nydam DV, Lindergard G, Guard CL, Schaaf SL, Wade SE, Mohammed HO. Serological detection of exposure to *Cryptosporidium parvum* in cattle by ELISA and its evaluation in relation to coprological tests. *Parasitol Res* 2002; 88: 797-803.
18. Chai JY, Lee SH, Guk SM, Lee SH. An epidemiological survey of *Cryptosporidium parvum* infection in randomly selected inhabitants of Seoul and Chollanam-do. *Korean J Parasitol* 1996; 34: 113-119.
19. Park JH, Kim HJ, Guk SM, Shin EH, Kim JL, Rim HJ, Lee SH, Chai JY. A survey of cryptosporidiosis among 2,541 residents of 25 coastal islands in Jeollanam-do (Province), Republic of Korea. *Korean J Parasitol* 2006; 44: 367-372.
20. Seo M, Huh S, Chai JY, Yu JR. An epidemiological survey on *Cryptosporidium parvum* infection of inhabitants in Chorwongun, Kangwon-do. *Korean J Parasitol* 2001; 39: 201-203.
21. Nichols GL. Food-borne protozoa. *Br Med Bull* 2000; 56: 209-235.