from patients who are taking salicylate of soda is not to be attributed to the presence of glucose in small quantity.

The Reducing Action of Chloral-Hydrate upon Cupric Oxide.—Dr. Sherwin (*Boston Med. Journ.*, November 18th, 1886, p. 487, and *Pharmaceutical Journal*, December 25th, 1886, p. 507) states that a sample of urine, specific gravity 1015, to which a few grains of chloral-hydrate had been added to prevent putrefaction, reduced alkaline copper solution like diabetic urine. He also found that a very weak solution of chloral-hydrate in water reduced Fehling's solution, and the urine of two patients who had taken a medicinal dose of chloral gave a similar reaction. I have repeated and confirmed Dr. Sherwin's observations, and I have further ascertained that, while chloral-hydrate has a powerful reducing action on the copper solution, it has none upon the picric acid test.

The Potassio-Mercuric Iodide Test for Albumen is Fallacious.—The Albumen Tests Committee of the Clinical Society, in their report, published in the last volume (xix) of the *Transactions*, state: "That a solution of potassio-mercuric iodide with citric acid, particularly when used after Heller's method, gives the most delicate and clearly marked reaction." It is very noteworthy that the committee make no mention of the fact which has often been recorded, and which must now be pretty generally known, that this test causes an opalescence, not to be distinguished from a slight albuminous precipitate, in every normal urine. This reagent apparently precipitates, not only mucin, but also some other ingredient of normal urine. For it is found that when the mucin in a normal specimen has been precipitated by citric or acetic acid, and then separated by filtration, the addition of the potassio-mercuric iodide to the clear urine causes a further precipitation and opalescence. It is obvious, therefore, that this reagent, if relied upon as a test for a minute trace of albumen, would be seriously misleading.

The Quantitative Estimate of Albumen.—Since attention has been drawn to the use of Esbach's graduated tubes by Dr. Veale (*BRITISH MEDICAL JOURNAL*, May, 1884), by Mr. Blomfield (*Lancet*, January 23rd, 1886), and by Dr. Sidney Coupland and myself (*Lancet*, July, 1886, p. 63), this simple and accurate method of ascertaining the actual amount of albumen has been coming into general use. Mr. Cetti, of 36, Brook Street, Holborn, who supplies the tubes, with printed directions for their use, at a small cost, tells me that there is a constant and increasing demand for them. The method combines simplicity and facility of working with accuracy of results in a quite remarkable degree.

The most convenient Pocket Test Case.—After trying a great variety of pocket test cases, I have found none so convenient as the "nickel-plated pocket urine test case," made by Ernst Jahneke, Canonbury Works, Dorset Street, Essex Road, N., and sold by most surgical instrument makers and pharmaceutical chemists. The case admits of some variety in its fittings and contents. In my own case, which, after two years' use, is as bright and clean as when it was new, I have two test tubes one within the other, and within the smaller tube a nipple pipette, which is very useful for transferring the urine to the test tube. Then there is a small spirit-lamp, the glass cap of which is so accurately fitted that no spirit leaks out, a stoppered bottle for powdered picric acid, another for grain pellets of caustic potash, and a drachm measure. There are also some blue and red litmus papers.

The case, therefore, contains all the materials required for bedside albumen and sugar testing. In testing for albumen, as much picric acid powder as can be carried on the end of a penknife (about a grain by weight) is dissolved in about a drachm of water by the aid of the lamp. To this is added an equal volume of urine, when the presence of albumen is at once made apparent. Then, to test for sugar, a grain pellet of potash is added to the mixed urine and picric acid, and the liquid is again boiled for a few seconds. If sugar is present, the liquid will be rendered so dark, that no red colour is visible through the test-tube. By diluting the urine in a definite proportion until the standard red colour is obtained, a proximate estimate of the proportion of sugar may be obtained, but, for an accurate analysis, the picro-saccharometer must be used.

The Albumen Tests Committee admit that the possibility of being able to test for both albumen and sugar, by picric acid, with the addition of potash, "renders this very portable test most useful;" but they add, "There is, however, some practical difficulty here arising out of the hygroscopic quality of the potash cubes recommended by Dr. Johnson." I can assure the committee, and all whom it may concern, that there is here no practical difficulty whatsoever. I have some potash pellets which were supplied to me six months ago by Mr. W. T. Cooper, of 66, Oxford Street, and they are as dry and hard now as when I first received them. Doubtless caustic potash exposed to the air will deliquesce, but, to prevent this, nothing more is re-

quired than to keep it in a bottle, the glass stopper of which is smeared with vaseline. This effectually excludes air and moisture, and also prevents the stopper from becoming fixed by the corrosive alkali.

PERIPHERAL NEURITIS IN RAYNAUD'S DISEASE (SYMMETRICAL GANGRENE).¹

By JOSEPH WIGLESWORTH, M.D. LOND.,
Assistant Medical Officer, Rainhill Asylum.

THIS communication had reference to a case of Raynaud's disease, under the observation of the author, in Rainhill Asylum, in which changes in the direction of chronic inflammation or sclerosis were found in the peripheral nerves after death.

The patient, a married woman, aged 26, was admitted from the workhouse December 27th, 1884, and died May 10th, 1886. She suffered from epilepsy, the monthly average of fits being about 2½. Her mental condition was one of mild dementia; but she occasionally had slight, transient, maniacal attacks, and exhibited at times hallucinations of hearing. Physically, she was a small, thin, badly-developed young woman, measuring 4 feet 10½ inches in height, and weighing only 79 pounds. There was a peculiar macular eruption scattered over face, trunk, and extremities, but most marked on the face, consisting of irregular blotches of a rather deep port-wine colour; these blotches faded completely under pressure, and were apparently caused by stasis in the capillaries; they underwent no change with time.

There was slight bronchitis; the heart was distinctly hypertrophied; and the urine contained permanently about one-fourth albumen. Both hands were very small and slender, the muscles were everywhere atrophied, and several fingers had undergone partial amputation. Thus, in the right hand, the thumb, first, and third fingers consisted of little more than the first phalanx, while the second finger consisted of this phalanx only, the fourth finger alone being entire. In the left hand, the second finger was reduced to the first phalanx, whilst the other fingers were entire, but contracted in flexion. The right foot had lost the big toe from amputation, and in the left foot the corresponding toe had undergone considerable loss of substance, and presented an unhealed sore. The patient herself stated, with reference to these losses of substance, that the parts mortified and were removed by a doctor—which was, without doubt, just what occurred; for, not only did the patient's mental condition permit of reliance being placed on her statements, but the appearance of the hands was quite characteristic.

Actual gangrene was observed whilst the patient was under observation. She suffered at irregular intervals from small excoriations, which formed chiefly on the tips of the fingers, or on the extensor aspects of the fingers and toes; at one time a superficial slough formed over the right olecranon, on the separation of which a sore was left about the size of a shilling. But the most marked example of gangrene was the formation of a slough on the extensor aspect of the second phalangeal joint of the left fourth finger; this rapidly spread upwards and downwards, opened into the second phalangeal joint, and necessitated amputation at the joint below. The patient's death occurred rather suddenly, after an epileptic fit, on May 10th, 1886. At the necropsy, thirteen hours after death, the brain weighed 1,176 grammes, and, with the exception of slight thickening of the membranes and slight wasting of the gyri, appeared quite normal. The lungs were moderately congested, and the heart (weight 9 ounces) showed considerable hypertrophy of the left ventricle. The liver and spleen were normal, but the kidneys (right 2½ ounces, left 2½ ounces) were diminished in size, with adherent capsules, granular surfaces, and mottled cortices, containing numerous cysts.

Raynaud described the disease as a neurosis, characterised by an exaggeration of the excito-motor centre in the cord, which presides over the vaso-motor nerves; and he attributed the local mischief to cramp-like contractions of the arterio-capillaries.

In the case above described, well-marked changes in the peripheral nerves were found on microscopical examination. The spinal cord and nerves were hardened in Müller's fluid and spirit; sections were made transversely, and stained with eosin and logwood. The spinal cord itself presented but slight changes. The nerve-cells of the anterior cornua were healthy, or nearly so, but those forming the posterior vesicular columns of Clarke presented more or less sphericity of outline. There was perhaps a little general thickening of the neuroglia. The nerves examined were the following: sciatic, internal popliteal,

¹ Abstract of a paper read at the Pathological Society, January 4th, 1887.

external popliteal, posterior tibial, median, and ulnar, on each side, and the right musculo-spiral—thirteen in all. The great majority of these nerves presented well-marked changes, and none of them were wholly unaffected. The changes found were summed up as overgrowth of the fibrous elements of the nerve, with atrophy and degeneration of the nervous elements. The left posterior tibial nerve was looked upon as the most diseased of the nerves of the lower extremities. This nerve, as a whole, was distinctly diminished in size, and the overgrowth of the fibrous framework was well marked; the hyperplasia affected all the fibrous elements—epineurium, perineurium, and endoneurium; the perineurium formed a series of richly nucleated concentric laminae, which in many cases encroached considerably upon the nerve-bundles; the nerve-tubules were more or less atrophied and degenerated. These changes were found, though slightly less advanced, in the other nerves of this extremity examined; and the corresponding nerves of the right lower extremity were affected in an exactly similar way, though in a slightly less marked degree. Of the nerves of the upper extremities, the left median was taken to be the most affected. The epi-, peri-, and endoneurium were all hypertrophied, but the chief characteristic of this section was the great widening of the perineurial lymph-spaces, caused by shrinkage of the nerve-bundles; these spaces were more or less filled with a loose spongy sparsely nucleated connective tissue, which was seen in many bundles to be connected with thickened bands of the endoneurium. The nerve-tubules were much atrophied and degenerated. This dilatation of the perineurial lymph-space, with loose connective tissue contents, was much more marked in this nerve than in any other, but most of the other nerves presented more or less evidence of it. Of all the nerves examined, the right musculo-spiral was the least affected, the changes here being only slight. On the whole, the changes described appeared to be a little more marked on the left side than the right. The disease also appeared to be more advanced at the distal than at the central ends of the nerves; and the conclusion was, therefore, hazarded that the disease had its starting-point at or near the periphery, and travelled upwards, but a sufficient number of sections was not made to prove this absolutely.

The question was raised as to the nature of the disease. Was it to be looked upon as a chronic interstitial inflammation, which had gradually invaded the nerve-bundles, and caused them to dwindle; or had we to deal with a primary degenerative atrophy of the nerve-tubules, with sequential hyperplasia of their fibrous framework? The evidence was not deemed sufficient to draw a positive conclusion; but the appearances presented by the left median nerve seemed rather to suggest a primary atrophy, for here the nerve-bundles had obviously shrunk out of proportion to the fibrous overgrowth, and the space thus left was occupied by a loose connective tissue, which might be looked upon as representing the earliest phase in the development of the concentric laminae of the thickened perineurium. There was, however, no reason to doubt that the localised symmetrical gangrene was directly dependent upon the changes in progress in the nerves; and should the above observation be confirmed by others, it would tend to take Raynaud's disease out of the category of neuroses by giving it a tangible material lesion to rest upon.

RUPTURE OF THE UTERUS DURING PREGNANCY,

WITH NOTES OF A CASE.¹

By ARTHUR H. N. LEWERS, M.D. LOND., M.R.C.P.,

Assistant Obstetric Physician to the London Hospital; Physician to Out-Patients, Queen Charlotte's Lying-in Hospital.

RUPTURE of the Uterus during Pregnancy stands in marked contrast to Rupture of the Uterus during Labour, partly because it is an accident of very much rarer occurrence, but especially because its causation is little understood.

In considering the subject, it is necessary to restrict the expression "rupture of the uterus during pregnancy" to cases where the fetus has not reached a considerable size, otherwise the causes producing rupture of the uterus during labour, causes that are well understood, come into operation. The causes of rupture during labour may briefly be described as follows: some obstruction exists opposing the advance of the child; whether the obstruction is pelvic contraction, unusual size of the child, or malpresentation does not matter: the uterus continuing to contract, thickening of the upper part of the uterus occurs, while its lower segment becomes thinned. If assistance is not given, the lower segment becomes more and more thinned, and finally ruptures. It is to be noted that rupture of the uterus during labour

almost always begins in the lower segment. No such explanation will help us to understand cases where rupture of the uterus occurs about the middle period of pregnancy, when the fetus is small, and when labour has not commenced, and where no mechanical violence has been used. Cases where the uterus ruptures during pregnancy, as the result of a severe blow, are not difficult to understand; clearly, the blow need only be hard enough, and sufficiently well-directed, to be in itself a satisfactory explanation of the accident. It is the residuum of cases where nothing of the kind has happened, where pregnancy is not far advanced, and where nothing unusual has been noticed up to the time of the rupture, that present a difficulty.

In Trask's monograph on Rupture of the Uterus, 303 cases are recorded; of these 38 only are classified as ruptures during pregnancy, the others all being cases of rupture during labour. On a careful examination of the notes in each of these 38 cases it appears that the number must be considerably reduced, first by taking away cases where the evidence that they were really cases of rupture of the uterus and not some other condition, for instance abdominal extra-uterine pregnancy, is unsatisfactory, and second, by removing those cases which were really cases of premature labour, the pregnancy having advanced nearly to term. When this reduction has been made there remains 16 cases where the evidence of rupture is fairly conclusive, 14 of spontaneous rupture, and two of rupture due to mechanical violence. Trask's monograph was published in 1847, and he appears to have collected all the cases he could find recorded from about the year 1700 to the time of writing. I have only been able to meet with records of two cases of spontaneous rupture, in addition to the 14 in Trask's paper—one by Lustgarten, quoted in the *British and Foreign Medico-Chirurgical Review*, and the other in the *Glasgow Medical Journal* for 1861. To these I will provisionally add my own case, making 17 in all. Though no doubt others have been recorded, and might have been found by a further search, still a group of 17 cases of a very rare condition may be taken as representative, and affording material from which some general conclusions on the nature of the accident may be drawn. I propose to read notes of two of the cases, first that from the *Glasgow Medical Journal* of 1861, and then the notes of my own case.

CASE I.—B. T., aged 37, unmarried, a farm-servant. The day before the accident, the patient had walked four miles and milked a cow, but there is nothing to suggest that she had undergone any unusual exertion. She went to bed, after a good supper, in good spirits, and apparently quite well. Soon after, about 11 P.M., she complained of pain in the stomach and bowels. When asked if she had walked too far, she answered that she had not. She vomited, and continued in pain till 7 the next morning, when she died. On *post mortem* examination, the abdomen was found to contain enough blood and clots to fill a washhand-basin. Projecting from the pelvis there was a large tumour, covered with clotted blood. This, on further examination, was found to be the uterus. A fetus of about the fourth month, covered by its proper membranes, was seen partially protruding through a rent in the fundus of the uterus. The rent was right across the fundus, and measured transversely $4\frac{1}{2}$ inches, and from before back $3\frac{1}{2}$ inches. The uterus at the seat of rupture appeared normal in structure, and at no part of it was there any appearance of disease. At the os uteri and in the neck of the uterus was found a plug of clear mucus, completely closing the cervical canal. It may be mentioned that no diagnosis was made during life; and that there was, till the *post mortem* examination, some suspicion that the patient had been poisoned.

The other case is that from which the specimen shown was taken and came under my care at the London Hospital.

CASE II.—A. D., age 33, had had six children, the last about a year and ten months before admission. At the time the accident happened she believed herself to be five months pregnant. She had not menstruated for five months. Nothing had occurred during the pregnancy different from what she had experienced on other occasions. On October 12th, about 3 P.M., while going upstairs, she was suddenly seized with severe pain in the abdomen, and became rapidly blanched and collapsed. When the pain seized her she felt as if something had given way in the abdomen, and stated that she heard a crack. Vomiting occurred soon after. She was seen by someone from a neighbouring dispensary, but after a time, the friends becoming alarmed, Dr. Wheeler Brown was called in. This was about 11 P.M. He at once recognised the gravity of the case, and brought the patient up to the London Hospital. She was admitted about one o'clock in the morning, and shortly afterwards Mr. Blaxland, the resident accoucheur sent for me. When I saw the case about a quarter past two the patient was much less collapsed than I was told she had been some time before. She was able to answer questions, and in fact sup-