be hard for the scientific pathologist to admit the cogency of these arguments, for he is tempted to assume that the epiblastic origin is true because it accords with *a priori* conceptions. For instance, in the case of the testicle, the assumption that the Wolffian duct arises from epiblast, has suggested a plausible explanation of the origin of testicular carcinomata; but under the circumstances such an inference seems premature. However clearly the epiblastic origin of the Wolffian duct is demonstrated, an obvious fallacy has to be guarded against. It is superfluous to point out that nothing but a very simple conclusion ought logically to be drawn from that demonstration; namely, that it may explain the origin of carcinomata from the parts developed from the Wolffian duct.

After describing the way in which the Wolffian duct attained its position in relation with the body cavity, the development of the Wolffian tubules was discussed. The simplest origin for these organs would be as outgrowths of the Wolffian duct, but in the rabbit they arise independently, and were produced from the mass of cell (intermediate cell-mass) from which the Wolffian duct grew. Anteriorly the rudimentary tubules are continuous with the peritoneal epithelium, but posteriorly they are separate (Fig. 3).

Fig. 4.—Front part of human Wolffian body, to show the appearances which may be supposed to indicate the existence of a fore-kidney. The embryo was at about the thirtieth day of intra-uterine life. C.V. Cardinal vein; P. Protrusions into peritoneal cavity; R. Recesses; W.T. Wolffian tubules.

Soon after their formation the tubules become continuous with the inner side of the duct, and eventually by a separation of the central cells acquired a lumen opening into the lumen of the Wolffian duct. The number of Wolffian tubules developed was equivalent to about two or three for each somite in relation with the Wolffian body.

After showing that in early human embryos the Wolffian tubules were continuous with the peritoneal epithelium, the question was raised whether the body cavity was, in mammals, prolonged into the Wolffian tubules. For this purpose, the anterior end of the rabbit's Wolffian duct was examined, and the following condition described (Fig. 3). After beginning as a solid column of cells, and after receiving two prolongations (nephrostomata) from the body-cavity, the duct ceased. After a few sections it began again, and had one nephrostoma, after which it merely received the Wolffian tubules.

The lecturer agreed with Janosik and Renson in considering that this indicated the existence of a head kidney or pronephros in the rabbit. The absence of external glomeruli, such as Renson mentions, was noted, but this fact was not thought to militate against the previous assumption.

After showing that nephrostomata seemed to be present in transverse sections of human embryos, the appearances seen at the fore end of the Wolffian body of a human embryo were described, in which there were three well marked protrusions from the fore end of the Wolffian body (Fig. 4), the recesses between these protrusions were in close relation with the Wolffian tubules. The lecturer thought that these appearances indicated the existence of a pronephros in the human embryo; and, in conclusion, drew attention to the various histological specimens and drawings with which the lecture was illustrated.

TINNITUS AURIUM, AND ITS TREATMENT BY A NEW METHOD OF ALTERNATE INJECTION AND EVACUATION OF AIR.¹

By JOHN WARD COUSINS, M.D. LOND., F.R.C.S.,
Senior Surgeon to the Royal Portsmouth Hospital, and to the Portsmouth and South Hants Eye and Ear Infirmary.

TINNITUS AURIUM is a symptom which occurs in connection with many obscure aural disorders, and it is also a frequent accompaniment of many of the common forms of middle-ear disease. It is generally absent, however, in cases of chronic otorrhœa, attended with perforation or ulceration of the membrana tympani.

In different cases we observe a marked variation in the character and intensity of the tinnitus. The noises are compared to many varieties of external sound, and their existence seems sometimes to augment the intensity and to modify the character of actual sounds; so that the distress of the patient is increased, and his head is constantly filled with vibrations which have no real existence in the surrounding air. Tinnitus is generally associated with some degree of deafness, but cases occasionally occur in practice in which the hearing power, tested by the ordinary method, appears to be scarcely below the normal standard. Sometimes the noise is referred to one or both ears; at other times it is described as sounding all over the head. Sometimes it is intermittent, and very variable in character. Sometimes it sets in gradually, and, at the onset, scarcely causes much annoyance; but it occasionally creates great alarm by the suddenness of its invasion. Fortunately tinnitus often passes away, and gradually subsides, and the patient is scarcely conscious of the time of its departure. On the other hand, it sometimes continues for years, and haunts many who are the unfortunate victims of the most hopeless forms of aural disease.

Tinnitus aurium can be artificially induced by simply pressing the tragus firmly over the aural orifice. In this way the column of air within the external passage is condensed, and the pressure on the drum augmented, and this increase of pressure is directly transmitted through the conducting apparatus to the endo-lymph and nervous structures of the labyrinth. Now this little experiment yields an explanation of the occurrence of tinnitus in association with the most common forms of middle-ear disease; for, in a vast majority of cases, it is wholly dependent upon abnormal tension within the tympanum. In acute catarrhal affections it is present, together with throbbing and distressing pain. An attack of this kind often follows exposure to cold, and the introduction of water into the ear during bathing. The hearing power is impaired, the lining membrane of the middle ear congested and swollen, and its cavity distended with mucous secretion. This sort of aural disturbance originates, too, in pharyngeal and bronchial affections, and generally soon passes off; but it is also the forerunner of a chronic disorder, by which the conducting apparatus is seriously damaged.

Again, in chronic catarrhal inflammation of the tympanum, the cavity becomes filled with thick and inspissated secretion, its mucous membrane gets thickened, and bands of adhesion slowly form around the ossicles; so that the membrana tympani loses its normal elasticity, the mobility of the bony chain is impaired, and tympanic tension is developed, attended with gradually increasing deafness and tinnitus. In many of these chronic cases, a disturbance of pressure is produced by obstruction of the Eustachian tubes, and this is of frequent occurrence as the result of catarrhal and specific inflammations of the

¹ Address delivered before the South-East Hants Medical Society.

throat. In the early stage, the pent-up fluids cause serious distension, but these become gradually absorbed; and then, unless the normal communication between the ear and the pharynx is re-established, the atmospheric pressure on the outer surface of the membrana tympani presses it inwards, and forces it against the fenestra ovalis. In this way, the organ undergoes permanent injury, and the drum, together with the contents of the tympanum, are damaged by structural changes. The membrane then appears to be deeply indented and concave, the margin looks thickened, the centre thin and transparent, and sometimes the short process of the malleus is visible through it.

Deafness and tinnitus are also intimately associated with diseases of the naso-pharynx. The tonsils are often enlarged, and their follicles filled with a yellowish fluid. Sometimes the mucous surfaces of the pharynx are swollen and covered with a thick muco-purulent secretion, which completely blocks the Eustachian orifices. In many chronic cases, the mechanical closure of the tubes is caused by out-growths of glandular tissue, which narrow the naso-pharynx, and obstruct the nasal passages.

Pressure on the outer surface of the drum is another source of tinnitus. It is frequently excited by the presence of a mass of impacted cerumen or the lodgment of a foreign body. In a case, recently under my care, immediate relief followed the removal of a large polypoid growth which projected from the external orifice. Under these circumstances, of course, efficient treatment is very easy of application.

Some aurists have ascribed the obstinate continuance of this unpleasant symptom to an altered condition of the intrinsic muscles of the tympanum. It is quite possible that, during a state of chronic inflammation, they may form adhesions with the surrounding structures, and so excite abnormal pressure during their contraction. On the other hand, they may lose their normal tonicity, or fall into a paralytic condition, through vascular and trophic changes. Now, any loss of contractile power must tend to aggravate any disturbance of the balance of pressure within the tympanum. The muscles of the ossicles are directly stimulated by sound, and, in the normal state of the organ, may regulate the tension of the membrana tympani, and the movements of the stapes in the foramen ovale; so it is at least probable that a loss of tone may be followed by an undue excitation of the auditory nerves.

I have now briefly considered tinnitus aurium in its relation to many common disorders of the conducting apparatus of the ear. In all cases of middle-ear disease in which it is present, it must be regarded as a symptom excited by abnormal pressure upon the internal auditory apparatus, and produced by mechanical causes within the tympanic cavity. But this disorder has sometimes a much deeper origin, and is essentially dependent upon disease of the nervous structures of the organ, and this, too, under many and very varying conditions.

During the course of brain or meningeal disease, the blood-supply to the internal ear may become deranged, and thus abnormal vibrations may be generated by a disturbance of the circulation within the cranium. The presence of persistent tinnitus, in association with general indications of atheromatous degeneration, must be regarded as a premonitory symptom of structural changes in the walls of the cerebral blood-vessels. Under such circumstances, the sounds appear especially depressing to the patient, exhibiting a wavy or beating character, and are often synchronous with the cardiac systole. Not long ago, a case of this sort occurred in my practice, and the patient, an officer in the army, died suddenly in a London street.

Tinnitus may result from hyperæsthesia or undue irritability of the auditory nerve. This condition is often induced by constant exposure to loud and harsh sounds. Many persons engaged in noisy trades experience for hours, after the cessation of their toil, a constant buzzing in the ears. Now, whenever this aural system constantly recurs, more or less deafness is sure to follow; and, if the exposure is continued, it will at length issue in the development of a very obstinate and persistent form of this affection.

In many cases of chronic aural disease, the branches of the auditory nerve which are distributed to the deep recesses of the labyrinth, appear to be the real seat of the disorder. Sometimes the fibrous membrane lining the bony cavities, and also the osseous structure itself, are found thickened and congested. The blood-vessels of the labyrinth are likewise dilated, and the membranes marked by the remnants of old capillary extravasations. Many forms of atrophy and degeneration of the nerves of the labyrinth are the result of arrested function, induced by disease of the conducting apparatus and tympanum. In these cases, from the long withdrawal of healthy stimulation, the ultimate fibres have lost their sensibility for sound, and a persistent tinnitus has been roused into existence dependent upon chronic changes

in the walls of the membranous labyrinth, which are capable of exciting the filaments of the auditory nerve, and creating disorderly discharges in the direction of the cerebral centres.

The most careful examination of the internal ear, in cases of obstinate aural disease, has sometimes failed to discover any very definite structural lesion; but, after all, opportunities for investigation in this direction are rare; at the same time, the complexity and delicacy of the auditory nervous network render the recognition of morbid changes a matter of considerable difficulty.

I have already remarked that very slight and varying conditions of the tympanic apparatus are often quite sufficient to modify the sensibility of the ear, and to kindle within it disturbing vibrations. Now, in some cases, these little variations are rendered acutely distressing through constitutional derangement, or by general disorder of the vascular system. Thus, tinnitus is sometimes present in anæmia and chlorosis, and is roused into activity and greatly intensified through the altered composition of the blood itself. At other times, it follows as the result of serious hæmorrhage, by which the blood-pressure is reduced and the vessels suddenly relaxed. It occurs in connection with venous hyperæmia, associated with cardiac obstruction and regurgitation. Occasionally it is a troublesome symptom during the course of some forms of chronic disease, in which the aural vascular system is disturbed by the high tension of the circulation. Many constitutional disorders are also observed to be important factors in the production of tinnitus, and this fact is frequently exhibited in patients who have long been the victims of gout, alcoholism, chronic dyspepsia, and general nervous depression.

It is evident, then, that tinnitus, of every character and intensity, must be regarded as the outcome of many disordered conditions of the ear, and that these conditions are aggravated by vascular and constitutional derangements. Sometimes it is an indication of temporary functional disturbance, at other times it points to structural change in some part of the organ. When considered apart from all its surroundings, it possesses very little diagnostic value. The important question to determine is the special form of aural disease of which it is a manifestation. Is it a symptom of middle-ear disease, or an affection of the nervous structures of the organ? We must not forget, however, that in practice the deep and superficial disorders are often intimately blended together, that changes in the sound-receiving apparatus are the secondary results of chronic middle-ear disease, and that it is sometimes a difficult problem to determine the exact seat and nature of the lesion which is the real source of the tinnitus. It is necessary, therefore, that we should in every case investigate the history and duration of the disorder, and carefully inquire into the constitutional condition of the patient. The conducting power of the ear must be tested with the watch and tuning-fork, and the external meatus and tympanic membrane inspected under good illumination. It is always, too, an important matter to examine the condition of the pharynx and nasal cavities, and then to prove the mobility of the drum, and the patency of the Eustachian tubes, by means of the air-douche and otoscope.

[To be continued.]

PROLAPSE OF THE UPPER INTO THE LOWER PART OF THE RECTUM WITHOUT EXTERNAL PROTRUSION.

By HARRISON CRIPPS, F.R.C.S.,

Assistant Surgeon to St. Bartholomew's Hospital.

THE term prolapse of the rectum is often used somewhat loosely, many writers making it include hæmorrhoidal protrusion as well as the simple eversion of the gut. When the protrusion chiefly consists of dilated or hypertrophied vessels, it is rightly described as hæmorrhoidal, notwithstanding that a certain amount of normal mucous membrane may be drawn down at the same time.

The word prolapse should be restricted to cases in which the bowel, except for its liability to protrude, is healthy, or, at any rate in which any morbid condition affecting it is the result, rather than the cause, of the protrusion.

The amount of bowel that prolapses varies greatly, sometimes being little more than a fold of mucous membrane, while, occasionally, many inches involving all the coats of the bowel are turned inside out. In the former case the prolapse is described as partial, in the latter as complete. The closeness of the attachment of the rectum to the surrounding structures varies greatly in different individuals, a fact often noticeable upon examining the part with the finger. In some patients the bowel, when empty, has a smooth uniform surface, while