the body into a complicated attitude of defence? This introduces far too wide a subject of discussion to take up at present; but it seems to me extremely likely that the thalamus is an educating optic ganglion, in the sense that the callosal fibres entering it have their functions concentrated within its substance, and that these fibres are the means of educating the opposite side of the brain through vision.

Where the corpus callosum has been destroyed in infancy, the invariable result seems to have been imbecility. There are certain congenital deficiencies of the body which have been recorded unaccompanied by any symptoms of note, more especially one described by Eichler. In these latter cases, however, it seems, from an examination of their records, that we have to do not with a deficiency in the actual callosal fibres, but with a malformation by which they have failed to decussate in the middle line, just as so frequently happens in the anterior pyramids.

On the supposition that the thalamus subserves the purpose of concentrating the fibres which educate the higher centres through the optic, I can quite easily see how, if it were destroyed in adult life, no very evident symptoms might follow. It has already subserved its purpose to a great extent. The higher centres have been educated, and are capable of discharging their functions apart from the channels through which that education has been imparted.

Nay, I could go so far as to grant that a large part even of the corpus callosum might be annihilated in adult life without occasioning any very marked phenomena. It has already played its part, so to speak, in infancy and youth, and may in a manner be now considered as functionally inert. The impressions made upon the cerebral cortex through it are quite possibly recalled by a perfectly different set of channels; for I do not see why in the brain-economy there may not be paths for educating the higher centres through vision, hearing, touch, and so on, and a whole set of other paths by which the results of this education may be brought into action. If such be the case—and I advance this simply as a surmise—I should regard the callosal system of fibres as the great educating system; while the other direct bundles to which I have adverted would constitute the means of adapting this education to a utilitarian purpose.

In the case of those born blind, this education, of course, could not go on through the optic circle of communications, but would take place by means of the nuclei of the nerves of special sense situated in the pons and medulla and the crossed callosal fibres connected with these.

AN ADDRESS

ON

SOME OF THE RARER SYMPTOMS PRODUCED BY GALL-STONES.

Read in the Section of Medicine at the Annual Meeting of the British Medical Association in Brighton.

By WILLIAM M. ORD, M.D. LOND., F.R.C.P.,
Physician to St. Thomas's Hospital

THE occurrence of gall-stones is so common, and the symptoms of biliary colic are, in a general way, so decisive, that, in dealing with the effects produced by gall-stones, I shall ask to be allowed to pass over many points which, in a complete treatise, would be, necessarily, subjects of primary consideration. Therefore, I do not intend to discourse either on the structure of gall-stones, or on the way in which they come to be formed; nor to describe biliary colic as commonly seen; nor to enumerate all the possible wanderings of gall-stones, either to exit by one path or another from the body, or to intrusion into various organs; nor to discuss treatment, operative or not operative, seeing that this relation is to be discussed in another Section.

I will ask you to follow me in two lines of thought. Accuracy of diagnosis occupies, in all cases, the primary attention of the practical physician. In this aspect, it appears to me, on the one hand, that cases in which gall-stones have existed without producing the symptoms commonly indicative of their presence may be found worthy of consideration; that, on the other hand, cases illustrating uncommon effects due to the presence of gall-stones may be found to have a value, a very different value, of their own. Around this thread of diagnostic investigation winds the thread of

inductive or explanatory investigation, every turn whereof may send us to contemplate some fresh and important relation to the whole organism of the small area affected by disease. In this aspect we are rather philosophic observers than practical physicians. If we are to do all that is required of us, we must be both; and we may be sure that it will ever come about that in this duality each comrade will minister to the other. I will first call your attention to gall-stones which give rise to no symptoms.

Gall-stones Existing without Symptoms.—It is a common experience of the *post-mortem* room that gall-stones, small and large, solitary or in hundreds, shall be found in the bodies of persons whose medical history contains no record of suffering or disease such as our present knowledge would refer to gall-stone; and that, structurally, no evidence of serious local mischief determined by the presence of concretions shall be apparent. Are there, one is tempted to ask, people who pass gall-stones unconsciously, and without symptoms, as there are people who pass renal or vesical calculi, without other signs than the migrations of the calculi into the chamber-pot?

Gall-stones Passed without Symptoms.—The occurrence of what is generally spoken of as biliary colic, with vomiting, with sequence of jaundice, and of the faecal and urinary signs of retention of the bile, constitute, I think you will allow, the most obvious signs of the existence of gall-stone. But gall-stone of large size may exist, and may find its way into the bowel, without producing any symptoms whatsoever of biliary colic. Here is a specimen which I showed some time ago at the Pathological Society of London. It was sent to me by my friend, Mr. Carr H. Roberts, of Kensal Town. I always call it the "Twin," because it was passed by a lady on the day after her confinement, with such difficulty as almost to constitute a second labour. This lady had, so far as I can learn, no precedent signs of gall-stone whatever, and it must be supposed that adhesion had taken place between the gall-bladder and an adjoining coil of intestine, and that a painless ulceration had offered an easy retreat to the concretion. But we all know that the obstruction met with in this case at the anus may occur in the intestine. Most of us have, no doubt, seen cases of the kind. I have seen several. One, which I saw with Mr. Bullock, of Isleworth, I should like to quote. A lady, aged between 60 or 70, was attacked suddenly with symptoms of intestinal obstruction. A careful investigation led to the belief that the obstruction was mechanical, and within the abdomen. Mr. Thomas Smith, of St. Bartholomew's, was accordingly asked to give his help. He first opened the abdomen at a point where most pain was felt. He then minutely examined the state of the intestines, bringing out through the small orifice the nearest coil of intestine, and working from this in both directions, a fresh coil being brought out after its predecessor had been examined and returned. A piece of jejunum, about 5 inches long, was at length extruded, in a state of intense inflammation, black and rough on the surface. Nothing abnormal could be felt within it, and we supposed that it had undergone some strangulation, which had been relieved by the operation. The patient died the next day, and a very limited *post-mortem* examination was allowed. A large gall-stone was, however, found, embraced strongly by slightly inflamed intestine, a little below the seat of inflammation already observed. The gall-stone had, of course, been moved on, without being felt, in the manipulation. Now, in this case, reviewed after the knowledge of the actual existence of the conditions stated, it may be observed that no ordinary symptoms of gall-stone had presented themselves. It follows that in considering the possible causes of any case of intestinal obstruction, we are not justified in excluding impaction of gall-stone, even though all story of previous biliary colic or of jaundice be wanting.

Gall-stones giving Rise to Pain, etc., without Jaundice.—The case just recorded leads on to the consideration of a fact of great importance, namely, that gall-stones may give rise to various kinds of pain and suffering to inflammation, and even to death, without the occurrence at any time of jaundice. Such cases are far from rare, and are to me of great interest. Let me quote two or three illustrative histories.

Some years ago, I was asked to see the wife of a clergyman in the East of London. I was informed that she was subject to irregularly recurring attacks of fearful pain in the region of the gall-bladder, with associated vomiting and faintness. She had never had jaundice, nor passed pale stools. Many physicians had seen her, whose diagnoses had ranged between uterine irritation at one pole, and cancer at the other. The absence of jaundice, and of the other signs of obstruction of the common bile duct, left the clinching of the diagnosis impossible, and the idea of duodenal ulcer was chiefly entertained. The patient died in an attack a few months later, and was found to have had a large biliary calculus, which had made its way through a perforation of the gall-bladder into the peritoneum.

Medicine, we may observe, is full of seeming paradoxes. One such paradox is here presented to us, in that, granted a disease having generally definite symptoms, death or danger occur frequently in such disease without previous presence of—rather with conspicuous absence of—the symptoms held to be most decisive of its presence. Let me, in passing, commend this seeming paradox for your consideration.

In a second case belonging to this class, the patient is an unmarried lady of 40, who is still under my observation. During the last three years she has had, from time to time, attacks of severe pain in the epigastrium, not related with food-taking, lasting for several days, almost continuously, but with recurring periods of intensity in which she has become faint. Vomiting has soon followed the pain, and has recurred at intervals, the vomited matter containing, besides ingesta, bile and mucus. There has been no jaundice; the fæces have been always of natural colour; there has been no pile-pigment in the urine. When I first saw her, a year ago, she was supposed to be suffering from new growth in the stomach. Her appearance did not support this idea. Although she had lost a good deal of flesh, she was still well nourished; her face was placid, and was free from all expression of anxiety; the tint of her skin was healthy, its texture soft and elastic. I saw her then in the interval of comparative ease following an attack. She still had uneasiness and distinct tenderness in the epigastrium. The abdominal wall was tense, from the xiphoid cartilage to the umbilicus, and through the tension it was possible to feel a somewhat extensive sensitive induration, corresponding in position to the right end of the stomach, the duodenum, and the gall-bladder. No special enlargement of the gall-bladder was made out, though, of course, carefully sought. The symptoms of gastric catarrh were present. Under treatment, chiefly directed to the relief of the gastric catarrh, she improved for some time; but, some months later, I was summoned to see her in consultation with Dr. Brockwell, of Gipsy Hill. She was then in the midst of one of the severe attacks of pain which I have described; there was again no jaundice or other sign of biliary obstruction. Two or three days later she passed, with a soft motion, a hard body, which attracted notice by the noise which it made in falling into the closet-pan. This hard body was broken up by her nurse. Fragments of it, subsequently brought to me, were clearly fragments of a gall-stone, and, as far as could be made out, of a gall-stone of considerable size. The passage of the gall-stone was followed by cessation of the urgent symptoms, and by rapid improvement in the general health. Within the last few weeks another attack of the same kind has ended in the passing of the stone which I now show. It is, as you see, a large stone, and presents at least five facets, indicating probably that it has had, at least, five companions of much the same size.

To these cases I will add a brief note of a third, which was under my care in St. Thomas's Hospital. In this case, after symptoms simulating cancer, but without jaundice, a gall-stone, escaping by ulceration into the duodenum, produced fatal obstruction in the upper part of the jejunum.

You will observe that, in three of the cases quoted, large gall-stones found their way into the intestine without the occurrence of jaundice or other sign of biliary obstruction.

The disproportion between the size of the stones and the size of the gall-duct renders it to the last degree improbable that the entry into the intestine was effected through the duct. The last case indicates the track probably taken by all—through an adventitious ulcerative perforation, leading, in the first case, into the peritoneum, in the rest, into the intestine, after the previous establishment of protective adhesions. In two cases, the process was attended with great suffering, and with signs of inflammatory thickening so marked as to suggest the existence of new growth. In the remaining case, which is far from standing alone in my experience, the process of ulceration and escape was attended by no sign, the obstruction being the first condition to attract notice.

These cases, and the recorded experience of many good observers, must lead us to remember that, although no signs of biliary obstruction be present, gall-stones must be kept in mind as possible factors in cases of recurring pain, vomiting, tenderness, and thickening in the epigastric region; that is to say, where cancer, ulceration of the pyloric part of the stomach, or of the duodenum, may be also suspected.

The class of cases in which no suffering is recorded, although large calculi have escaped from the gall bladder, is certainly puzzling. The passage of the stone into the intestine must have been permitted by processes of ulceration of the gall-bladder, of adhesions of the gall-bladder to the intestine, and by subsequent perforation of both parietes. Yet the several histories tell us of no symptoms of the occurrence of

such processes. Here is another paradox less easy to solve than the former.

In our considerations, we have, so far, been fixing our attention mainly on the nomadic gall-stone. We might, if time should have served, have followed it in many an excursion. A reference to text-books and monographs will readily indicate a great number of tracks by which a gall-stone, once escaped from the gall-bladder, has been known to travel to exit from the body, or to intrusion, mostly fatal, into viscera or into serous cavities. These tracks are so fully set forth in many books, that I do not propose to deal with them now. I leave them for the (to me) more interesting study of little noticed and more remote evils wrought by gall-stones.

Gall-Stones producing Intermitting Pyrexia.—First, let me speak of the occurrence in relation with gall-stones of a form of fever, with precedent rigors, occurring at more or less irregular intervals, so as to present some resemblance to an ague, but to an ague not of ordinary character. My attention was first called to this complication by some remarks of the late Dr. Murchison, having reference to the case of a distinguished Indian medical officer, who, after his return to this country, was attacked with paroxysms of shivering, followed by fever and sweating, at regular weekly periods. He was supposed, at first, to have a recurrence of an old intermittent; later on, to have hepatic abscess; till, at last, his symptoms indicated, and the necropsy proved, that his actual and only disease was gall-stone, so impacted as to produce great irritation, but not complete obstruction of the common bile-duct. Similar cases have been noticed by M. Charcot (*Maladies du Foie*, 1877), who argues that “la fièvre hépatalgique, comme celle de l’angiocholite, résulterait de l’introduction dans le sang de l’agent pyrérogène hypothétique provenant de l’altération de la bile.” Dr. Murchison was of opinion that such attacks were not of poisonous or septic origin, but were due to nervous irritation; and he compared them to similar intermittent febrile seizures observed in connection with the passage of renal calculi. Charcot has touched upon these also, and has evidently accepted in them the idea of a uro-septic fever. For my part, I believe that the paroxysmal fever here considered is due to local irritations of mucous membrane, propagated to the central nervous system, and resulting in pyrexia, mostly in persons apt to take on febrility, and particularly in persons who have been previously the victims of intermittents.

The whole subject of the tendency of irritation of mucous membranes to produce remittent or paroxysmal pyrexia is one which has yet to be studied completely; but, as a clinical observer, I believe that I can recognise such tendency, not only in enteric fever, but in catarrhs of all kinds, specially in children as regards mucous surfaces generally, specially in persons of all ages in as regards catarrhs of intestine, or in catarrhs of the urethra, particularly when the prostatic portion thereof is involved.

Gall-stones producing Glycosuria.—In relation to this question, there is much to be taught by the case of a gentleman of the age of 60, who first consulted me in December, 1884. He had lived many years in Ceylon, where he had severe jungle-fever. After returning to England—after, indeed, some time of residence at Bedford—he had become subject to regularly recurring attacks of biliary colic, with jaundice, and with “ague-fits” of great sharpness, coinciding with the biliary attacks. The liver and the spleen were found to be enlarged. He was very thirsty at all times, was passing much urine, and he was wasting. The urine at this time ranged in quantity from fifty to seventy ounces daily; its specific gravity ranged from 1026 to 1032; it contained much sugar, and much albumen. The case, after exclusion of all other conditions productive of febrility, seemed to me, so far as the intermittent febrility was concerned, to be like Dr. Murchison’s case; to illustrate the power of a limited, but intense, irritation of a mucous-membrane-lined duct, to arouse the febrile excitement to which the system had been accustomed in the tropics. But the glycosuria was a complication which had to be examined, both in diagnosis and in respect to treatment. I set to work to reduce, as far as possible, all catarrhal conditions, at the same time enforcing a diet intended to contain as little sugar-making stuff as possible. I found that, up to the time when I saw him, he had been taking large doses of quinine in alternation with arsenic, but had, notwithstanding, steadily gone down. On the hypothesis that the mucous membrane irritation was the real basis of his illness, I gave him alkaline sulphates, carbonates, and iodide of potassium. Before the action of the treatment could be well established, the patient had fallen in weight from an original 13 stone to 10 stone. Within two months from the institution of treatment, the liver and spleen had returned to normal size, the jaundice and glycosuria had disappeared, and the patient had increased to 12 stone 12 lbs. in weight. The improvement had commenced soon after the beginning of treatment, and was soon followed by the passage

of a gall-stone. It is important to notice that two attacks of "fever" had occurred without signs of colic. This patient subsequently gained flesh, till he reached the weight of 14 stone. I have just heard of his death from an attack of acute illness, the nature of which has not been communicated to me.

I have published, in *St. Thomas's Hospital Reports* of 1881, the case of a man who had attacks of fever of agueish character, occurring at first pretty regularly in conjunction with similarly regular attacks of biliary colic, and ceasing after the colic had yielded to treatment comprehending the use of no antiperiodic or antipyretic, but consisting in the free use of salines. In this case the symptoms left little reason for doubting the existence of biliary calculus, and of its causal relation to the attacks.

Reviewing this class of cases, I may point out that the glycosuria, increased quantity of urine, and thirst which formed no inconsiderable part of the symptoms of the case last but one quoted, disappeared when the febrility ceased to reappear. No one, I think, would find any ground for regarding glycosuria and polyuria, as the work of septicity or poisoning. Nevertheless, they coincided in their duration with a period in which constantly recurring block of the common bile duct was indicated, in which a distinct enlargement and painful condition of the liver was observed. I would venture to regard them as signs of a quasi-inflammatory condition of the liver, in which an active arterial hyperæmia was induced by the irritation in the gall-ducts, and by the incidental over-distension of the whole duct-system of the liver. They appear to me to illustrate the production of glycosuria by irritations of the liver. Glycosuria, as we know, may be brought about by irritation of the central nervous system, by paralysis of certain parts of the sympathetic system, by certain peripheral irritations outside the liver. In this case, the irritation of the liver has not been reflected to other organs, as is often the case, but has been turned back upon itself, with resulting arterial hyperæmia of the liver and glycosuria.

A case still under my observation is akin to the above. A labourer, aged 74, was admitted to St. Thomas's Hospital, for jaundice, which had commenced, suddenly, about six months before, after an attack which, as described, was an attack of biliary colic. From the time of that attack, there had been persistent jaundice, with equally persistent absence of bile in the fæces. It is necessary to state that this patient had several forms of disease. He had aortic obstruction, thickened arteries, enlarged heart, hypertrophy of the left ventricle, and albuminuria.

The analysis of the meaning of particular symptoms is certainly made more difficult by these complications. But his history makes it quite clear that soon after the establishment of jaundice, he began to pass a much larger quantity of water than he had been accustomed to pass, and that, having been a stout man, he began to lose flesh. When he came under observation, he was still fairly well nourished; his liver was greatly enlarged; but there was absence of pain, of tenderness, and of vomiting. He was passing from 60 to 80 ounces of urine daily; the specific gravity was generally about 1030, and the urine contained sugar, as well as some albumen. The history begins with biliary colic, and goes on to polyuria, with thirst and wasting, which proved, when observation began, to be associated with glycosuria. None of the other morbid conditions were such as would produce glycosuria.

Surely here again we have evidence of the reflection upon the liver of an irritation arising in its own domain—to all appearance an irritation started by gall-stone.

The argument of the preceding remarks is that irritation in the gall-duct may, besides producing pain and local inflammation, act in a reflex way, so as to produce arterial hyperæmia of the liver, and thereby glycosuria; or in another—possibly reflex—way, to produce a pyrexia corresponding with storms of local irritation. Have we, it is well to ask, any other indications of reflex action taking different directions from these? There is clearly a curious association of renal troubles with biliary colic, which has been noticed by Murchison and others. But I should like to give a short history of a case where acute pneumonia was, to all appearance, set up by impacted gall-stone, altogether apart from any sign of septic poisoning.

Gall-stones and Reflected Visceral Inflammations.—A gentleman, of middle age, had consulted me for some years on account of recurring attacks of what appeared, from all the symptoms, to be biliary colic. He was kept under very careful observation, but no calculus was ever found in the fæces. After two courses at Carlsbad, he seemed much better, and remained free from attacks for a year. Then he had a series of excessively severe attacks of biliary colic, increasing in duration. I went down into the country to see him in consultation with my friend, Dr. Horley, of Hoddesden, during one of these attacks,

with a view to coming to a decision as to cholecystotomy. Up to the time at which I saw him, he had no signs beyond those belonging to biliary colic; he had no irregular fever, no perspiration, no shortness of breath, but he had had some rise of temperature and some quickening of respiration for the last twenty-four hours. When I saw him, he presented, besides intense jaundice and the other results of obstruction, a pneumonia occupying the lower lobe of the left lung. At first I thought that this pneumonia must have arisen by continuity, and I feared that it might indicate either the formation of abscess in the gall-bladder, taking an unusual direction, or extension of inflammation; but no signs of suppuration, or of inflammation or tenderness in the epigastric or right hypochondriac region, were present. Moreover, there was no pain attending the contraction of the diaphragm. In respect of external causes, it may be stated that the patient had been lying in one room, which was well warmed and free from draughts, for several weeks, and had been sedulously protected from all exposure. On the other hand, it must be noted that the attacks of colic had been of increasing severity, and that the last, which had preceded the pneumonia by only two days, had produced intense shock. The pneumonia then appeared to me to be of reflex causation, and to be an alternative of the glycosuria, indicating hepatic hyperæmia, observed in the preceding cases.

Now we all know that, in advanced diabetes, pneumonia is a fairly common complication, and that when pneumonia occurs in diabetes, the sugar in the urine diminishes, or altogether vanishes. The disappearance of the sugar is often attributed to the fever accompanying the pneumonia, and is then regarded as an indication of increase of combustion in the body, whereby the sugar is consumed before it can reach the urine. I believe this hypothesis to be as far from the truth as the original hypothesis of Bernard assuming the combustion of sugar in the lungs—the hypothesis so completely demolished by Dr. Pavy. As I read the order of phenomena in these cases, the pneumonia is induced by a transference of arterial hyperæmia from liver to lung, variously determined by position of body, by cold, by disorders of venous circulation, by previous weakening of the lung. I have record now of three cases of diabetes complicated by pneumonia, in which fever preceded the appearance of the physical signs of pneumonia for twenty-four hours or more. In these cases, so long as there was only pyrexia, the sugar remained at its previous amount; when the signs of pneumonia became manifest, the sugar at once fell in two of the cases, and disappeared in the third. I venture to submit the probability that, in the case under consideration, the intense conditions of irritation in the liver tended, *per se*, to produce a primary reflex arterial hyperæmia, and, therefore, glycosuria; this was carefully looked for, but was not found. The absence of glycosuria was compensated by the access of pneumonia, presumably, in the circumstances of the cases, the result of a secondary or remote reflex arterial hyperæmia, determined by local or general conditions which cannot be identified.

It is well to note that the patient got over his pneumonia, which, of course, precluded operation; and, soon after, passed a large number of rather small, faceted, gall-stones, of which nearly seventy were obtained by regular washing of the fæces.

Gall-stones and Malignant Disease of Liver.—The co-existence of gall-stone with malignant disease of the gall-bladder and parts immediately adjoining has been recorded frequently enough to give rise to speculation as to how far the presence of gall-stones would be capable of giving rise to malignant disease. I exhibit a specimen from the museum of St. Thomas's Hospital, which has been described by Dr. Murchison. The patient, a woman, aged 64, had suffered for four years from attacks of severe pain in the region of the gall-bladder, without vomiting or jaundice. Six months before her death an unusually severe attack of pain was followed by jaundice, which persisted till her death. The stools were at first pale, but during her stay in hospital, for seven weeks before death, they contained a distinct evidence of bile. At the necropsy the gall-bladder and common duct were found filled with colourless (nearly clear) mucus, and the duct was blocked with a small calculus, at a point about an inch from its termination. A large cancerous mass occupied the centre of the liver, extending forwards to the neighbourhood of the gall-bladder. The bile-ducts were dilated throughout the liver, and filled with thick bile. The anterior wall of the stomach was adherent to the liver; and at the adherent portion were three or four rounded orifices, through which dilated bile-ducts on the surface of the liver opened into the cavity of the stomach. This was the source of the bile found in the fæces.

There is nothing in the case to indicate whether, on the one hand, the gall-stone was an incident to the malignant growth in the liver, or a result itself of retention of bile in the gall-bladder. Here, however, is another gall-stone which effectually answered my question,

in a case of malignant disease evidently beginning in and around the gall-bladder, and extending into the liver. I will not enter fully into the details of the case, but will merely say that the patient had violent recurring pain in the region of the gall-bladder, and for some time before death there was jaundice. At the necropsy the gall-bladder was found embedded in a mass of adherent growth, which completely closed the hepatic and cystic ducts. The cavity was rather small, and contained clear, pale, mucus, together with this calculus. It is so clear from the dissection that the growth had started at the gall-bladder, that I did ask myself the question whether the presence of the gall-stone might have been the original determining condition.

Chemical examination showed that the calculus was composed of carbonate and phosphate of lime, with a large proportion of animal basis, apparently altered mucus. It contained no trace whatever of any constituent of bile. This result showed, clearly, that the calculus was formed subsequently to the obliteration of the duct, and that it was a consequence, and not a cause, of the malignant disease.

Gall-stones and Hemorrhage.—In two cases coming under my observation of late, the passage of gall-stone has been attended by sharp hemorrhage. In the first place, the hemorrhage preceded the passage of a large gall-stone without biliary obstruction. The bleeding might here be supposed to be produced by the tearing of the opening between the gall-duct and the bowel.

In the second case which was under my care in St. Thomas's Hospital last spring, hemorrhage of a considerable volume occurred directly after an attack of biliary colic, with jaundice. After the cessation of the hemorrhage, a ragged gall-stone, of such size as might have allowed it to traverse the gall-duct, was found in the fæces. Another was found a few days later, after a smaller hemorrhage.

The various effects of gall-stones which we have passed under review may well make us pause and meditate. We have to deal with biliary colic very often; to know it as a process involving intense pain, and many other distresses; to see it make its progress commonly to a satisfactory end. As regards the suffering produced, we are impelled to do all that we know towards relief; as regards the outcome, we may be generally hopeful, if not confident. Biliary colic being the most commonly apparent indication of the presence of gall-stones, we may be ready to see in gall-stones things productive of much suffering, but having in their aspect more of the troublesome than the deadly. Gall-stones, indeed, travel so commonly by the same line, that we tend to believe that, in spite of delay occurring now and then, the terminus must be reached. The consideration which I have very imperfectly put before you may remind us of many possible accidents of the journey. A permanent block occurring on the line sends the travelling stone into many by ways of disaster. We may find the stone in such case causing abscess, causing perforation of the gall-bladder, causing intermitting pyrexia, causing glycosuria, causing distant inflammations, probably malignant disease, and certainly serious hemorrhage.

I rise from the contemplation with increased respect, born of sad experience, for the gall-stone. If biliary colic be no very terrible illness, the gall-stone is to be dreaded and feared, in proportion as there is less of the colic and more of accessory symptoms.

parts are acting, combination of their movements occur. The special combinations of movements are simply the result of the time of the action of each of the moving parts.

In observing movements of the digits, two modes of description may be employed: the series of movements of each digit may be described, or the series of coincidences or combinations of acts may be recorded. Either plan may be used, as convenience may dictate; thus it may be said that each of the five digits flexed at the same moment, or it may be said that the hand assumed "the convulsive posture." What we commonly call an action, such as writing or walking, is really an observed series of combinations of movements; this applies equally to spontaneous reflex, or voluntary action. When a child seizes an orange, we observe a series of movements, each of which can be performed separately; while the character of the action depends upon the time and quantity of each of the component movements.

Series of movements in plants were then described. The quantity of a movement is its second intrinsic attribute; it indicates the quantity of action in the nerve-centre. Quantities of motor action are studied in observing postures of the body, as signs of the ratios of force in the nerve-centres. In "the straight intended hand,"² we observe a sign of the temporary equilibrium of the nerve-muscular apparatus for flexion and extension; the posture may change, the fist being clenched; the change of hand posture is a sign of a change in the ratio of force acting in the apparatus for flexion and extension, the force of flexion being increased. Thus we see the outcome of the balance, or ratio of quantities of motor force, demonstrated by the observation of postures. Examples were given concerning other hand postures. It was argued that postures often indicate the condition of the nerve-system; that they are often changed by the sight of objects; that they are due to the ratios of force exerted by opposing portion of nerve-muscular mechanism; and hence, that postures are signs of ratios of action that can be controlled by incident forces, such as light, sound, mechanical irritation, etc.

The total quantity of movement seen in man may depend upon the quantity of blood-supply, and upon the amount of stimulation of the subject; the blood-supply and stimulation are needed to produce movements. In anæmia, and in starvation, the quantity of movement is lessened. In darkness but little movement occurs in a well-nourished man; the sight of an object may be followed by the head turning towards it, and by other series of movements such as result in the seizure of the objects. The quantity of movements in plants is mainly determined by light and heat. The proposition was then enunciated that every movement depends on some change in the time or the quantity of action in the nerve-muscular apparatus producing it, and that every such change in the apparatus must be preceded by some stimulus acting on it, immediately antecedent, or more remotely antecedent.

Among the forces which, acting upon a man, may stimulate and control the undivided movements, the following were mentioned and illustrated: light, heat, sound, mechanical irritation, etc. Every movement must have some sequence, whether it be to supply food, or be expended as physical energy. The sequence of a movement must be carefully separated from the cause or antecedent. To say that the bee visits the flower to get honey is to imply that the honey causes the visits. It was argued that the bee sees the flower, and afterwards gets the honey; that it is not the honey but the light reflected from the flower which determines the movements of the bee. The forces which control the time and quantity of the acts composing a series of movements are really the antecedents of the results of the action.

The sequences, outcome, or results of movements are often said to be useful, intelligent, voluntary, purposive, etc. What is meant by the outcome of an action being useful? What is the connotation of the term "useful" expressed in terms of physical criteria that can be observed? Concerning the meaning of the word "useful" as applied to the outcome of an action or movement, the primary idea seems to be that the outcome produces some physical change that can be observed. A movement which brings food to the subjects is said to be useful, partly because it produces some effect or impression on the subject. The movements of deglutition, respiration, and all the movements of organic life are useful, because, among other things, they impress the subject considered. We do not say that all acts which produce an impression are useful; but to be useful there must be survival in the subject producing the act, or in others considered at the same time. It is insisted that usefulness is not an intrinsic character of any movement, but a relation of its outcome to surrounding objects. Impressionability as a property in man may be indicated by movements. If the movements of a man are regulated and controlled by light or sound,

ABSTRACT OF LECTURES

ON

THE ACTION OF NERVE-CENTRES AND MODES OF GROWTH.

Delivered at the Royal College of Surgeons of England, February, 1887.

By FRANCIS WARNER, M.D., F.R.C.S.,

Hunterian Professor in Comparative Anatomy and Physiology at the Royal College of Surgeons; Physician to, and Lecturer on Botany at, the London Hospital.

LECTURE I.—ON THE ACTION OF NERVE-CENTRES.

MOVEMENTS are here studied as signs of action, observations being recorded by verbal description or by the graphic method.¹

To study a movement involves the consideration of the part moved, the muscles moving it, and the nerve-centres which stimulate them; while the movement itself indicates the time and quantity of action in the nerve-centre corresponding; these are called the intrinsic attributes of the movement.

The time of a movement often affects its outcome or sequence; movement of a single part is a comparatively simple fact: when several

¹ See *Journal of Physiology*, Vol. iv, No. 2, and Vol. vii, No. 4.

² See *Physical Expression, its Modes and Principles*, Chapter ix.