Conclusion.—These analyses, I hope, will show to the profession to what extent colchicum will alter the renal secretion, by supplying it, when deficient in its normal constituents—urea and uric acid—from the blood. From practical observations and experiments, I am inclined, therefore, to believe that to this property of colchicum its remedial action is in a great measure to be referred. Further analysis of the blood may, however, be thought necessary, but want of time has allowed me no means for this purpose. The analyses of the urine recorded are only a few selected from many others made by me, and were effected according to Becquerel's method. Colchicum has been employed in all forms of the disease, but it appears to me more particularly useful in articular rheumatism. My belief is that in proportion as the synovial symptoms predominate or mix themselves distinctly with the fibrous—in proportion as the disease approaches in its character to gout—we may expect to be successful with colchicum.

N.B.—Large doses are not requisite; 20 minims of the tincture of the wine may be given every six hours until some relief be obtained, or a grain of the inspissated juice or of the acetic extract of colchicum every four hours. Under this treatment the disease sometimes vanishes within three days, the medicine producing sickness and purging, and the rheumatism or rheumatic gout rapidly declining; occasionally, the same favourable event takes place, although there has been no disturbance of the stomach and bowels.

The cases in which colchicum appears to me to be more pre-eminent useful are those in which the disease attacks the joints and is of an erratic character, for example, suddenly disappearing from one joint, and as suddenly appearing in another; these, I believe, are the cases in which the heart and other internal organs are chiefly affected, and probably if the active operation of colchicum is secured early, these formidable secondary diseases are less likely to occur.

REMOVAL OF UTERINE APPENDAGES FOR THE RELIEF OF CONSTANT OVARIAN PAIN, INCESSANT VOMITING, AND DYS- MENORRHOEA: RECOVERY.

By A. C. BUTLER-SMYTHE, F.R.C.S. Ed.,

Senior Surgeon to the Grosvenor Hospital for Women, Westminster.

A. M., widow, aged 40, came under treatment in 1884, suffering from constant pain in the ovarian region and troublesome vomiting. She was a short, slight, dark-featured woman, with the following history. Had been married four years, but had never been pregnant. First menstruated at the age of 15, and ever since she has suffered from dysmenorrhœa. Seven years ago she had diphtheria. For more than two years she had been troubled with sickness at the time of her periods. Within the last seven months she has complained of constant gnawing pains in both iliac regions, and has been sick several times daily, the vomiting occurring at all hours of the day and night. No disease of the stomach could be detected. Physical examination revealed tenderness over both ovaries, pressure causing vomiting. The uterus was anteflexed, cervix elongated, and the os patent. The sound passed three inches. For nine months she remained under treatment, and during that time every known remedy was tried for her relief, but without avail, and she rapidly lost flesh and took to her bed. Her next period, July, 1885, was attended by aggravated symptoms. At her urgent request I agreed to remove her ovaries and tubes as a last chance of relief; so five days after the period had passed she was put under ether, and the abdomen opened. The appendages were removed without much difficulty, and the patient made a rapid recovery, the pain and vomiting ceasing from the date of the operation. A catamenial flow appeared on the second day, and lasted till the ninth, but she had no dysmenorrhœa. The ovaries were the smallest I have ever seen removed from an adult. They were hard and fissured on the surfaces, and about the size of small beans. The tubes were thick and congested.

Two years have elapsed since the operation, and the patient remains well. The symptoms throughout the illness, and their immediate cessation after the operation, taken together with the condition of the ovaries, would lead one to suppose that they were due to the cirrhotic state of those organs. Inasmuch as the operation was performed in order to save the patient's life, the result seems to justify the treatment adopted.

By permission of the Benchers, the gardens of Lincoln's Inn will be opened on evenings for the benefit of the poor children living in the surrounding districts.

SALIX NIGRA AS A SEXUAL SEDATIVE.

By J. HUTCHISON, M.D.,

Physician to Anderson's College Dispensary, Glasgow.

In many women pain in the ovarian region is a constant attendant upon the menstrual epoch. In some this is due to organic disease, but in a large number it is one of the manifestations of the neurotic temperament. Such cases are met with in all degrees of severity, from a slight amount of discomfort along with indications of globus hystericus, up to hystero-epilepsy in its most pronounced forms.

It has always been a slur upon our profession that when a method of treatment becomes popular or fashionable other methods are entirely discarded. At present, massage and isolation from relatives is the popular mode of treatment, and drugs occupy a secondary place, if, indeed, they have any place at all. Pecuniary difficulties, however, stand in the way of isolation or massage ever reaching the masses, and drugs will always be in employment. Several drugs are in daily use against the ailment, but with only partial success, and it is with the view of bringing before the profession a remedy which in my hands has produced results which I never had before while I was employing the bromides, valerian, assafoetida, etc., that I have sent this short notice.

Salix nigra, or the pussy willow, is a tree growing from 15 to 20 feet high. It is met with along the streams in the Southern States of America, and is credited with possessing tonic, carminative, and stimulant properties, besides being an astringent and antiperiodic. In the *Transactions of the Texas State Medical Association*, Mr. Pain reports many cases treated successfully with the drug. He prescribed it in cases of ovarian hyperæsthesia, uterine neuralgia, etc., and also in spermatorrhœa and nocturnal pollution. His verdict upon the drug is that it is a powerful sexual sedative, similar in its action to bromide, but without its depressing qualities.

Through Messrs. Thomas Christy and Co., of London, I obtained a supply of the fluid extract, and have been employing it for some months. The most numerous class of cases in which I exhibited the drug were women of a nervous temperament, in whom the nervous irritability reaches its height at the menstrual period, when, along with the general *malaise*, is added a very decided pain in one or other ovary. They also suffered from hemicrania, the pain being situated above the left eyebrow, and resembling the feeling as if a nail were being driven into the skull (*clavus*). Many of them, too, complained of a pain underneath the left breast, and extending round to the back. On one or two occasions I have noticed patients complaining of the above symptoms, and in only a moderate degree, under favourable conditions—as, for example, long-continued anxiety or alcoholism—go from bad to worse till they became hystero-epileptics. In cases of this kind, it is supposed that the centre of inhibition has in some way got out of gear, and the severity of the symptoms depends upon the amount of disturbance in this nerve-centre.

In cases where the ovarian distress was the symptom for which advice was sought, as being in the patient's eyes the most prominent, I usually succeeded in eliciting other indications of an irritable nervous system, and placed them upon half-drachm doses of the fluid extract of *salix nigra* three times a day. In quite 75 per cent. of patients so treated a great amount of relief was obtained after two or three days' treatment. Not only was the ovarian hyperæsthesia relieved, but the nervous palpitation of the heart was abated, and the patient felt in every way stronger.

I have also given the drug in two cases of nocturnal emissions with marked benefit. The pollution ceased entirely while the drug was being taken and for several months thereafter. Virile power and passion were not much if at all diminished, but the relief from the ailment gave them great satisfaction.

A CASE OF CHRONIC PROSTATITIS: A PATHOLOGICAL CONTRIBUTION TO THE STUDY OF THE PHYSIOLOGY OF THE PROSTATE GLAND.

By WILLIAM ANDERSON, F.R.C.S.,

Assistant-Surgeon to St. Thomas's Hospital.

THE patient, a healthy man, aged 23, came under treatment in May, 1884, with symptoms of chronic inflammation of the prostate of eighteen months' duration: the condition having supervened upon an acute abscess of the gland, the result of gonorrhœa.

The only symptom of which the patient complained when first seen was a persistent sensation of fulness and discomfort in the perineum,