was stitched to the abdominal wall, a drainage-tube inserted, and dry dressing applied. For a few days the urine was black like porter, but then it cleared, the patient made a capital recovery, and remains perfectly well.

CASE OF SUPRAPUBIC LITHOTOMY IN A CHILD.

By WILLIAM SYKES, M.R.C.S., Moxborough.

As few cases of suprapubic lithotomy in private practice have been reported, I venture to send the following, declaring at the same time my opinion that, from the comparative simplicity of the operation and from the generally good results obtained, it is destined to replace the lateral operation, in the same way as it was itself replaced by the latter.

J. T., aged 9½ years, but looking not older than 6, with lateral curvature of the spine and a strumous appearance, came under my care in November, 1886. He was suffering from all the usual symptoms of stone in the bladder. The urine was loaded with pus and mucus, and the child seemed in great pain. The history of the symptoms extended over two years. On sounding, a stone was felt; indeed, the sound had some difficulty in entering the bladder, as it at once impinged on the calculus, which appeared to be fixed.

I operated on November 25th, making the incision of the length recommended by Sir H. Thompson. I found that the bladder was not sufficiently distended, and had to make my final incisions dangerously near the peritoneum. I had no difficulty in extracting the stone between my two fingers when once I had entered the bladder. There was very little hæmorrhage. I put only one suture in the upper part of the wound, and I afterwards regretted placing this one in, as it caused some dragging. The wound was dressed with lint dipped in carbolised oil, merely laid on, and frequently renewed during the first three days; afterwards only changed night and morning, and moistened externally with the oil.

For the first week the urine passed exclusively through the wound, then began to come through both the natural and artificial passages, and finally ceased to trickle from the wound about one month after the date of operation. The wound itself, however, took on the appearance of a strumous ulcer, and did not finally heal until the end of January of the present year.

The stone was phosphatic, of the dumb-bell shape, with one end much larger than the other. It appeared as though it had been tightly grasped in the middle, and the resistance to the passage of the sound gave one the same impression. It weighed, when dried, a quarter of an ounce.

REMARKS.—The only difficulty in the operation was caused by the bladder not rising sufficiently over the pubes, owing to insufficient distension. The ivory instrument recommended by Sir H. Thompson for tearing the connective tissue could have been well dispensed with, and is of an unfair cost. After enjoying fair health and entire freedom from any difficulty with the bladder until the beginning of the present month (June), the child was attacked by broncho-pneumonia, and died from that disease, having only been ill three days.

SUPPURATIVE OTITIS MEDIA: PERFORATION OF MASTOID: TREPHINING AND EXPLORATION OF THE TEMPORO-SPHENOIDAL LOBE.¹

By JAMES BLACK, M.B., F.R.C.S.

H. H., aged 22, a lamplighter, was admitted into the Bouverie Ward of the Westminster Hospital on November 30th, under the care of Dr. Allchin. The note taken by the house-physician, Dr. Scott Sanders, at the time of his admission, states that: "On November 27th, the patient shivered and felt very sick, but did not vomit. For some days he had not been feeling well, 'had a dull sort of pain in his head, and lost his appetite.' The headache became very severe, and had been getting worse ever since. From a child he had had otorrhœa, for which he could not attribute a cause; he had never had scarlet fever, small-pox, nor syphilis. There was no history of a blow on the head."

¹ Abstract of a paper read on this case before the Medical Society of London on February 25th.

On December 8th (week after admission) the thermometer registered a temperature of 104° at 4 A.M., but showed a decline to 99° four hours later; and subsequent to this similar elevations and remissions frequently occurred; for example, on December 11th, there was a rise to 104.5°, followed by a rapid fall to 99°. The next day the patient had a rigor lasting twenty minutes. On December 14th (a fortnight after admission) I was first asked to see the patient. He had then a very offensive otorrhœa from his left ear; after cleansing the external auditory meatus of pus, I detected a large perforation at the anterior part of the membrana tympani. The patient informed me that at the age of 17 he became so exceedingly annoyed by the offensive otorrhœa, that he used effectually to plug the external auditory meatus with cotton-wool, so as to prevent the escape of the fetid discharge. This was naturally followed by inflammation about the mastoid process, and a sinus eventually ensued, from which matter for a time gained exit. He never sought advice for his ailment, but there was a depressed scar left at the site of the sinus, which was tender, and which bore out the veracity of the man's statement. Taking into consideration the repeatedly occurring elevations of temperature, and the fact of there having been a well-marked rigor, I was at once convinced of the serious nature of the case, and arranged to perforate the mastoid antrum on the following day, ordering in the meantime a saturated solution of boracic acid to be injected pleasantly warm along the meatus every three or four hours.

On December 15th, at 1 P.M., the patient having been put under the influence of an anæsthetic, a crucial incision was made over the mastoid process, and the soft parts reflected down to the bone. Not being able to detect any signs of a previously existing sinus under the depressed scar, I proceeded to very cautiously perforate the bone at the spot recommended by Mr. Barker, half an inch behind and above the auditory aperture, in a direction forwards and inwards, so as to open the "antrum," and at the same time avoid wounding the lateral sinus. HgCl₂ solution (1 in 2,000) was syringed into the perforation, and as it was believed that the fluid was seen to well up in the external auditory meatus, there appeared to be no doubt that the operation had succeeded in effectually opening up a communication with the tympanic cavity. No pus, however, gained exit through the opening made, and, on December 17th, a severe rigor, lasting twenty-five minutes, occurred. I then arranged with Dr. Allchin to trephine for intracranial abscess.

On December 18th, at 11.45 A.M., having explained the nature of the operation to the patient, and obtained his consent, I had him anæsthetised with chloroform, and, under the carbolic spray, dissected back the soft parts to the mastoid foramen, as advised by Mr. Barker, for the purpose of detecting if there were any matter thus gaining exit, as might possibly occur if suppuration existed beneath the dura mater corresponding to a thromboid lateral sinus, or if the disease had extended in a backward direction sufficiently to have given rise to cerebellar abscess; but as there was no such exit of pus, I proceeded to thoroughly syringe out the external auditory meatus and the perforation in the mastoid portion of the temporal bone with boracic acid lotion, and sprinkled the parts very thoroughly with iodoform, and then made an incision upwards from the original wound, and dissected up a V-shaped flap of soft parts, including the periosteum. The bone was perfectly healthy and normal in appearance. With a trephine having a crown of the circumference of a sixpence, I carefully removed a disc of the skull an inch and a quarter above, and only slightly behind, the external auditory meatus. I chose this in preference to the spot selected by Mr. Barker, namely, an inch and a quarter above, and a like distance behind, the auditory aperture, for I found that in a skull I examined this exactly corresponded to the position of one part of the course of the lateral sinus. The posterior branch of the middle meningeal artery crossed the middle of the hole made in the skull; this I cleared by nicking the dura mater on either side of the vessel, when I was able, with a bent aneurysm-needle, to carry a ligature across beneath the artery, and tie in two places and divide between. I then reflected the dura mater, which was normal in appearance, and exposed the brain, and, with a perfectly clean aspirating syringe, pierced the cerebrum slowly in a direction downwards and inwards, until it impinged on the bone corresponding to the roof of the tympanum. On drawing up the piston, no pus appeared. So soon, therefore, as all bleeding had thoroughly stopped, I brought the edges of the wound into apposition with silver-wire sutures, and, having dredged well with iodoform, applied salicylic-wool dressing.

On December 19th the patient had a rigor lasting twenty minutes, and the temperature mounted to close upon 105°. The rigors were attended with copious sweating, and the patient was getting much emaciated; he had a dry brown fissured tongue and sordes on the lips and teeth. I now, with Dr. Allchin's concurrence, commenced to

push quinine, giving the patient at first five grains with dilute hydrobromic acid every six hours, and a few days later the same draught every four hours, and ten days later, still increased the dose to seven grains every four hours; this was continued for eight days, when the quantity was gradually diminished. A large amount of stimulant was simultaneously employed; five ounces of brandy being given daily for three weeks, and then as much as eight ounces for the ensuing four days, when the quantity was again reduced to five ounces; and as much nourishment in the form of milk, beef-tea, eggs, and even chicken, was given as could be assimilated. The healing of the wound progressed favourably, in spite of the very severe constitutional symptoms, and with the exception of the small sinus, which still remains, was complete in about a month from the date of the operation. For a fortnight after the quinine was pushed in large doses, although the temperature repeatedly shot up, and on three occasions rose to close on 105°, yet not a single rigor occurred, and the fact that quinine will effect this result in pyæmia I noticed mentioned in *Erichson's Surgery*. At the end of this time, however, that is January 7th, a note taken by Dr. Humphreys who had come on duty as house-physician, states, "there was a rigor at 4 o'clock this morning, lasting thirty-five minutes, followed by profuse sweatings; the tongue dry, small, and red; impaired resonance at the right base behind with prolonged expiration;" and a note of the following day, January 9th, adds, "rigor at 5 a.m., lasting twenty minutes; the temperature taken after the rigor 105.8°, wakeful and very restless on the preceding night."

On January 10th, however, he commenced to improve and to put on flesh rapidly; he gained twelve pounds in weight in a little over a fortnight, and was able to leave the hospital apparently quite well, excepting for the existence of the small sinus, which has, I suspect, since healed. He could hear a watch held two inches from the affected ear, and had a very slight amount of otorrhœa.

OPHTHALMOLOGICAL MEMORANDA.

"OPHTHALMIC REPAIR AND SENILE DECAY" AND "A NEW METHOD OF TREATING AN OLD COMPLICATION."

I CAN thoroughly endorse Mr. Anderson Critchett's observation that the healing process after operation will in many cases go on satisfactorily while the physical powers are in the last stage of exhaustion and decay.

Some years ago I operated most successfully for cataract upon a gentleman in his 79th year, who was suffering from diabetes, complicated with gangrene of the left foot. I have also operated upon a considerable number of patients over 80 and 85 years of age, and one upwards of 90. In the last case the left eye was first submitted to treatment, and although the operation (which restored most excellent sight for all purposes) was performed without anæsthetics, and before the discovery of cucaïne, the patient suffered so little that, two months later, he insisted upon the removal of the cataract from the right eye, an operation which was likewise perfectly successful. Another case of special interest was that of a lady in her 78th year, who caught cold on her journey to Nottingham, developed the day after the operation double broncho-pneumonia, and died on the eighth day; although this patient was moribund from the first, and suffered greatly, the process of repair was uninterrupted throughout, and the result, so far as the eye was concerned (as ascertained on *post-mortem* examination by Dr. Morris and myself), as perfect as could be.

With regard to Dr. Mules's "new method of treating an old complication," I may mention that I have for years been in the habit of performing iridectomy with the knife wherever it has appeared to me that the patient's chances of recovery from cataract extraction would be increased by abscission of a portion of the iris. With a little practice this may be done with great precision, and a coloboma obtained in no way inferior to that secured by any other method, while the pain is incomparably less, either with or without cucaïne, and the risk to the patient from involuntary spasm or untoward motion reduced to zero. When an iridectomy as part of cataract extraction is necessary, it appears to me that this slicing off with the knife while making the preliminary incision is the best way to do it, but, in truth, it very seldom is necessary; for instance, I have, within the last three years, extracted for cataract in upwards of 500 cases without iridectomy, and, in the great majority of these, have secured a central and movable pupil, infinitely superior to anything obtainable by the more complicated operation.—I am, etc.,

CHARLES BELL TAYLOR, M.D., F.R.C.S.,
Surgeon to the Nottingham and Midland Eye Infirmary.

CATARACT-EXTRACTION: DR. MULES'S NEW METHOD OF TREATING AN OLD COMPLICATION.

DR. MULES is to be congratulated on his manipulative skill in the case he describes. Nevertheless, I think the proceeding he adopted is one to be deprecated as a routine practice in such emergencies. Even if the patient be absolutely quiet, the difficulties of the operation must be enormous when the cornea and lens capsule are in apposition, and the former so relaxed that it will not resist, but be pushed before the puncturing knife, making it very difficult to appreciate when the point has penetrated its whole thickness. Then if the lens be wounded, a complication not at all unlikely to occur, the section will be too small to admit of its extraction.

A very simple plan which I have successfully adopted, and for which I am indebted to Mr. Hussey, Consulting Surgeon to the Radcliffe Infirmary, consists in drawing the eyeball forward on the flat of the knife after the counter-puncture has been made, when the iris will roll off the edge of the knife, and the section can then be completed.—I am, etc., ROBERT W. DOYNE, Surgeon Oxford Eye Hospital.

DR. MULES, of Manchester, asks the question, "What shall we do if during corneal section (referring to the operation for extraction of cataract) the iris falls over the knife?" My answer would be: Never allow the iris to fall over the knife, by always performing the iridectomy as a preliminary and separate operation. In my own practice I now invariably adopt this proceeding, with certainly better results and a higher percentage of success than when excising the portion of iris at the same time as the extraction of the lens.

In the days of chloroform and ether, no doubt there was some excuse for performing the operation at one sitting, as the dread of the anæsthetic was not infrequently so great, that rather than submit to it a second time, the patients preferred to remain blind. Now, however, owing to the introduction of cucaïne, all this is changed, and when the patient finds that during the preliminary iridectomy he has experienced no pain, and exceedingly little discomfort of any kind after it, he approaches the second operation of extraction with good nerve.

The behaviour of the eye during and after the iridectomy gives some idea as to the way in which it will probably bear the second operation, and shows very certainly whether during the cucaïne anæsthesia there is any tendency to involuntary muscular spasm. By this means I was quite recently enabled to predict that it would probably be necessary during the extraction of the lens to give ether. In addition: 1. The operation of extraction is made shorter, and also unquestionably easier. 2. The iris cannot fall in front of the knife whilst making the corneal section. 3. The size of the piece of iris to be excised can be better regulated through the small incision made with a kerotome, than in the larger wound necessary for the extraction of the lens. 4. There is no bleeding from the iris, and consequently less risk of soft matter being left behind, as may sometimes be the case where the blood cannot easily be removed. 5. There is less fear of iritis. 6. The risk of hæmorrhage between the retina and choroid and into the vitreous is probably less, owing to the intra-ocular tension being diminished more gradually. 7. There is less fear of septic infection.

I am constantly asking myself, Why is it that this method of operating is comparatively so seldom adopted? Why is it, though well known, the exception rather than the rule?

ERNEST D. BOWER, Ophthalmic Surgeon to the
Gloucester. Gloucester General Infirmary.

TOXICOLOGICAL MEMORANDA.

A CASE OF ATROPINE POISONING.

A CHILD, James H., aged 2 years, was admitted into St. Mary's Hospital on January 9th, 1887. Two hours previous to admission he had swallowed the contents of a half-ounce phial containing a solution of atropine, which the mother had obtained on the previous day from an eye hospital. The amount of atropine taken was one grain.

On admission, that is, two hours after taking the poison, he presented the following symptoms:—The pupils were widely dilated and scarcely reacted to light, the muscles of deglutition were paralysed, the child being quite unable to swallow, and he was, on admission, in an excited state, the excitement, in the course of half-an-hour, giving way to a comatose condition. Three hours later, that is, five hours after taking the atropine, a red rash appeared on the face, arms, and upper part of the chest.

Treatment.—On admission I injected one-sixth of a grain of mor-