OBSERVATIONS ON A METHOD OF PROPHYLAXIS, AND AN INVESTIGATION INTO THE NATURE OF THE CONTAGIUM OF SCARLET FEVER.

[ILLUSTRATED BY COLOURED PLATE.]

By W. ALLAN JAMIESON, M.D., F.R.C.P. EDIN.,

Extra-Physician for Diseases of the Skin, Edinburgh Royal Infirmary; Consulting
Physician, Edinburgh City Hospital:

AND

ALEXANDER EDINGTON, M.B., C.M.,

Assistant, Surgical Department, Edinburgh University.

It is admitted that scarlet fever is little if at all communicable, in its early stages, from the sick to the sound. On this first principle rests the advantage of prompt isolation, since a fair chance is thereby afforded of arresting the further extension of the disease from this its primary source. It is equally agreed that the flakes of skin cast off during desquamation contain the active contagium, and these, if inhaled or swallowed, reproduce scarlet fever in those unprotected by their having already passed through the ailment. Dr. Jamieson has endeavoured to prove that, without separation, it was possible to prevent scarlet fever spreading from one member of a family to others of the same who had not had it, even though in close contact with him, or occupying continuously the same apartment with him. The two sources of infection are probably the exhalations from the mouth and throat in the early stage, certainly the particles of dry cuticle cast off in the later. The method recommended was to disinfect the throat by painting it frequently with a strong solution of boracic acid in glycerine (a saturated solution of boroglyceride in glycerine). In dealing with the skin more exact methods were available. Those consisted in the employment of warm baths every night from the very first, and in the application to the entire surface of the body, including the head, of an ointment composed of carbolic acid gr. 30, thymol gr. 10, vaseline 3j, simple ointment 5j, night and morning. In this way he believed that the scales of epidermis would never become contagious, and actual experience completely bore this opinion out.

Dr. Jamieson's theory, stated concisely, was, "That the contagium of scarlet fever present in the scales when inhaled or swallowed produced such changes in the blood and tissues as to lead to the symptoms characteristic of the disease; that this contagium, which was in all likelihood an organism, in due course of time reached the skin and there ripened, so that when the dry flakes of keratine were cast off, these contained the organism in a state ready for immediate multiplication when placed under suitable conditions. Arguing from the domain of cutaneous parasitic diseases, it seemed to me permissible to anticipate that, were the soil kept continuously in a state unfavourable for the ripening of the organism, it would either perish, or at least be thrown off immature. It also appeared evident to me that for the full development of the particulate contagium of scarlet fever air was necessary, since it seemed that the late desquamation contained it in largest amount." It may be argued that the urine in post-scarlatinal nephritis, the pus from suppurating cervical glands, or the muco-purulent discharge from the ear or nose, in instances where these sequelæ occur, might all be sources of contagion. This may indeed be true, yet it is well known that mild cases, in which the only abnormal condition is the desquamation, do spread the disease. Hence as "peeling" is the sole constant element, it is also the chief source of danger.

It is self evident that it is only in cases treated at home, among those unprotected by having had scarlet fever, and young, that the efficiency of such a method can be determined. During the three years since this method of prophylaxis was first published, many opportunities of testing it have occurred, and while in no one instance did the plan fail the following examples are specially illustrative.

CASE I.—C. H., aged 15, was one of a family of thirteen children, none of whom had had scarlet fever. There was a case of scarlet fever next door, and there had also been some at the school which he

attended. There was some suspicion that the milk-supply might have been the source of infection, as several families who obtained milk from the same dairy were nearly simultaneously affected, but careful inquiry elicited nothing. He shivered on January 28th, and on the 29th the rash was well out. He occupied a room on the ground floor of a large house, and was kept there so as to be further from the nursery, which was situated at the top of the house. M. H., aged 17, at that time an anæmic girl, who had had a troublesome cough all winter, went on January 30th to meet her sister at the Waverley station in the evening. On the 31st she had a sharp attack of diarrhoea, and at the same time the period came on, but scantily. Her evening temperature on January 31st was 101.8°, on February 1st, 102.6°, and she complained of sore-throat. It was not till February 2nd that a minute punctate red rash appeared on the backs of the hands, the back, and the ankles, and then gradually became more characteristic, while the temperature slowly fell. She suffered a good deal from rheumatism in the wrists, knees, and ankles, and this, a persistent tendency to diarrhoea, combined with the other circumstances mentioned above, prevented bathing being carried out quite so systematically as was desirable; but she was anointed regularly, and bathed as often as possible, and made an excellent recovery. The cough, which had lasted some months, also entirely ceased. Her elder sister frequently sat in the room beside her while she was desquamating. She occupied a room at the top of the house near the nursery. J. H., aged 19, shivered on February 2nd, rash out and temperature highest (103.6°) on the 3rd. He had a room adjoining his sister's on the top flat. All these three cases occurred within six days. The first case was from the outset separated from the others, and thus it is nearly certain the source was the same for all three, the periods of incubation being longer, but all within the generally admitted limit. No other case occurred in this family, though two servants, one who had had and one who had not had scarlet fever, suffered from slight follicular tonsillitis, in the desquamative stage of the scarlet fever cases, but associated with no rash or other symptom. This may have been a mere coincidence, since such sore-throats are common at all times in Edinburgh. From the position of the rooms inhabited by the three, one on the ground floor and two on the top, while there were no wet sheets hung up, but only the most careful treatment of the cases themselves; the scales must have been diffused to some extent through the house, yet the sole result was slight sore-throat of doubtful origin in two servants; all the other eleven children and servants remained quite well.

CASE II.—In this series the method was subjected to the most severe test. H. M., aged 9 years, became sick early in the morning of March 9th, and shortly after complained of sore-throat. He was seen on the 10th, when the throat and tongue presented the usual appearances of scarlet fever, and the rash was well out. His temperature was 102°, and pulse 100; next day the temperature had risen to 103.5°, but subsequently continued to fall. The family consisted of ten children, distributed through three apartments, and none of these ten had had scarlet fever. H. M. began perceptibly to peel on March 17th. A. M., aged 6, slept in the same bed with H. M., but with his head the reverse way. He was sick early on the morning of March 11th, and the rash was well out on the 12th. He also was desquamating on the 17th. D. M., aged 3, became ill on March 14th, and was more feverish than his elder brothers; the rash, too, was more vivid, and the throat more inflamed. He began to desquamate on the 24th. Thus all three were attacked within six days, and though two younger and five older children were all day in the same room with those affected, none took the disease. One sister, aged 14, had a slight follicular tonsillitis, which lasted two days, from March 21st to the 23rd, during, that is to say, the early desquamative period of her brothers' illness. Was this the ordinary scarlatinal sore-throat, so often seen in those who are nursing cases of scarlet fever; or may we regard it as an instance of attenuation of the virus, due to an arrest of development in consequence of the baths and inunction? The treatment was very faithfully carried out by the mother, then herself not strong; by no means an easy task under all the circumstances, but as she was assured that only in this way could the spread of the disease be arrested, if she kept her children at home, as she was desirous of doing, she was encouraged to persevere. There is little room for doubt that all three brothers contracted the fever from the same source.

CASE III.—T. W., aged 25, felt ill on December 30th, 1884, had sore-throat, and on the following day the rash came out. He was not seen till January 3rd, 1885, when all the symptoms of a well-marked but mild case of scarlet fever were present. His wife, who occupied the same room, as well as an infant under 1 year, and a brother-in-

TO ILLUSTRATE THE PAPER ON
A METHOD OF PROPHYLAXIS AND AN INVESTIGATION INTO THE NATURE OF THE CONTAGIUM OF SCARLET FEVER
BY W. ALLAN JAMIESON, M.D., AND ALEXANDER EDINGTON, M.B.

law, who lived in a room opening off the apartment, had not had scarlet fever. His wife attended to the bath and inunction, and cooked the food, yet no further case occurred, though desquamation was abundant.

CASE IV.—J. B., aged 3, had been playing near a house standing in its own grounds, where several cases of scarlet fever had been treated to the stage of convalescence. He had gone inside the house within a week before April 10th, 1885. At 4 A.M., on that day, he was sick, vomited, and complained of pain in the throat on swallowing. On the 11th the rash was distinctly out. The case was a mild one, desquamation was first noticed on the legs on the 19th. He was at once removed upstairs, to another part of the house, and treated in the same manner as the previous examples. There were three other children: two older had had scarlet fever, one, 2 years old, had not. The eldest girl had had the fever when two years old, and was not in contact in any way with her brother after the 10th, but she also had been in the neighbouring house. On the night of the 17th she became sick, and on the 18th complained of sore-throat, her tongue was red, with the papillæ elongated, and there was slight symmetrical congestion of the side of the fauces. In her case the disease was also mild, though the rash was well-marked. Apparently she had outgrown the protective influence afforded by her previous attack. No other case occurred in the house.

CASE V.—A. T., aged 7, took scarlet fever on December 1st, 1886, was attended by an aunt and servant, both however over 40, neither of whom had had it. They bathed, painted, and anointed her, yet had not even a sore-throat.

CASE VI.—J. and A. W., took scarlet fever on February 13th, 1887. The elder was aged 7, and the younger 4. The cases were mild, and were treated similarly to the others. A sister, aged 6, was constantly in and out of the room during desquamation, and yet did not take the disease. Neither did the father or the mother, both under 30; who were in constant attendance, and have not suffered.

CASE VII.—H. and D. took scarlet fever on February 8th and 10th respectively. The former was aged 4, the latter 7. Neither their mother nor a trained nurse who was engaged to attend them had had fever, and these were in constant association with them during their entire illness. The baths and inunction were continued till the end of February, when hæmaturia appeared. After this the bathing was persevered in, but the carbolic ointment was replaced by vaseline. No other case occurred.

CASE VIII.—In September, 1886, Dr. Jamieson saw a child aged 3, who had a smart attack of scarlet fever. In the same house there was a younger child, then aged 2, which could not be kept out of its sister's room. The treatment by baths, inunction, and application to the throat, was carried out from the first, and desquamation was used, and some months after that there had been no other member of the family attacked.

Mere insusceptibility will not explain the immunity enjoyed by those who were constantly near the patients during the desquamative period. It might possibly account for some, but comparing these and others similar with what happens in a family when no such precautions are adopted, the difference in favour of the plan is too great to be accidental. Clinical experience therefore supports the first assumption in the theory that the activity of the contagium is destroyed if the soil—the superficial layers of the skin—be kept continuously in a state unfavourable for its development. In this the removal of the desquamation as it forms by baths is one factor; the alteration in the character of the cast-off epidermis by antiseptic ointments is the other. Whether the employment of such substances as hasten cuticular exfoliation, such as sulphur or resorcin, would still further assist, must be decided by future observation.

The second assumption is, that the late desquamation contains the particulate contagium in largest amount, and that this argues that air is necessary for its maturation or ripening. The following cases tend to throw some light on this problem.

CASE I.—A nurse was sent on December 30th, 1886, to wait on three children who had been ill with scarlet fever for a fortnight. The temperature of one rose that day to 105°, and she died. The nurse remained well to January 20th, 1887, the fifth week of the children's illness, then took a slight sore-throat on January 21st, and noticed the rash the same day. On January 22nd she was admitted to the City Hospital with well-marked scarlet fever.

CASE II.—A mother, who had never had scarlet fever, occupied at night the same bed as her child suffering from it, during the whole course of its illness, yet did not herself become affected till the fifth week of the fever in her child.

CASE III.—A nurse was sent to the country to attend a case of illness of a non-infectious character. In a part of the house which com-

municated directly with the rest by a straight single stair, slept a man who had scarlet fever. The nurse, unprotected by a previous attack, passed the bottom of the stair several times daily, yet she remained free till between the fifth and sixth week of fever in the man, then returned home ill, and was admitted to the City Hospital with scarlet fever. There was no other known source from which she could have caught the disease. She never was in the man's room, only passed the stair which opened directly on his apartment, and continued well till he began to move about.

CASE IV.—A family of five persons residing in a district of Edinburgh, consisting of small self-contained houses, where there were no cases of scarlet fever at the time, took the disease on February 11th, 1887; and a young man in an adjoining house also fell ill on the same day. These got their supply of milk from the same dairy, while the other families in the terrace were supplied from a different one. All the families remained well till after March 13th, when the first of those affected began to go about. Nothing in the way of baths or inunction had been employed in their case. On March 19th a little girl, aged 7, was taken ill. She had been playing with some of those who had just recovered, and were probably still peeling.

In the late desquamation the cuticle becomes more freely separated, is older and drier, and consequently more perfectly oxygenated; the contagium, therefore, reaches its maturity at that stage in preference to an earlier one. The evidence submitted would not in the present position of medical science carry conviction. The positive existence of a micro-organism must be demonstrated; clinical experience does not hold the place it did once. Soon after entering on his duties as consulting physician to the City Hospital, in the end of last July, Dr. Jamieson mentioned his intention to Professor Chiene, who at once, in the kindest and most generous manner, not only placed his splendid bacteriological laboratory in the new University buildings freely at Dr. Jamieson's disposal, but also the services of his enthusiastic assistant, Dr. Edington. The questions to be answered, if possible, were:

1. To discover the organism in the desquamation, to the presence of which the contagious properties of these cast-off flakes may be inferred in some way to be due.

2. To ascertain approximately the period at which the organism first manifests itself in the desquamating epidermis.

3. Are the methods already described which have been found by clinical experience capable of neutralising the contagiousness of scarlet fever sufficient to destroy this organism, or, at least, to render it incapable of reproduction, though under suitable conditions otherwise?¹

The answer to the first two queries will be found in the statement adjoining, which represents the results of the cultivation and inoculation experiments from the blood and desquamating cuticle, which have been carried out by Dr. Edington in the Professor of Surgery's laboratory.

It will be found that a peculiar organism has been discovered to be constantly present in the blood up to a certain date, and in the desquamation after a certain date, in the progress of the disease. This organism, when inoculated into rabbits and calves, is capable of producing a disease in all respects identical with scarlet fever in the human subject.

The calves so inoculated were under the constant observation of Dr. Wood, the medical superintendent of the City Hospital, who has during his eight years' tenure of that office had exceptional opportunities of knowing what is and what is not scarlet fever. Dr. Wood has expressed his conviction that the disease which these calves suffered from as the result of inoculation was scarlet fever, and not any mere septicæmic or erysipeloid disease.

In endeavouring to solve the third question, some modifications in the duration of inunction had to be made, for the following reason.

In practice the baths and inunction which had been commenced on the first or second day of the disease, were continued until visible desquamation on the body had, as a rule, terminated—commonly, therefore, till the end of the fourth week of the disease. Desquamation visibly commences, on the average, at the ninth or tenth day of scarlet fever—it may be earlier—immediately on the subsidence of the rash; it may be considerably delayed, but the average period is as above. Now, as it was necessary to obtain the desquamating flakes at a period in the course of the disease at which the organism which had been discovered would certainly, if uninterfered with, be present in them, we had to select a time beyond the twenty-second or twenty-

¹ Or expressed otherwise, if a parasiticide capable of killing the organisms were applied to the skin at sufficiently short intervals.

third day, and we fixed on the thirtieth day to be quite sure. At that date the characteristic bacillus had been constantly found.

Seven cases were selected; the baths and anointing were persevered in from the second day of the disease till the eighth day of desquamation, the seventeenth or eighteenth day of the disease. An ointment for inunction containing carbolic acid in the proportion of one in sixteen had formerly been employed. Now and again slight evidences of absorption of carbolic acid had taken place, and, therefore, in these seven cases, an ointment of only one in thirty-two was made use of. On the eighth day of desquamation one leg was once more carefully anointed, enveloped in a thick layer of sterilized cotton wool, bandaged and covered with a stocking, and allowed to remain undisturbed till the thirtieth day.

The wool was then removed by Dr. Edington, with the same precautions as in the other cases, and the scales transferred to sterilized tubes for cultivation. It will be seen that the wool was put on at a period of the disease before the bacillus had been obtained from the scales, in cases where no disinfectants had been applied to the skin. This method subjected the procedure, therefore, to a rather severe test.

In cultivations of scales from five of the seven cases no bacillus was found. In two it appeared in the cultivating medium; but whereas, under ordinary circumstances, the bacillary pellicle is formed in thirty-six hours, in these two instances it took six days to develop any evidence of its presence in the jelly. In five, therefore, the powers of reproduction of the bacillus had been checked by the method of treatment adopted, while in the two others a remarkable retardation of the virus had resulted. There is little doubt that, had we continued the bathing and anointing for a few days longer, the bacillus would have been absent in all. We submit, therefore, that proof, clinical and experimental, has been furnished that by such simple methods we can neutralise the contagiousness of scarlet fever, so far as that arises from the desquamating flakes of cuticle. As the organism develops centrifugally, the outer and denser layers of the cuticle cast off at the commencement of desquamation contain little of it; but when these have been broken up and displaced, and access of air is afforded to the organism rising from beneath, it rapidly attains maturity and the power of reproduction when placed under suitable conditions. But apparently, at this stage in its growth, it is easily rendered sterile if brought directly in contact with a parasiticide such as carbolic acid. The organism, as is shown by Dr. Edington, and as Dr. Jamieson had inferred from clinical observation, is highly aerobic, and apparently ripens with great rapidity when the epidermic cells in or between which it is contained come in contact with the air, as they are successively thrown off in desquamation.

The valuable assistance rendered by Dr. Wood, Medical Superintendent of the City Hospital, who selected specially typical cases for our observations, and saw that all instructions were carried out to the letter, deserves our acknowledgment.

REPORT BY DR. ALEXANDER EDINGTON

(From the Bacteriological Laboratory of the Surgical Department, Edinburgh University).

Towards the latter end of October, 1886, acting upon instructions received from Professor Chiene, I began to make investigations into the desquamation of scarlet fever, with a view to establish, if that were possible, the specific nature of its infectiveness. It will no doubt be asked why such investigations in the first instance should not have been conducted on the blood. This it was intended to do later on, but meanwhile it was deemed fitting that the organisms found in protected desquamation might be each separately inquired into, and their particular infectiveness, if any, made clear by actual experiment on animals.

As a preliminary step in this inquiry, the epidermis of scarlatinal patients was submitted to a microscopical examination, aided by the various aniline staining methods, in order to find out the precise visible nature of the organisms it was afterwards intended to cultivate. A series of cases in the scarlet fever wards of the Edinburgh City Hospital were selected, and before desquamation had begun the part to be examined—whether arm or leg, as the case might be—having been carefully washed with a solution of carbolic acid (strength 1 in 30 of water), was covered over with a piece of gutta-percha tissue, previously washed with solution, the edges being fixed down to the skin by means of flexible collodion, while over all was placed a roller-bandage, in order that as far as possible the protective tissue might not be liable to displacement, nor contaminations allowed to pass beneath it. On the termination of the period selected, and when desquamation had become

pretty far advanced, the tissue was carefully raised, and some of the desquamation removed to an empty sterile test-tube.

Owing to this epithelium being dry and greasy, and further, containing many air-globules, it was found to be somewhat difficult to stain it perfectly. In order to overcome this, it was first treated with ether and caustic potash, to remove the oily matter, and then on having been transferred to the vessel containing the stain, the whole was placed under a bell-jar connected with an air-pump, and on a nearly complete vacuum having been created, it was found that the air-globules had given place to the staining fluid. The particles of epithelium were then washed, cleared in clove oil, and mounted in Canada balsam.

On microscopic examination micrococci and many minute stained points, smaller than micrococci, were usually to be seen, but nothing of the nature of bacillus or bacterium. Subsequently, small blisters were made on scarlatinal skin, and examination of the serum so obtained gave absolutely the same results.

But as it was possible, and even probable, that the infection from the skin might be due not to fully developed organisms but to spores, a series of investigations was instituted, such as might be likely to lead to the development of the supposed spores into their adult forms. A variety of methods were tried, requiring great precautions to be taken to prevent the intrusion of air-contaminations, but as such precautions were of too stringent a nature, and gave negative results; a simpler method was adopted, which, though it could not be expected to preclude entirely the entrance of contamination, yet was found to give fairly successful results. This method was as follows: the part having been washed first with carbolic soap and then with a solution of carbolic acid, 1 to 50 (we found that a stronger solution often gave negative results), was then enveloped in a sheet of sterilised absorbent wool sufficient completely to cover the part washed, which, in the case of the leg, included all the skin from above the knees downwards, toes included, and in the arm from above the elbow; then a roller-bandage having been carefully applied, a cotton stocking was drawn over all and made secure. The sterilisation of the wool used was effected by a dry heat of 160° C., kept up for an hour on each of two successive days. The sterility of such wool was tested by cultivation. On the termination of various periods of time, which varied from the nineteenth to the thirty-fifth day of the fever, the wool was removed by dividing it on the lower surface and in the long axis of the limb, and then, while an assistant gently raised it by grasping it by the middle of the upper surface, I, having previously sterilised my hands, introduced them beneath until I had reached the most central parts, carrying in them two tubes of sterile Koch's jelly which were, whilst here under cover, unplugged, and their open mouths pressed against the under surface of the wool, which was at the same time gently tapped upon from above, and then the tubes were replugged. In this way a small quantity of dusty desquamation was obtained, pretty free from air-contaminations, although we fairly anticipated that organisms, which might have been contained in the sweat-ducts, might be present. The tubes then were cleansed externally with a solution of corrosive sublimate, and their plugs singed with flame, in order that any germinative material here present might be destroyed and thus kept from contaminating experiments which might be conducted in the laboratory. The tubes were then placed in the incubator and kept there for two or three days at a constant temperature of 31° C., until, as a rule, more or less opalescence developed in the jelly, when cover-glasses were prepared from this usually impure cultivation, and sub-cultures, and plate-cultures made.

The examination of the blood was found to give the most satisfactory results only when made during the first three days of the disease, and at the bedside of the patient. It was found that otherwise the blood very rapidly coagulated, and that fibrinous threads were found in many cases around the organisms, which thus precluded either satisfactory examination or cultivation.

Cultivations from the blood were obtained as follows: Test-tubes of half an inch in calibre and ten inches in length were used; into each, three excessively fine capillary tubes, each nine inches in length, were placed, and the whole sterilised and plugged. A drop of blood was drawn from the finger, which was for two hours previously covered with a wet carbolic bandage (carbolic lotion, 1 in 20), care being taken while the drop was issuing that organisms did not alight upon it, by having a sheet wet with lotion held above it. The moment the drop issued a capillary tube was pushed into it, and on a little blood ascending, it was withdrawn, and the outer surface being cleansed with lotion, the tube was then gently passed into a tube of Koch's jelly, the plug of which was only so slightly moved as to allow of its entrance without coming in contact with it. The contained blood

was then ejected by a syringe into the jelly, care being taken that a small quantity was allowed to remain in the tube, so that contamination did not thus enter from the air in the free end of the capillary tube. The jelly was then incubated, and its further treatment conducted in the same way as those derived from the skin.

The method of isolating the various organisms demands some little attention. "Damp chambers" of the usual form, for example, flat glass bell-jars, fitting into shallow, flat glass dishes, were thoroughly cleaned, and in the latter were placed sheets of bibulous paper and enough of a solution of corrosive sublimate (1 in 1,000) poured in as would more than saturate the paper. Flat plates of window-glass, each five inches square, were cleaned and sterilised in the hot-air chamber, and thence, on cooling, transferred to the damp chamber, where they were placed upon supports. Small flasks of distilled water were sterilised by boiling on two successive days, and to each was added one or two drops of the original impure culture, which were then each freely agitated, in order to separate the individual organisms; this being done, a drop of this diluted seed-material was used to inoculate a tube of Koch's jelly, previously liquefied, and on being, by gentle agitation, thoroughly mixed, an assistant gently raised the bell-jar of one of the "damp chambers," and the liquefied jelly was gently emptied upon the plate and the cover replaced. This method, as a rule, gives excellent results, but one little point requires attention, that is, that if the flask of diluting water be quite cold, in adding to it the drop of seed-material, the jelly of this latter solidifies as extremely fine threads, which no amount of shaking will dissolve.

The following is a detailed description of the various organisms which have been isolated, and is taken to apply to cultivations made in jelly that has been neutralised by caustic potash, unless otherwise stated. It has been found that cultivations made in such jelly differed in some cases from that made in soda jelly, and this has been of considerable use as a means of distinguishing various organisms. The form of growth does not materially alter, but in potash jelly it is in almost all cases more luxuriant and more rapid, and, in addition, in some cases a peculiar and individual darkening of the deep needle-track growth results after a time.

ORGANISMS ISOLATED FROM THE DESQUAMATION AND BLOOD.

1. *Sarcina Lutea*.—This well-known organism occurred in 15 per cent. of the original cultures from desquamation.

2. *Streptococcus Rubiginosus* (provisional name).—Small spheres occurring frequently in short threads and as tetrads, measuring $\frac{1}{4}\mu$. Grown on gelatine plates it develops in two or three days as minute yellow points, if they be ever so little beneath the surface of the jelly, but if on the free surface are of a snowy whiteness. Sown in test-tubes of potash jelly, if the needle has not much disturbed the surrounding jelly, it develops a growth which is nearly of an equal calibre throughout, being made up of small cocoons closely appressed, and usually terminating somewhat abruptly in a cocoon of larger size. The deep growth is of a yellow colour at first, but rapidly passes through varying shades of brown, until, after several weeks' growth, it is of a rusty colour. The surface-growth is usually little, confined to the area bounded by the sides of the needle-track, and of a snowy whiteness. In older tubes this surface-growth frequently spreads in the superficial jelly, as white vertical lamellae. It occurred in 20 per cent. of the original tubes. Inoculations into rabbits, guinea-pigs, and pigs, gave negative results.

3. *Micrococcus Capsaformans* (provisional name).—Spherical or oval bodies, frequently united as dumb-bells, which are very often enclosed within a capsule, see Fig. VIII, and some also may be seen in Fig. XI. This capsule is well seen on staining with alkaline methylene-blue. In many cases these capsules appear empty, and when thus are usually kidney-shaped. It would seem that this is a form of endogenous, free cell-formation (not spore-formation), and that the depression corresponding to what would be the hilus of the kidney is the part from which the micrococci have been extruded. In one case I saw the micrococci lying together with this capsule half enclosing one of them. As a rule these capsules seem only to enclose two cocci, but in others one can notice division taking place, just as in a sarcina, for example, into four cocci; but I have never noticed the capsules corresponding to the latter. The single spheres measure 1.0μ — 1.2μ . On gelatine-plates it develops as white points of pretty rapid growth, having a very moist appearance.

In test-tubes of Koch's jelly the deep growth is usually somewhat scanty, and made up of cocoons, having a yellowish colour, but the surface-growth is distinctive; it is usually plentiful, forming a circular, flat, coin-like growth, on which concentric rings are beautifully seen. If the growth be ever so little impure, these lines are not to be seen. It occurred in 40 per cent. of the original tubes. Inoculations

gave negative results in rabbits, guinea-pigs, and pigs (see Fig. VIII).

4. *Diplococcus Scarlatinae Sanguinis* (provisional name).—Spheres measuring 1.0μ — 1.2μ , frequently, especially in fluid cultures, existing as dumb-bells. On gelatinous plates it forms creamy points of fairly rapid growth. On test-tubes of Koch's jelly it forms a deep growth of very closely appressed minute cocoons; while the surface-growth, usually irregularly heaped up around the needle-track, is of a slightly creamy white translucent colour. It occurred in 45 per cent. of the tubes taken from the desquamation, and in 30 per cent. of the tubes made from scarlatinal blood. Inoculations gave negative results, but this will be referred to when speaking of the bacillus scarlatinae.

Ascobacillus (provisional name).—Small rods in their fully developed form, measuring 0.8μ in length and 0.2μ broad, often as dumb-bells, made up of long ovoid spheres. This organism is peculiar in that the spores are contained in large sausage-shaped capsules many hundreds of times larger than the bacilli themselves (see Fig. VII). Sown in gelatine plates it grows as canary-yellow points of rapid growth, which in the course of a few days may be one-eighth to one-quarter of an inch in diameter, when liquefaction of the jelly usually begins. In tubes of Koch's jelly the deep growth is usually seen as a dense growth of closely appressed minute cocoons; the surface growth is of a canary yellow, with an irregularly sinuous outline. After some days liquefaction is observed in the jelly. This was obtained only in one case, with the bacillus scarlatinae, in which death resulted in twenty-four hours, and was obtained from the blood.

Bacillus Fulvus (provisional name).—Very small oval or rod-shaped motile bodies, measuring 1.0μ — 1.5μ in length, and 0.2μ in breadth. (See Fig. IV.) Sown in fluid broth it develops with great rapidity a creamy coherent pellicle, rapidly becoming wrinkled, but which is never very deep. This pellicle is easily detached from the sides, and if the flask be a little shaken it falls to the bottom, a new pellicle being in each case formed until the nourishing fluid becomes used up. Sown in test-tubes of Koch's jelly it forms a deep growth, made up of small cocoons, which are larger near the surface and smaller but better defined towards the bottom of the needle-track. The surface growth is semi-transparent and of a golden-brown colour, and somewhat slow in growth. Inoculation gave negative results.

Bacillus Arborescens (provisional name).—Rods measuring 0.8μ in thickness and 4.5μ — 12.0μ in length, most usually the former. This organism usually forms excessively long, jointed leptothrix filaments, either in jelly or in fluid media. It is non-motile (see Fig. IX). It is excessively difficult to isolate, as if a pure culture be grown on a gelatine plate, the resulting cultivation is quite invisible while the plate remains in the damp chamber; but if it be withdrawn and viewed by transmitted light, there is seen a ground-glass-like appearance in the jelly, caused by a fine reticulation of the filaments of growth. If one is working with a mixed cultivation containing this and unaware of its presence, one may fail to obtain it from the plate; but as in its growth on the plate it usually contaminates everything else, it may make its presence visible in the tubes made from other growths. Sown in test tubes of Koch's jelly, it develops little or no needle-track growth, but in a day or two at most there appear most beautiful arborescent and excessively fine filaments, shooting, in the first instance, toward the periphery of the tube, but later in all directions. It then looks just as if some very fine cotton-wool had been placed in the jelly and dispersed throughout. It does not cause any liquefaction of the jelly. If it be sown in the inclined surface of a tube of jelly it rapidly spreads over it, and in the course of twenty-four hours the surface of the jelly looks as if a piece of lint had been applied to it and withdrawn, leaving the woolly material in the jelly, but later it penetrates throughout. This usually occurred with the streptococcus rubiginosus, and was then easily isolated from the tubes by passing a needle down into the arborescent growth.

Inoculation into rabbits and pigs gave negative results. In guinea-pigs a well-marked local erythema developed, but did not spread far from the point of inoculation, and was unattended by any sequelae.

Bacillus Scarlatinae (provisional name).—Rods measuring 0.4μ in thickness and 1.2μ — 1.4μ in length, most usually forming excessively long-jointed and curved leptothrix filaments, motile (Fig. X). Sown on gelatine plates, it forms little points of liquefaction after several days. Sown in fluid broth and incubated, it rapidly forms a coherent parchment-like film, which is not easily broken up, and, even if the flask be well shaken, will yet float on the surface. Later, it develops an extraordinary deep wrinkling, which, if it be microscopically examined, is seen to be made up of a dense felted network of bacillary filaments. Sown in test-tubes of Koch's jelly, it rapidly liquefies it, but with no distinct growth-formation. The fluid thus formed is crowded with the motile bacilli, but a pellicle is not formed until the liquefaction is well advanced.

It occurred in every case but one² of the tubes made from the desquamation, if taken after the termination of the third week, but never before this. It also occurred in every tube made from scarlatinal blood if taken before the third day of the fever. Inoculation made into rabbits was always attended with a slight but marked erythema if they were young, but if older the erythema was better developed and most vivid at the point of inoculation. After two to five days a fine desquamation began, and usually lasted for a week to ten days. The temperature usually rose slightly in the younger rabbits, but more in the older; for example, 103° to 104° and in one case 106° F. During the first two days the animals huddled themselves in a corner of the cage and did not seem much to wish food, but usually after this they got quite lively again. The bacillus can be reobtained from the animal's blood. None died.

In guinea-pigs the same result obtained, but in them the desquamation is more copious and the flakes much thicker than in rabbits; and further, the hair if taken hold of came out.

A calf was then inoculated, and at the same time given some of a culture in milk. The calf was in good health at the time, and had a temperature of 99.5°. Six hours from the time of the inoculation, the calf developed great sickness, and the temperature taken in the axilla registered 103° F. (this was at 10 o'clock P.M.); the calf was then left for the night, but in the morning it was found to have died. At the necropsy, which was kindly made with me by Dr. Thomson, assistant to the Professor of Pathology, the following results were noted. General congestion of all the abdominal viscera; liver, well marked cloudy swelling, with many small petechiæ; spleen congested, pulp very fragile; kidneys, well marked cloudy swelling; heart, petechiæ numerous in serous surface of pericardium, and also in myo- and endo-cardium; cardiac muscle, well marked cloudy swelling; lungs, a considerable quantity of freshly exuded lymph was found at the base of the left pleura. On cutting into the tongue, with the contiguous portions of the pharynx and the larynx, there was seen to be an intense congestion of the posterior two-thirds of the tongue, and extending into the larynx, the rest of the mucous surfaces being pale in comparison. The fungiform papillæ stood out like red beads on the surface of the tongue. Examination of the tissues with the microscope showed a few bacilli here and there, but very sparsely distributed; while, on the other hand, there were many minute stained points, exactly as was found in scarlatinal blood. So far this agrees with *post-mortem* examinations made in the early stages of human scarlet fever; and if we had found bacilli plentiful there would have been suggested to me the probability of its being an organism either purely septicæmic, or one at least not specifically associated with scarlet fever.

Small pieces of tissue taken from the spleen and kidney were placed in Koch's jelly and incubated, with the result that the, to me, now familiar pellicle formed by this bacillus was developed. So far, then, this experiment and its *post-mortem* result show, first, that in all likelihood this organism had caused death, and while the state of the abdominal and thoracic organs would imply that the fatal result was due either to a high fever or to blood-poisoning, yet the state of the tongue, together with the suddenness of the attack and its rapid termination, suggests scarlet fever.

A second calf was inoculated, when only one day old, with the bacillus, care being taken that the inoculation was made with absolutely pure material as before, while at the same time the blood was examined with the microscope, and found to contain nothing of the nature of an organism. The inoculation was made in this case with a very carefully sterilised hypodermic syringe. At 6.30 P.M. this was performed, the temperature *per rectum* then being 99.6°. At 10 P.M. the animal took milk freely, and the temperature remained practically the same. Next morning, temperature 104°; sickness, slight diarrhoea, and complete prostration; throat, as far as could be seen, was in an inflamed condition. Towards the afternoon, the skin covering the thorax, upper part of abdomen, and inner side of forearm presented a general redness, which towards evening became still better marked. Evening temperature, 10 P.M., 102.8°. The animal now got a little milk. Next morning, the animal was still unwell, but the sickness was almost entirely gone, and it took a little milk. Rash very vivid, specially in throat. Temperature 101.8°, evening temperature 102°. Throat and posterior part of tongue very much inflamed. After this time the temperature became practically normal, and on the sixth day of the disease desquamation began on the chest and inner sides of forelimbs.

Further, blood was taken from a case of scarlet fever, on the third day of fever, with the care previously described, and, to prevent its coagulation, was passed into a tube containing a little liquefied sterilized

Koch's jelly; of this a part was injected into a guinea-pig and part cultivated by incubation. The guinea-pig developed the erythema as before, with succeeding desquamation, and the tube was found to contain the bacillus and diplococcus. But we have found that the diplococcus sanguinis, as well as the micrococcus capsuliformis and streptococcus, while they can exist in the blood for some days, yet of themselves produce no symptoms in the animals; thus, we may conclude that the injection of scarlatinal blood, taken during the period immediately preceding the height of the fever, is, in the case of guinea-pigs, and probably also in the others, attended with the same result as the injection of a cultivation of the bacillus itself.

This bacillus, then, is to be seen in scarlatinal blood if it be taken during the first two or three days of the fever, and if the cover glasses be spread immediately after the blood is drawn. It is also to be obtained from the desquamation after the twenty-first day of the fever—that is to say, in ordinary cases, when the desquamation begins, somewhere between the ninth and twelfth day. Of course, in the severest cases, where the desquamation begins earlier, it necessarily follows that we may anticipate its earlier appearance in the desquamation. I have examined the blood of some cases from day to day, taking care to control my experiments by performing them twice in the same day, with an interval of an hour or two intervening.

To take a typical case—a young lady, who had an extremely well-marked rash. On the second day the blood, being used for cultivation as presently described, was found to contain at the end of twenty-four hours' incubation only the bacilli; on the third day, bacillus and diplococcus sanguinis; fourth day, the same; fifth day, no bacilli, only diplococcus; seventh day, diplococci and streptococci rubiginosi; on the eighteenth day, cultivation made from the desquamation by removing the wool from one leg showed that no bacilli were present; on the thirty-first day the wool was removed from the other leg, and cultivation resulted in giving a good crop of bacilli, with the diplococcus and streptococcus.

In addition, seven cases were subjected to the treatment described by Dr. Jamieson, the wool being put on on the eighth day of desquamation, and removed, as a rule, on the thirtieth day, and cultivations then made as in the other cases. As a result, it was found that only two contained bacilli, and in these cases it required six days' incubation for development of the bacillary pellicle. Thus, while the treatment did not entirely destroy the bacilli, it nevertheless evidently had a distinct action in retarding their development from the spores, as usually, in other cases, a profuse pellicle is developed in twenty-four to thirty-six hours.

If desquamation be placed in milk kept at a gentle heat for twenty-four hours, the bacilli developed can be recognised by their motility and the pellicle which they soon form. Thus it is evident that, if milk be taken as food, containing infective desquamation, the spores will have a perfect nidus and suitable temperature afforded them for development.

From what has been stated, it is pretty well proved that this bacillus scarlatinæ is the specific cause of scarlet fever, and probably the other organisms found in the blood during and at a later period than the bacillus, are merely concomitants, and pass into the blood only after the vitality of the system and tissues has been lowered by the entrance into them of this specific organism. But if one injects or otherwise inoculates a perfectly healthy animal with the diplococcus scarlatinæ sanguinis, one finds that it is able to live for some little time in the blood without, in the case of animals, giving rise to any marked symptoms. This being so, then it is quite logical to suppose that if such be carried in infecting desquamation along with the bacillus, a double infection may ensue. Whether, in such a condition, the action of the bacillus is modified by it, cannot yet be definitely stated; but if we are to rely on what has been observed during the course of this investigation, then it might be stated that in those cases where one finds an extremely vivid and general rash, there is usually present from the first the bacillus and diplococcus. Unfortunately, much cannot be said of those extremely rapid and excessively malignant cases, because, owing to various reasons, I was unable, with the exception of one case, to get the blood investigated, and in it there was present, in addition to the bacillus, the ascobacillus mentioned and described. One thing which is very certain is that this bacillus scarlatinæ is able to form spores in the blood of a living animal, thus differing from the bacillus anthracis, which only does so on the near approach of death.

The rapidity of the growth of this organism—which is such that, if one inoculates a flask of broth, the diameter of which is two inches and a half, and it be incubated, the pellicle will develop and cover it entirely over in the course of four hours—suggests an explanation of the very short period of incubation in scarlet fever, and thus

² This tube was accidentally broken before time had been afforded for its development, if it had been present.

the particular pabulum in the body suited for its growth may be very rapidly used up, and so spore-formation ensue.

It has been endeavoured thus to find out, by a long course of careful investigations, embodying the results of several thousands of cultivations, the specific cause of scarlet fever, and these, the records of such, are now placed before the medical fraternity.

NOTES OF A CASE OF PORRO'S OPERATION.

By SIR T. SPENCER WELLS, BART., F.R.C.S.,
Surgeon to the Queen's Household.

So far as I have been able to ascertain, the case I now publish is the third in the United Kingdom where mother and child have both been saved by Porro's operation. The first was Dr. Godson's, the second Dr. Galabin's. In both these cases the impediment to parturition was deformity of the pelvic bones, the uterus not being diseased. In my case there was no pelvic deformity; but the presence of a large, solid, immovable uterine tumour rendered delivery by the natural passage impossible.

On May 7th, 1887, I saw a lady who had arrived in London that morning from Leeds. She was 37 years of age, and was married in April, 1880. She gave me a letter written the day before by Mr. Scattergood, of Leeds, containing the following history of her case:

"On Wednesday last I was sent for to see her at her residence four miles from Leeds, and met in consultation Dr. Glaister, her old medical attendant, who was leaving the village that day, and Dr. Stevenson, his successor in practice, who had only been in residence about a week. Having known the lady before her marriage, I had seen her in consultation about a year ago, and knew her to be the subject of extra-mural fibroids. In the beginning of November last, I saw her again, the question which had then arisen being a possible pregnancy dating from August 8th, 1886. Dr. Glaister and I then came to the conclusion that in all probability she was not then pregnant. The pelvis contained a large, hard tumour almost filling its brim, projecting low into the vagina, and making it almost impossible to feel the os uteri high up behind the pubic bone. I heard nothing more of the patient until a few days ago, when her husband called upon me and said that his wife was believed to be near her confinement. An immediate inquiry into her condition seemed imperative, and accordingly we met on Wednesday, May 4th. On examining her, the fibroids apparent on uncovering the body showed a movement which appeared to be the result of intermittent and almost rhythmical contractions of an enlarged uterus. We could not detect a foetal heart-beat, nor feel active movements; but the lady asserted that she regularly felt the latter, and Dr. Glaister is confident of having heard the foetal heart about a fortnight ago. It seemed to us that, as delivery *per vias naturales* could not take place, owing to the firm blocking of the pelvic brim, the probable alternative is Porro's operation."

The patient arrived in London in great discomfort, but not having suffered much from the railway journey; and I found, as Mr. Scattergood described, the pelvis completely blocked up by a solid tumour, the os uteri quite out of reach, the abdomen filled by a solid tumour, on the surface of which were several separate and movable masses, apparently outgrowths from the main tumour. To the right of the umbilicus was a solid mass, *balloting* freely, and capable of being pushed downwards and to the middle line. This, I felt confident, was the foetal head. I could not clearly detect foetal heart-sounds, but the placental murmur was very distinct to the left of the umbilicus and in the left iliac region.

By my desire Dr. Matthews Duncan made a careful examination of the patient on the following day (May 8th), and we saw her together on the 9th, both agreeing that Porro's operation offered the best prospect of life both to mother and child; and Dr. Duncan saying very distinctly that he believed delivery by the vagina (by craniotomy, evisceration, or any obstetric operation) was impossible. The chief point for consideration was whether, the full period of pregnancy expiring on the 13th, we should wait for that time, or for signs of commencing labour, or operate without delay. We decided upon the latter course, mainly on account of the very natural anxiety of the patient and her husband.

On the 11th of May, at 9 A.M., Dr. Day administering bichloride of methylene, and, assisted by Mr. Meredith and Dr. Godson (who undertook the care of the child), Mr. Scattergood, of Leeds, and Mr. Bishop, of Manchester, being present, all the preparations were made precisely as for ovariectomy, with all the usual antiseptic precautions, including the use of spray. I divided the abdominal wall in the

median line, midway between the umbilicus and pubes, to an extent of about five inches. The uterus was free, but was covered by layers of broad ligament extending upwards from the bladder. Some of these were divided and pushed aside and downwards, and the uterine cavity was opened by careful puncture. As the liquor amnii escaped I tore an aperture in the thin uterine wall large enough to enable me to draw out the child's head. The body followed immediately on very slight traction. The funis was compressed by two pairs of forceps and divided between them. Dr. Godson took charge of the child, which cried loudly. I did not interfere with the placenta, and finding that the opening in the abdominal wall was not long enough to allow the uterine tumour to pass out, I enlarged it as high as the upper border of the umbilicus, so that the opening was about seven inches in length. I was then able to draw out the uterus with the main tumour which had filled the pelvis, and several sub-peritoneal outgrowths, with both ovaries. As just stated, the bladder was partially separated from the uterus before opening the cavity. I now completed this separation, carefully pressing the bladder forwards as Mr. Meredith passed a constricting wire around the base of the tumour, and gradually tightened it by the screw. I was naturally uncertain as to the exact position of the ureters, under such altered conditions, and therefore kept the wire more upon the base of the tumour than was perhaps quite necessary. But when it was tightened, and I had cut away the whole mass some two inches above the wire, the stump was not much larger than a five-shilling piece. I cut away as much of the solid tumour, after further tightening of the wire, as could be done safely, and passed a long pin through the remaining stump across from side to side just above the wire. On proceeding to sponge the peritoneal cavity I found a free tumour as large as a flattened orange, surrounded by and closely connected with the cæcum, the appendix cæci, and a piece of omentum. The separation of this tumour from its connections had to be effected with extreme care, and added considerably to the time occupied in the operation. Several ligatures were applied during the process, and parts of the separated omentum were cut away. The opening in the abdominal wall was then closed by silk sutures precisely as in ovariectomy. The stump was fixed at the lower angle of the wound, the lowest suture being passed not only through the parietal peritoneum and abdominal wall on each side, but also through the peritoneal coat of the stump just below the constricting wire. Salicylic wool was then pressed all round the stump under the pin, and the surface of the stump was saturated with perchloride of iron. The abdomen was then covered with pads of salicylic wool and supported by strips of adhesive plaster and a flannel belt.

No provision was made for drainage. The time occupied from the commencement of inhalation of the anæsthetic until the patient was in bed was fifty minutes. From three to four pints of liquor amnii are supposed to have escaped, but the quantity was not measured. The child weighed six pounds. The uterus with the tumour and placenta weighed nine pounds. It and the detached outgrowth were sent to the museum of the College of Surgeons on the day of operation. They may be seen there, and it may be observed that the placenta is partially detached, the uterine cavity nearly closed, and the cervical canal open. As the wire must have been constricted at least an inch below the spot where the cervix was divided, it is almost certain that the whole uterus must have been removed.

Nothing could have been more satisfactory than the progress of the patient after operation. The breasts gave no trouble, and there was no sign of milk. There was no vomiting, and no interference with bladder or bowels from the depressed stump—nor any flatulent distension of the abdomen, although there was a good deal of pain at times for several days, always relieved by the passage of flatus, and requiring rather more opium than usual. This was given by the rectum in doses of fifteen minims of laudanum. The highest temperature noted was about thirty-six hours after operation, 101.2° F. It remained about 99° until the fifth day, and then became about normal. The pulse was about 100 to 116 until the fifth day, when it gradually fell to about 80. The urine was for some days loaded with lithates, and after a week contained a good deal of mucus for a few days, but it was quite normal by the time the pin and wire came off on the fourteenth day. I had not touched the dressing until the seventh day. Then the wool was changed, the wire tightened, some shreds of the stump cut away. After this, dry cotton and iodoform in powder were applied fresh once a day, and the screw occasionally tightened. The sutures, except the lowest, were removed on the seventh day, union of the abdominal wall being perfect. Four weeks after operation—June 8th, 1887—both mother and child are, as they have been throughout, as well as after a natural labour. About half an inch remains unclosed at the lower end of the cicatrix, but the granulations are quite healthy.