diathesis, especially as it is commonest in men over middle age. Some support is afforded to this by Case IV, where uric acid gravel was often passed; and if Dupuytren's contraction of the fingers be of gouty origin, as sometimes alleged, Case I also supports the possibility of a gouty origin.

It is possible that in some cases the irritation giving rise to the pulsation may originate in subacute arteritis or other changes in the coats of the vessel, for in most cases there is evidence of degenerative arterial changes. The *pathological nature* of pulsating aorta is little understood; *post-mortem* examinations have revealed no changes in the aorta itself (though I think the vessel is very commonly atheromatous), and the affection must be classed among the group of so-called functional diseases till our knowledge of it is increased.

It is usually, and probably correctly, ascribed to a disturbed action of the vaso-motor nerves or ganglionic plexuses connected with the aorta. A somewhat analogous local pulsation of vessels is seen in Graves's disease, and it has been noticed that occasionally pulsating aorta and Graves's disease co-exist. But their analogy does not explain very much.

It would seem that there is a paralytic condition of the vessel-wall arising from the disturbed vaso-motor influence, and that the vessel in consequence throbs unduly, but this alone cannot account for the violence of the throbbing whilst the heart is acting but feebly.

It will be asked how this vaso-motor disturbance is set up. Apparently it is a reflex result in some cases of lesions or irritation in the alimentary tract; in others, the reflex disturbance may result from changes in the arterial wall itself, possibly a gouty sub-acute arteritis.

The formation of a tumour of distended intestine is very probably due to the same cause as the pulsation of the aorta, namely, a paralysis of the tone of the intestine from disorder of the solar and intestinal plexuses.

The connection between the pulsation and the gastro-intestinal hæmorrhages is possibly, but very improbably, due to the violence of the pulsation rupturing small arterioles or capillaries, and it is more probably due to a vaso-motor paralysis of small arterioles, allowing them or other vascular areas to be gorged with blood until their walls give way. Hæmorrhages from the kidney, which are only ascribable to some such causation, are seen in locomotor ataxy. I have already suggested the possibility of the dryness of the mouth, and the occasional alteration of the pupils, being due to an occasional irritation of the sympathetic. The co-existence of Graves's disease with pulsating aorta tends to confirm this.

The pain and discomfort occurring with pulsating aorta are probably due to the stretching of the nerve plexuses by the pulsating vessel.

The *diagnosis* between the pulsating aorta and abdominal aneurysm cannot be made without care. In both there may be a violent impulse, with systolic *bruit* and thrill, and in both, as I have pointed out, there may be gastro-intestinal hæmorrhages and a tumour felt in the abdomen by the surgeon, and pain in the abdomen and back by the patient. The chief point on which the diagnosis must rest is the different character of the pulsation in the two cases; in aneurysm the pulsation is expansile, in pulsating aorta it is a violent jerking, and its direction is directly forwards (though if the vessel is very tortuous this may not be evident at once); it will, I think, in pulsating aorta, be almost always possible, by gradual, equable, firm pressure on the abdomen, to feel the vessel, and to ascertain that there is no aneurysmal tumour connected with it.

In some cases extreme tenderness over the aorta may prevent this, and then the diagnosis must depend on the course of the disease. In simple, pulsating aorta the pulsation is very apt to intermit at times, in aneurysm much less so: and aneurysms tend gradually to enlarge, and the symptoms become more severe, while the simple pulsation may continue for months or years without further development.

I think with care difficulty will rarely arise, though I should be very loth to say diagnosis is always possible, seeing an observer like Dr. Walshe writes that "to distinguish between incipient fusiform dilatation or sacculation on the one hand, and dynamic pulsation on the other, is, I believe, often a matter of impossibility."

In the *treatment* of pulsating aorta I have no specific remedy to recommend. If any mechanical cause be present, such as accumulation of feces, this must be removed where possible. Where impossible, as in the case of tumours, all that can be done is to relieve the pain by sedatives. In some cases, where due to gastric or intestinal irritation, it soon passes off when this is removed by ordinary remedies. I have found it relieved for a time by bismuth and opium, but the opium requires to be pressed. Bromide of potassium, as Dr. Walshe observes, may control the pulsation; but it, like the opium, is often required in somewhat large doses. Iodide of potassium with

alkalies certainly gave relief in the case of G. L., but here there were evidences of syphilis in the patient. In cases where it is persistent and no cause can be found for it in men of advanced life, the gouty diathesis should be borne in mind.

It is always to be remembered that the affection may—indeed, usually does—pass off in time; but many cases are very persistent and very intractable, and there is always present a tendency to recur, as the cases I have recorded tend to prove.

The distress often met with in these cases, the resemblance to aneurysm they bear, the co-existence of some symptoms which, though before observed, have dropped out of recent writings on the disease, and the very little notice given to it in many of the medical textbooks, will, I hope, be sufficient to justify my bringing these remarks on pulsating aorta under your consideration.

THE PRODUCTION OF ANKLE-CLONUS UNDER VARIOUS CIRCUMSTANCES.

By ANGEL MONEY, M.D., M.R.C.P.,

Assistant Physician to University College Hospital, and to the Hospital for Sick Children.

I AM greatly interested in the paper, published in the JOURNAL of September 24th, by Dr. Buxton, on the development of the ankle-clonus during the administration of nitrous oxide gas. The facts adduced are strongly in favour of the theory sketched in the first volume of Dr. Gowers's manual, and I may add that this theory has appeared to me to present many points in favour of its being a true explanation of many clinical facts.

For seven years past I have been engaged in the clinical study of ankle-clonus, and I may say with truth that I scarcely ever see a patient, either at hospital or at home, without attempting to elicit the foot-phenomenon.

In this brief communication I do not intend to give details, but merely to present some of the chief conclusions at which I have arrived, in order that other workers may compare my results with their own, and thus the general progress of knowledge may be facilitated.

The ankle-clonus can never be obtained when its neuro-muscular apparatus is in sound health by the passive method of dorsal flexion. Ankle-clonus developed by the dorsal flexion of the ankle always indicates one and the same pathological state of the lumbar centres, or corresponding nerve-paths of the spinal cord. So long as the muscles concerned are in a fit state for the production of the phenomenon, they exercise no influence in the direction of increasing the clonus, which is the same thing as saying that the presence of ankle-clonus signifies the presence of an abnormal state (of irritability) of the lumbar nerves and centres. The commonest cause of ankle-clonus is neurasthenia, and this neurasthenia may be developed in a rapid manner by the administration of nervine sedatives, by nitrous oxide gas, by chloroform, ether, etc., or it may be the result of hyperpyrexia, when its development may be exceedingly rapid; or, again, it may occur as the result of many, of perhaps all, continued fevers. Any wasting illness, such as cancer, may cause it; and, lastly, it may develop as the result of chronic nervous debility, or neurasthenia due to excessive work of the intellectual, emotional, moral, or physical nervous apparatus, but this seldom or never happens, unless there be the so-called neuropathic disposition, which again may be acquired as a neurotic product direct from the individual's ancestors, or be made by and in the individual himself either as a pure and simple effect of excessive expenditure of nervous energy, or as the side product of gout, rheumatism, and other toxæmic states.

I conceive, with Dr. Gowers, or rather with some adaptations of his theory, that this neurasthenia is most prone to exhibit its manifestations in that tract of nervous tissue whose source of nutrition rests on a more precarious basis than perhaps any other tract of nervous tissue. I mean the leg-fibres of the pyramidal tracts, which, reaching as they do from the "motor" cells of the cerebral cortex all the way through the internal capsule, crus cerebri, pons, medulla, and spinal cord, down to the lowest part of the lumbar swelling, must be at a disadvantage so far as their nutrition is concerned—seeing that the cortical motor nerve-cells have to energise, innervate, and nourish them from their origin in the cortex to their destination in the lumbar cord—with any other set of motor or sensory nerve-fibres. I conceive that the same explanation holds good when sclerosis of the lateral columns coexists (or is the efficient cause of the ankle-clonus) with the neurasthenia. My belief is that it is not the sclerosis, but the poorly nourished nerve-fibres of the lateral tracts, that permits the production of the foot-phenomenon.

One other suggestion I wish to make, and that is to the effect that

the administration of nitrous oxide gas, ether, or morphine, and some other drugs, may be used with a view to diagnosis. I have grounds for believing that individuals who are likely ultimately to develop undoubted nervous disease would, under such above-mentioned influences, more easily develop ankle-clonus than individuals whose nervous systems are wonderful in being perfectly sound and whole. As bearing at least at one point and in one direction on this question, I may mention that a case of locomotor ataxy first consulted me for symptoms which could not be classed as other than mere neurasthenia, and for which I prescribed, amongst other things, a hypodermic of one-sixth of a grain of morphine. The patient complained for two days after this dose of morphine, and only for two days, that he had great difficulty in commencing to make urine, but it was not till several months afterwards that genuine trouble in micturition was developed, and associated with other phenomena (lost knee-jerk and sexual power, and ataxic gait) of locomotor ataxy. I could not help thinking that the morphine acted upon a *locus minoris resistentiae*, and, in fact, had diagnosed a weakness which was not in the least suspected. In suspicious cases of organic nerve disease would it be justifiable to employ such drugs as diagnostic agents? This case was interesting also from the circumstance that the knee-jerks were certainly excessive when first I saw the case.

REMARKS ON RÖTHELN.

By HENRY DAVIS, L.K.Q.C.P., L.R.C.S.I., TUAM.

RÖTHELN frequently resembles ordinary measles; occasionally it still more closely resembles scarlet fever; yet röteln is not a hybrid. Measles alone, or scarlet fever alone, or both diseases in the same subject will not protect against it; and on the other hand, röteln confers no immunity, neither against measles nor against scarlet fever, nor, I am persuaded, in the least degree against a recurrence of itself. During the continuance of a lingering epidemic, I have seen every member of a large family, nine months of perfect health intervening, twice attacked by röteln. From what I have observed of this affection, it would surprise me little to see it seriously put forward that an attack of röteln rather increases than diminishes the liability to recurrence and to the invasion of other diseases. I had once the opportunity of observing röteln in a parturient woman; it was but a single instance and insufficient as an argument, still it is worthy of note that the complication in no way interfered with the normal course of labour, nor did it give rise to any unpleasantness afterwards, such as would be expected to follow an attack of measles or scarlet fever.

Some years ago, in Manchester, I saw a good deal of an epidemic of röteln. The invasion was suggestive of measles, accompanied by sneezing, lachrymation, photophobia, fever, general *malaise*, a slight sore-throat, and cough. About the end of the second day the eruption appeared without amelioration of the other symptoms; on the contrary, the cough became frequent and harassing, the throat was much complained of, the temperature rose often to 105°, and prostration was pronounced. The character of the eruption was not usually the same on the face and over the body. On the face, especially the prominence of the cheek, it appeared as a number of dusky circular or oval slightly elevated blotches, grouped without regularity. Over the body and limbs it was fairly uniform, much the colour of the scarlatina efflorescence, with, upon close inspection, many minute elevations. The palate, fauces, and tonsils were of a deep red, also presenting minute elevations; the tonsils were swollen. About the fourth day of the disease, with quickened breathing, increased cough and restlessness, with accelerated pulse and burning skin, it was usual to find at one or both sides of the spine a distinct area of broncho-pneumonia. I believe it was this complication which gave to the epidemic its very serious nature. The deaths which occurred during the continuance of the primary affection were, in my experience, all to be referred to broncho-pneumonia. The eruption faded in about five days, and was followed by coarse, branny desquamation and shedding of the hair. Convalescence was slow. Dangerous sequelæ were very apt to ensue.

As a very curious coincidence, if not something more, I remarked that many of those who recovered from röteln immediately contracted a set of symptoms exactly resembling the paroxysms of whooping-cough.

This epidemic left upon my mind the impression that röteln was a very serious malady, more serious than either measles or scarlet fever as they are usually seen. How different the epidemic which I have now briefly to describe!

About twenty cases of sore-throat collected from the same locality were brought under my notice. They were all very similar in appearance. They came one after another. They were communicated from

one to another. Age seemed to make no difference whatever in the liability. Sickness was hardly complained of, only considerable pain and difficulty of swallowing. On the throat alone was there any rash; the palate, fauces, tonsils, and the root of the tongue were closely studded with minute, bright-red elevations; the tonsils were swollen.

I diagnosed epidemic herpetic sore-throat, and I heard of epidemic tonsillitis in the practice of others. I watched my cases closely. The throat symptoms soon subsided. There was no suppuration. In a few instances, there remained for a long time enlargement of several small glands of the neck. In one case, the skin peeled from the index and middle fingers of both hands. Distinctly traceable to these there soon began to flow in upon me a straggling list of patients, all with sore-throats showing the characteristic elevated points; some with yellow patches on the tonsils. Many of these latter cases complained of rheumatism, both fugitive and stationary, and in not a few swollen joints were exhibited. In several there was a distinct rash, which generally occurred as patches of a rose-red miliary eruption especially on the forearms or beneath the knees. These patches might appear in the morning and be gone before the end of the day, or they might remain undergoing little change for several days; occasionally they faded and came out again; they seldom appeared upon the face.

In cases of this type, desquamation of the cuticle was most uncertain. It did not seem at all to depend upon the eruption. It occurred just as frequently when there was none, and the presence of an eruption was no indication that desquamation would follow. Again, the fingers alone might peel in a case where the rash had appeared only on the legs. Desquamation from the body was usually in light scales, from the hands in entire pieces. The disease was röteln. One of my sore-throat patients brought me to his house where every stage of röteln was fully developed. Subsequently I saw enough of the epidemic to enable me with confidence to enumerate the following distinctive appearances which the disease might assume:—

1. Slight sore-throat; without *malaise*, eruption, desquamation, or sequelæ.

2. Severe sore-throat; with moderate fever, rheumatic pains, sometimes desquamation of the hands and fingers, a liability to chronic glandular enlargement (frequently sub-occipital), but no eruption.

3. Symptoms similar to the last, with patches of rose-coloured miliary eruption, generally on the limbs, sometimes extending over the trunk and face, uncertain in duration; sometimes decidedly itchy and often followed by branny desquamation.

4. Considerable fever, some coryza, cough, aggravated sore-throat, a general eruption scarcely to be distinguished from that of scarlatina (the tongue in many cases also becoming scarlet), often outlasting both the sore-throat and *malaise*; desquamation, branny on the body, in whole pieces from the hands. Health impaired for some time after the attack.

5. Lastly, the attack may be ushered in by severe rigors and vomiting, or even by convulsions and protracted unconsciousness. The temperature may range above 106°. The eruption may assume the appearance of purple blotches on the face and a mixed character over the body. There may be a foul tongue, with red papillæ projecting; acute sore-throat, with regurgitation of liquids through the nose; a distressing cough; great prostration; desquamation, both branny and in pieces; a tendency to dropsy and to chest complications.

According to my experience of this epidemic, röteln may be followed by delicacy of the throat and chronic enlargement of the tonsils; delicacy of the eyes; chronic enlargement of many small sub-occipital and cervical glands, two or more of which may unite to form a considerable swelling; moist eruptions over the face and ears; protracted suppression of the catamenia. In one case, there was a distinct relapse, with reappearance of the eruption after fourteen days. In another, the attack was followed by erythema nodosum, which, however, may have been an affection independent of the röteln, or possibly brought on by menstrual derangement, the consequence of röteln. In two cases I thought I observed the characteristic eruption on the throat. I then lost sight of my patients. Subsequently I learned that they both had had rheumatic fever, and that the skin had peeled from their hands during the course of the fever.

I will conclude this sketch with a brief notice of four cases of undoubted röteln in two adjoining rooms. The first in sequence was a little boy who lay perfectly unconscious, passing from one attack of convulsions into another; temperature 106°; a foul tongue; an eruption of livid, slightly elevated blotches; and a running pulse. Beside him his sister presented almost the type of scarlet fever, with scarlet uniform rash, scarlet tongue, swelling of the neck, and burning skin. In the next room the parents were lying almost as sick as the children, complaining bitterly of their throats; the mother, without a particle