Published in final edited form as:

Bioorg Med Chem Lett. 2008 July 1; 18(13): 3724–3727. doi:10.1016/j.bmcl.2008.05.049.

Design, Synthesis and Evaluation of Bisubstrate Analog Inhibitors of Cholera Toxin

Guangtao Zhang^{a*,§}

*a*Department of Chemistry, University of Washington, Box 351700, Seattle, Washington 98195, USA

Abstract

Bisubstrate analog inhibitors in which a nicotinamide mimic is attached to a series of structurally diversified guanidines (arginine mimics) were synthesized and evaluated for inhibition of cholera toxin. The mechanism-based bisubstrate inhibitors were up to 1400-fold more potent than the natural substrate NAD⁺ and 400-fold more potent than the artificial substrate diethylamino (benzylidene-amino) guanidine (DEABAG) in an assay toward an intrinsically active mutant of wild-type cholera toxin.

Cholera is a devastating diarrheal disease that is caused by infection of the bacterium *Vibrio cholerae*. It is responsible for thousands of deaths each year.¹ People are usually infected by drinking contaminated water and they often suffer from profuse watery diarrhea.^{2–4} The massive intestinal fluid loss is primarily due to the release of the heterohexameric cholera toxin (CT). CT is composed of a catalytically active A subunit and five identical B subunits responsible for receptor binding.⁵ CT binds to the exterior of the human epithelial cell by its B subunits and gets internalized into the cell.⁶ It is first transported retrograde from the plasma membrane to the *trans*-Golgi^{6,7} and then to the endoplasmic reticulum (ER)^{6,8,9}. In the ER, the A subunit separates from the B subunits and the A1 chain is reduced,^{10,11} freed from the carrier holotoxin, and gets translocated to the cytosol,^{12,13} where the active A1 domain gains access to its substrate, including the α subunit of the stimulatory G protein (G_{sa}). A1 modifies G_{sa} through an NAD-dependent ADP-ribosylation reaction.¹⁴ ADP-ribosylation of Arg201 of G_{sa} locks the G protein in its GTP-bound state and persistently stimulates adenylyl cyclase. The consequent dramatic production of cAMP activates the Cl[−] ion channels.^{7,15} The imbalance of ions causes an efflux of cellular water to the gut, which eventually produces host dehydration and diarrhea.

Oral rehydration therapy is the most important treatment for cholera,¹⁶ however, its applications are limited by its reliance on large amounts of clean water in areas where outbreaks occur. On the other hand, substantial efforts have been made to develop vaccines, but they are only effective for a short period of time;¹⁷ there has been no known marketed drug that directly targets cholera toxin. Several promising strategies have been revealed for the design of therapeutic agents for CT. These include blocking the enzyme active site in the A subunit,^{18,19} disrupting the assembly of the AB₅ holotoxin,²⁰ intercepting the receptor binding of the B pentamer,^{21–24} and inhibiting adenylate cyclase.²⁵

*Corresponding Author. Tel: +1 206 465 0852; email:windzgt@gmail.com.

§Current address: Department of Biochemistry, Weill Medical College of Cornell University, New York, NY 10065

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Since very few studies have focused on the inhibition of the enzymatically active cholera toxin A subunit (CTA), our goal was to design inhibitors to block the CT active site that resides in A1. There have been disagreements regarding the reaction mechanism of ADP-ribosylation by CTA.^{26–28} However, the latest study on kinetic isotope effects suggested that the reaction followed a dissociative concerted mechanism (i.e. S_N2).²⁸ Lacking the crystal structure of substrate-bound CT when we initiated our work, we designed a series of bisubstrate analogs as potential inhibitors, which incorporate key moieties from both the ADP-ribose donor (NAD^+) and the ADP-ribose acceptor (Arg201 of G_{sa}). Our design of bisubstrate analogs is shown below (Figure 1). In this work, we selected benzamide as the analog for the nicotinamide moiety of NAD^+ , which is subsequently linked to a substituted guanidine, mimicking Arg201 of G_{sa} . The substitutions on the guanidine moiety allow one to test diverse molecular moieties to improve inhibitor potency. Our design is based on the assumption that both NAD^+ and guanidine have a well defined binding site and that linking them together could be an effective strategy to enhance inhibitor affinity.

The guanidine library was synthesized on the solid phase using Rink amide MBHA resin and the synthesis is summarized in Scheme 1, taking advantage of the solid phase synthetic method developed in our laboratory that employed an arylsulfonylthiourea-assisted guanidine synthesis.²⁹ In a typical synthesis shown in reaction sequence 1, 3-(N-Fmoc-aminomethyl) benzoic acid was coupled to the solid support under standard PyBOP/DIPEA conditions in DMA. After the removal of Fmoc, resin-supported **1** was suspended in methylene chloride and reacted with pentafluorophenyl chlorothioformate. The resin-bound thiocarbamate **2** reacted with 2,2,4,6,7-pentamethyl-dihydrobenzofuran-5-sulfonamide (Pbf- NH_2) in DMSO with the presence of potassium *tert*-butoxide and formed thiourea **3**. Subsequent reaction with an amine nucleophile and EDC in DMF gave the target molecule. Following reaction sequence 2, target compounds can also be prepared by the reaction of resin-bound **1** under either EDC or Mukaiyama reagent conditions with the corresponding thiourea precursors that are prepared in solution phase. Guanidine compounds without an alkyl linker were synthesized using the same method, with the only difference being the use of 3-(N-Fmoc-amino) benzoic acid instead. The product was cleaved off the solid support by a cocktail mixture (TFA/TIS/ H_2O v/v 94:3:3), and purified using reversed-phase HPLC in the presence of 0.1% TFA. The structural interpretation of the desired compounds was carried out by 1H NMR and mass spectrometry.

The target compound library was examined in an enzymatic assay toward inhibiting CTY30S (Table 1), an intrinsically active CT mutant which has displayed similar activity as wild-type CT.³⁰ An artificial substrate diethylamino(benzylidene-amino)guanidine (DEABAG) was used as an alternative ADP-ribose acceptor.²⁷ The reaction progress was monitored by analytical HPLC because all important components in the assay are chromogenic. Calibrated by the UV absorbance of the internal standard theophylline (Figure 2), the inhibitory activity of our compounds can be evaluated by the progress of ADP-ribosylation as compared to the control. The library compounds were screened at 1.0 mM concentration in 200 mM PBS buffer at 37°C and pH 7, and the screening results are described in Table 1. Kinetic studies were conducted for those compounds that demonstrated a percentage inhibition of over 90%. The K_m values of NAD^+ and DEABAG were determined to be 14 and 4.0 mM respectively, which are in good agreement with previous reports;^{31–34} while the IC_{50} values of compound **5a**, **5j**, **5l**, **5m**, and **5q** varied in a range between 40 and 270 μM (Table 1), with **5q**³⁵ being the most potent. Its estimated K_i is $\sim 10 \mu M$ and this result is better than the best CT inhibitors reported previously.¹⁸

On the basis of these results, bisubstrate analog **5q** is 1400-fold more potent than natural substrate NAD^+ and 400-fold more potent than DEABAG toward CT. Data analyses indicate that hydrophobic functionalities are preferred as R group. However, when we introduced some

other hydrophobic groups, such as biphenyl and 1-naphthyl into our analog, no affinity gain was obtained (data not shown). We did observe that analogs with a one-carbon alkyl linker inserted between benzamide and guanidine are consistently more potent in their inhibitory activities than those who share the same R yet without any spacer. It is worth mentioning that dynamic light scattering studies (DLS) have been carried out for some of the inhibitors with high potency to check for potential compound aggregation caused non-specific inhibition.³⁸ The DLS results indicated that the polydispersity of CT control was around 10.5% and the intensity of the CT peak represented 83% of all solution species. The DLS results for the assay mixture of **5q**, CT (at 70 nM), and all the other components showed a polydispersity of 12% and a percent intensity of 92% for CT. To verify the solubility of **5q**, its 2-bromo and 3-bromo isomers were also prepared. DLS measurements of solutions of CT with these isomers showed low polydispersity and high percentage intensity too (data not shown). This suggested that these mixtures are free of inhibitor aggregation, ruling out the possibility of nonspecific inhibition in kinetic assays with compound **5q**.³⁸ As a comparison, DLS of assay mixtures with compound **5a** showed an additional peak and the intensity of the CT peak dropped dramatically to 12.5% of all species. The new particle was calculated to be 2.2 μm in diameter, indicative of the existence of compound induced aggregation.

In summary, we have designed, synthesized, and evaluated a series of bisubstrate analog inhibitors toward CT. Our results demonstrated that the best compound **5q** is 1400-fold more potent than natural substrate NAD^+ . With the recently published crystal structure of a quaternary CTA1- NAD^+ : ARF6-GTP complex, it could shed new light on designing optimized bisubstrate analog inhibitors with improved potency.

Acknowledgement

I acknowledge the NIH for financial support (AI34501). I thank Profs. Erkang Fan, Christophe Verlinde, and Wim Hol for their stimulating discussions. I thank Dr. Claire O'Neal for providing the CTY30S mutant. I also thank Drs. Zhongsheng Zhang and Jason Pickens for technical assistance.

References and notes

1. World Health Organization. Weekly epidemiological record. 2006. p. 297
2. De SN. Nature 1959;183:1533. [PubMed: 13666809]
3. Dutta NK, Panse MV, Kulkarni DR. J. Bacteriol 1959;78:594. [PubMed: 13818912]
4. Finkelstein RA, Norris HT, Dutta NK. J. Infect. Dis 1964;114:203. [PubMed: 14183392]
5. Zhang R-G, Scott DL, Westbrook ML, Nance S, Spangler BD, Shipley GG, Westbrook EM. J. Mol. Biol 1995;251:563. [PubMed: 7658473]
6. Lencer WI. Int. J. Med. Microbiol 1994;293:491. [PubMed: 15149023]
7. Kahn RA, Fu H, Roy CR. Trends in Biochem. Sci 2002;27:308. [PubMed: 12069791]
8. Majoul IV, Bastiaens PI, Soling HD. J. Cell Biol 1996;133:777. [PubMed: 8666663]
9. Lencer WI, Constable C, Moe S, Jobling MG, Webb HM, Ruston S, Madara JL, Hirst TR, Holmes RK. J. Cell Biol 1995;131:951. [PubMed: 7490296]
10. Tsai B, Rodighiero C, Lencer WI, Rapoport TA. Cell 2001;104:937. [PubMed: 11290330]
11. Bastiaens PI, Majoul IV, Verveer PJ, Soling HD, Jovin TM. EMBO J 1996;15:4246. [PubMed: 8861953]
12. Schmitz A, Herrgen H, Winkeler A, Herzog V. J. Cell Biol 2000;148
13. Teter K, Holmes RK, Schmitz A, Herrgen H, Winkeler A, Herzog V. Infect. Immun 2002;70:6172. [PubMed: 12379695]
14. Tsai B, Rapoport TA. J. Cell Biol 2002;159:207. [PubMed: 12403808]
15. Salmond RJ, Luross JA, Williams NA. Expert Rev. Mol. Med 2002 October 1;:1. [PubMed: 14585161]

16. Pulungsih SP, Punjabi NH, Rafli K, Rifajati A, Kumala S, Simanjuntak CH, Lesmana M, Subekti D, Sutoto FO. *J. Health Popul. Nutr* 2006;24:107. [PubMed: 16796157]
17. World Health Organization. Weekly epidemiological record. 2004. p. 281
18. Zhou GC, Parikh SL, Tyler PC, Evans GB, Furneaux RH, Zubkova OV, Benjes PA, Schramm VL. *J. Am. Chem. Soc* 2004;126:5690. [PubMed: 15125661]
19. Parikh SL, Schramm VL. *Biochemistry* 2004;43:1204. [PubMed: 14756556]
20. Hovey B, Verlinde CLMJ, Merritt EA, Hol WGJ. *J. Mol. Biol* 1999;285:1169. [PubMed: 9887271]
21. Fan E, Zhang Z, Minke WE, Hou Z, Verlinde CLMJ, Hol WGJ. *J. Am. Chem. Soc* 2000;122:2663.
22. Zhang Z, Merritt EA, Ahn M, Roach C, Hou Z, Verlinde CLMJ, Hol WGJ, Fan E. *J. Am. Chem. Soc* 2002;124:12991. [PubMed: 12405825]
23. Fan E, Merritt EA, Verlinde CLMJ, Hol WGJ. *Curr. Opin. Struct. Biol* 2000;10:680. [PubMed: 11114505]
24. Pukin AV, Branderhorst HM, Sisu C, Weijers CAGM, Gilbert M, Liskamp RMJ, Visser GM, Rol HZ, Pieters J. *ChemBioChem* 2007;8:1.
25. Zenser TV. *Proc. Soc. Exp. Biol. Med* 1976;152:126. [PubMed: 1265071]
26. Oppenheimer NJ. *J. Biol. Chem* 1978;253:4907. [PubMed: 209022]
27. Soman G, Narayanan J, Martin BL, Graves DJ. *Biochemistry* 1986;25:4113. [PubMed: 3017413]
28. Rising KA, Schramm VL. *J. Am. Chem. Soc* 1997;119:27.
29. Li J, Zhang G, Zhang Z, Fan E. *J. Org. Chem* 2003;68:1611. [PubMed: 12585918]
30. O'Neal CJ, Amaya EI, Jobling MG, Holmes RK, Hol WGJ. *Biochemistry* 2004;43:3772. [PubMed: 15049684]
31. Larew JS, Peterson JE, Graves DJ. *J. Biol. Chem* 1991;266:52. [PubMed: 1898725]
32. Galloway TS, Van Heyningen S. *Biochemistry* 1987;244:225.
33. Moss J, Manganiello VC, Vaughan M. *Proc. Natl. Acad. Sci. USA* 1976;73:4424. [PubMed: 188038]
34. Mekalanos JJ, Collier RJ, Romig WR. *J. Biol. Chem* 1979;254:5849. [PubMed: 447682]
35. N-(4-bromobenzyl)-N'-(3-aminocarbonylbenzyl)guanidine (**18**). From 0.078 mmole of resin-supported **A**, **18** was obtained in 82% yield as the TFA salt (30 mg). MS:361.0 and 363.0 (M+H)⁺. ¹H NMR (CD₃OD) δ 4.46 (2H, s, CH₂), 4.55 (2H, s, CH₂), 7.31 (2H, d), 7.62 (4H, m), 7.95 (2H, m).
36. Segel, IH. *Enzyme kinetics: behavior and analysis of rapid equilibrium and steady state enzyme systems*. New York: Wiley; 1975. p. 106
37. Both product peaks were collected separately, and N-(ADP-ribose)-N'-diethylamino(benzylidene-amino) guanidine was verified by both MS and ¹H NMR: MS: 775.2 (M+H)⁺. ¹H NMR (CD₃OD) δ 1.04 (6H, t), 1.81 (4H, q), 3.93–4.55 (8H, m), 5.53 (1H, m), 6.05 (1H, d), 7.52 (2H, m), 7.85 (2H, m), 7.86–8.17 (1H, m), 8.18–8.27 (1H, m), 8.30–8.51 (1H, m).
38. Seidler J, McGovern SL, Doman TN, Shoichet BK. *J. Med. Chem* 2003;46:4477. [PubMed: 14521410]

Figure 1.

Design of bisubstrate analog targeting CTA. This analog prototype includes three features: 1. benzamide would mimic the nicotinamide portion of the substrate NAD^+ ; 2. incorporation of substituted arginine (R_1 and R_2), mimicking Arg_{201} moiety of $\text{Gs}\alpha$ as ADP-ribosyl receptor; 3. insertion of the methylene group as a spacer (in parentheses).

Figure 2.

A typical chromatogram for an HPLC-based assay of CT. The one in blue is a control run; the one in red is a run with 0.33 mM of **5q**. Each peak has been labeled. The two peaks overlapped in the product area have both been verified by mass spectrometry and ^1H NMR to be the reaction product.³⁷ As a result, they were both monitored.

Scheme 1.

Solid phase synthesis of CT inhibitors. For analogs without alkyl linker, 3-(N-Fmoc-amino) benzoic acid was used in step b instead. Both sequence 1 and sequence 2 are applicable. Reagents: a) 20% piperidine/DMA; b) PyBOP, DIPEA, DMA; c) EDC or Mukaiyama reagent, DMF; d) 94:3:3 TFA:H₂O:TIS; e) pentafluorophenyl chlorothioformate, DIPEA, CH₂Cl₂; f) PbfNH₂, potassium *tert*-butoxide, DMSO.

Summary of screening results of all bisubstrate analogs^d and IC₅₀ and estimated K_i for selected compounds.

Table 1

Structure	Compounds	R1	R2	n ^b	% Inhibition ^c	IC ₅₀ (μM)	Est. K_i ^{36d} (μM)
5a ^e	ⁱ Pr	ⁱ Pr	1	94	115±18	30±5
	5b	Et	Et	1	24	-	-
	5c	ⁱ Bu	H	1	39	>2000	>500
5d		H	0	22	-	-
	5e ^e		H	1	57	-	-
	5f		H	1	68	-	-
5g		H	1	76	-	-
	5h		H	1	34	-	-
	5i		H	1	57	-	-
5j		H	1	93	272±16	71±4
	5k		H	1	97 ^f	-	-
	5l ^e		H	1	96	49±6	13±2
5m	Bn	H	1	93	192±7	50±2
	5n	ⁱ Pr	ⁱ Pr	0	15	350	90
	5o	ⁱ Bu	H	0	77	-	-
	5p	Ph	H	0	12	>2000	>500
5q ^e		H	1	98	31±12	8±3
	5r	Bn	H	0	10	-	-

Structure	Compounds	R1	R2	n ^b	% Inhibition ^c	IC ₅₀ (μM)	Est. K_i^{36d} (μM)
	5s		H	0	58	-	-

^a All the synthetic compounds were evaluated as opposed to a random non-inhibitor, galactose, which serves as negative control.

^b n represents the number of methylene group as a spacer (also see Figure 1).

^c The screening was performed at 1 mM in PBS buffer; only selected compounds were studied for IC₅₀ measurement.

^d The K_i values are estimated from the equation: $K_i = IC_{50} / (1 + [S] / K_m)$.³⁶ The IC₅₀ are those at 40 mM NAD⁺. $K_m = 14$ mM was used in the calculation.

^e These compounds have already been examined in dynamic light scattering (DLS) studies.

^f IC₅₀ of this compound was not studied due to insufficient amount of materials.