The terror of television

Made worse by family stress

EDITOR,—D Simons and W R Silveira describe post-traumatic stress disorder in two 10 year old boys, which was apparently precipitated by the programme *Ghostwatch* shown on television on Hallowe'en (31 October) 1992.¹ We have treated an 11 year old boy who was referred with a 12 month history of sleep difficulties, nightmares, fear of the dark and of sleeping alone, fear of ghosts, intrusive thoughts, and panic attacks; the onset of these symptoms had been sudden, occurring immediately after he had watched *Ghostwatch*. He had become phobic about going upstairs to his bedroom (he said that the house reminded him of the haunted house in the television programme) and had begun to stay overnight in an aunt's flat. He had had no symptoms before watching the television programme, although his mother described him as having a slightly anxious personality. His older sister, however, had a history of encopresis and enuresis and emotional disorder, which had been diagnosed after a lengthy and acrimonious parental separation several years previously.

The boy responded to cognitive and behavioural strategies addressing his anxieties and behaviour, including gradual reintroduction to his bedroom, and ultimately he could sleep in his own bed without a light being left on. This boy's disorder seems undoubtedly to have been precipitated by the television programme, although it was maintained by factors within the boy and the family.

FIONA FORBES
IAIN MCCLURE

Department of Child and Family Psychiatry,
Royal Hospital for Sick Children,
Edinburgh EH9 1LL

1 Simons D, Silveira WR. Post-traumatic stress disorder in children after television programmes. *BMJ* 1994;308:389-90. (5 February.)

Anxious children at greatest risk

EDITOR,—D Simons and W R Silveira report two cases of post-traumatic stress disorder after the television programme *Ghostwatch*.¹ We have recently seen four children with symptoms apparently precipitated by television programmes.

An 11 year old boy was referred with a one year history of panic attacks, which started after he saw *Ghostwatch*. He had always been a nervous child, unable to stay in his own bed at night. His mother had also seen *Ghostwatch* but had not realised that it was a hoax and had been unable to reassure her son about his fears.

An 11 year old girl was referred urgently in November 1992 because of panic attacks that occurred during school assembly. Her mother blamed *Ghostwatch* but said that the girl had always been nervous and shy.

A 14 year old boy continued to suffer from anxiety, persistent sleep disturbance, and intrusive visual images of ghosts at night one year after viewing *Ghostwatch*. He was regarded by his parents as nervous and emotional.

An 8 year old girl presented with episodes of panic, clingy behaviour, and complaints of generalised pains three months after becoming distressed while watching an episode of the tele-

Priority will be given to letters that are less than 400 words long and are typed with double spacing. All authors should sign the letter. Please enclose a stamped addressed envelope for acknowledgment.

vision programme *Casualty*. Closer questioning elicited the fact that the girl had witnessed her mother suffering an episode of severe chest pain two weeks before the programme. There was no evidence of abnormal temperament or of family dysfunction. The parents had not linked their daughter's symptoms with seeing her mother in pain.

We suggest two aetiological considerations that may be important in understanding sustained psychological reactions in children after they have watched television programmes. Firstly, Bloch *et al* described how parental reactions at the time of the trauma determine children's subsequent psychological recovery.² The realistic quality of both *Ghostwatch* and *Casualty* may have prevented the parents from adequately containing their children's anxieties. Secondly, the four children we saw either had anxiety prone temperaments or had had recent experience of other psychological trauma. The diagnosis and aetiology of prolonged psychological reactions after television programmes seem to require further debate.

MARJORIE BAILLIE
ANNE THOMPSON
CAROLE KAPLAN

Fleming Nuffield Unit,
Newcastle upon Tyne NE2 3AE

1 Simons D, Silveira WR. Post-traumatic stress disorder in children after television programmes. *BMJ* 1994;308:389-90. (5 February.)

2 Bloch DA, Silber E, Perry SE. Some factors in the emotional reaction of children to disaster. *Am J Psychiatry* 1956;113:416-22.

Distress not confined to children

EDITOR,—D Simons and W R Silveira describe an important, probably underrecognised, adverse effect of television.¹ Adverse effects can occur in other susceptible groups. Elderly people, who often use television as a social substitute, suffer substantial psychological effects, including that of 24 hour television on temporal orientation. This contributes to day-night reversal, increased demand on general practitioners' time, and impaired compliance, as described in psychoses.²

The "deludogenic" influence of television is also important.³ These delusions principally emanate from the person's interpretation of programmes but may be influenced by the programme's content. We have observed considerable distress from such delusions both in the affected person and in his or her friends, relatives, and carers. An example of this is seen in the reidentification syndrome described by Fulton *et al*.³ In this, a televised character's personality and behaviour are transposed on to someone with whom the affected person is familiar. There are obvious, potentially distressing, implications of this reidentification. For example, one woman was convinced of the

transvestitism of her consultant after watching a television programme. In these circumstances, explanations have to be convincing, but down regulation and cure of such delusions are possible through withdrawal of television.

Elderly people in hospital and in the community may be exposed to many hours of television daily. The possibility of an adverse reaction to television should be considered, particularly in people with affective or psychotic disorders or with organic brain disease.

YVONNE FOGARTY
FIONA MORRISON
JAMES D FULTON

Stracathro Hospital,
Brechin, Angus DD9 7QA

1 Simons D, Silveira WR. Post-traumatic stress disorder in children after television programmes. *BMJ* 1994;308:389-40. (5 February.)

2 Howard R. All night television can damage your health. *Br J Psychiatry* 1991;158:577-8.

3 Fulton JD, MacDonald E, Scott PJW. Television: fantastic reality or realistic fantasy? *J Am Geriatr Soc* 1990;38:829.

Post-traumatic stress disorder cannot follow television viewing

EDITOR,—D Simons and W R Silveira report possible repercussions when children are exposed to horrific images in the media.¹ They do not, however, describe two cases of post-traumatic stress disorder. According to the criteria of the *Diagnostic and Statistical Manual of Mental Disorders*, third edition, revised, a person with post-traumatic stress disorder has "experienced an event outside the range of usual human experience and that would be markedly distressing to almost anyone."² Examples of this cited include threats to your life and the sudden destruction of your home or community.

The rapid resolution of the children's symptoms suggests that the children suffered a brief anxiety reaction to the television programme; although they may have exhibited some of the features of post-traumatic stress disorder, this diagnosis in their cases is inappropriate.

SIMON THACKER

Department of Health Care of the Elderly,
Medical School,
Queen's Medical Centre,
Nottingham NG7 2UH

1 Simons D, Silveira WR. Post-traumatic stress disorder in children after television programmes. *BMJ* 1994;308:389-90. (5 February.)

2 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 3rd ed. Revised. Washington, DC, APA, 1987.

**We received a further seven letters questioning the diagnosis of post-traumatic stress disorder in these children.

Management of breast cancer

Refer women to multidisciplinary breast clinics

EDITOR,—In their audit of patients with breast cancer treated in the Thames regions A M Chouillet and colleagues show and comment on wide variations in the treatment delivered in early 1990 to women of the same age and with disease of

the same stage.¹ Reasons for these variations are not hard to identify. Women are referred to many general surgical clinics, which lack the range of skill required for managing women with breast problems. Multidisciplinary breast clinics achieve a measure of uniformity and quality by working according to agreed protocols, whose implementation can be audited prospectively.

Referral to specialised breast clinics will not eliminate variation in treatment altogether because of differences of opinion between experts. Differences in treatment practices reflecting clinical uncertainty should act as a spur to participation in large randomised trials rather than to pressure for consensus guidelines. Where the adjuvant systemic treatment of early breast cancer is concerned, a worldwide meta-analysis carried out in 1992 showed conclusively that tamoxifen, chemotherapy, and ovarian suppression each reduce 10 year mortality in premenopausal and perimenopausal women.² About 5-10 extra women with early breast cancer are alive at 10 years for every 100 women treated with any of these modalities. Tamoxifen and chemotherapy are each effective in postmenopausal women. Tamoxifen has a relatively favourable toxicity profile and is simple to administer. Five years' treatment at 20 mg a day costs just under £200. It should therefore be considered for most women with early breast cancer, regardless of age. The optimal duration of tamoxifen treatment is being tested in the Cancer Research Campaign's trial and the "adjuvant tamoxifen treatment—offer more?" (aTTom) trial, and participation in these should be encouraged.

The overview carried out in 1992 also raised the possibility of appreciable further reductions in mortality from the addition of chemotherapy or ovarian suppression, or both, in women taking prolonged adjuvant tamoxifen. This hypothesis is being tested in the adjuvant breast cancer trial conducted by the United Kingdom Coordinating Committee on Cancer Research. Major trial groups in Britain are collaborating in this trial, which was launched last October. The trial's design is novel in offering a choice of randomisation options, depending on the uncertainty of the specialist, while retaining clinical freedom for the requirements of the individual patient. Several thousand women will be invited to participate to test the added survival gains of combined chemoendocrine treatment in patients taking tamoxifen. Associated studies aim to study the trade off between survival gains and side effects related to treatment.

JOHN R YARNOLD (Royal Marsden Hospital, Sutton, Surrey), JUDITH M BLISS (Institute of Cancer Research, Sutton, Surrey), MURRAY BRUNT (North Staffordshire Royal Infirmary, Stoke on Trent), HELENA EARL (Cancer Research Campaign Trials Unit, Queen Elizabeth Hospital, Birmingham), STAN KAYE (Beatson Oncology Centre, Glasgow), MALCOLM MASON (Velindre Hospital, Cardiff), JEAN MOSSMAN (United Kingdom Coordinating Committee on Cancer Research, London), TIM PERREN (St James's University Hospital, Leeds), MIKE RICHARDS (Guy's Hospital, London)

1 Chouillet AM, Bell CMJ, Hiscox JG. Management of breast cancer in southeast England. *BMJ* 1994;308:168-71. (15 January.)

2 Early Breast Cancer Trialists' Collaborative Group. Systemic treatment of early breast cancer by hormonal, cytotoxic, or immune therapy. 133 randomised trials involving 31 000 recurrences and 24 000 deaths among 75 000 women. *Lancet* 1992;339:1-15, 71-85.

Incomplete case notes hamper research

EDITOR,—As a research worker who has studied clinical decision making for 20 years and has also developed breast cancer, I was concerned by A M Chouillet and colleagues' paper.¹ My own treatment has followed the guidelines mentioned and has been fully discussed with me; from talking to fellow patients, I believe that this is common practice in Leeds.

My concern is on behalf of patients who are not receiving optimal treatment. Firstly, some doctors do not follow guidelines recommended by their peers. Here, it is perhaps easy to be overcritical. Guidelines are useful in general, and when they are not followed doctors should be prepared to defend their failure to comply. On the other hand, guidelines are rarely infallible, and experience in clinical research teaches the value of professional judgment, especially by experienced surgeons.

Far less easy to defend is the situation concerning case records. Clearly, those members of the medical profession who seemingly cannot be bothered to collect information and enter it into the case records need to change. In over half of the cases studied by Chouillet and colleagues the cancer could not be staged from the information recorded. This information not only would benefit the patient at the time but is vital for research if we are to improve the treatment and prognosis of this disease.

I support strongly the remarks of S J Karp that collation of data and comparison of centres are possible only if centres begin to record their data in a standardised way.² This applies to the whole of medicine: similar problems have been encountered in studies in acute abdominal pain, in which structured data collection is widely associated with improved diagnosis and decision making³ but young doctors often fail to collect data crucial to patients' management. If there are doctors who are unwilling to collect such information (much less follow guidelines produced to enable patients to benefit from recommendations of those doctors' peers) then what hope is there for audit and what hope is there for optimal patient care?

S E CLAMP

Clinical Information Science Unit,
University of Leeds,
Leeds LS2 9LN

1 Chouillet AM, Bell CMJ, Hiscox JG. Management of breast cancer in southeast England. *BMJ* 1994;308:168-71. (15 January.)

2 Karp SJ. Clinical oncology information network. *BMJ* 1994;308:147-8. (15 January.)

3 Adams ID, Chan M, Clifford PC. Computer-aided diagnosis of abdominal pain: a multicentre study. *BMJ* 1986;293:800-4.

4 de Dombal FT, de Baere H, van Elk PJ, Fingerhut A, Henriques J, Lavelle SM, et al. Objective medical decision making. Acute abdominal pain. In: Beneken JEW, Thevenen V, eds. *Advances in biomedical engineering*. Amsterdam: IOS Press, 1993:65-87.

Neurological symptoms may not be due to radiotherapy

EDITOR,—As surgeons and clinical oncologists who treat the axilla by surgery or radiotherapy for early cancer, we sympathise with the patients who have experienced major morbidity after loco-regional irradiation, whose cause Karol Sikora espouses.¹ Diagnosing brachial plexopathy induced by radiation is never easy. In a continuing but retrospective review of 773 women treated between 1979 and 1984 with radiotherapy after mastectomy we have identified only nine patients with symptoms suggesting brachial plexopathy, in two of whom radiation was identified as the cause. In five patients tumour was eventually identified as the cause, while in two others a final diagnosis cannot be made.

Our units are conducting a randomised trial of axillary node sampling (with axillary irradiation for women with positive nodes versus axillary clearance in which morbidity in the arm is being measured prospectively). From the response so far to a questionnaire completed by patients we cannot ascertain whether all neurological symptoms are secondary to radiotherapy. We would caution against any premature conclusions. Even computed tomography may not be helpful.² Long periods of careful follow up may be needed to exclude recurrence as a cause of this debilitating problem.

Sikora states that radiotherapy to the nodal areas

was probably unnecessary in many of the women. Patients who have had a full level III clearance (up to the apex of the axilla) will not require postoperative axillary radiotherapy. A clinical oncologist, however, is often faced with inadequate information about the pathological involvement of the axilla in patients referred for an opinion on the need for axillary irradiation.³ Commonly, this is due to inadequate sampling (when fewer than four nodes have been sampled) or occurs when no axillary procedure has been performed. A randomised trial in this unit showed that axillary sampling of at least four nodes provides an accurate assessment of the nodal status as established at a level III clearance.

It is also essential to follow up all patients to audit morbidity.

I H KUNKLER

Department of Clinical Oncology,
Western General Hospital,
Edinburgh EH4 2XU

U CHETTY

Edinburgh Breast Group,
Edinburgh Breast Unit,
Western General Hospital,
Edinburgh EH4 2XU

A RODGER

Department of Radiation Oncology,
William Buckland Radiotherapy Centre,
Alfred Hospital,
Prahan,
3181 Victoria,
Australia

1 Sikora K. Enraged about radiotherapy. *BMJ* 1994;308:188-9. (15 January.)

2 Cooke J, Powell S, Parsons C. The diagnosis by computed tomography of brachial plexus lesions following radiotherapy for carcinoma of the breast. *Clin Radiol* 1988;39:602-6.

3 Chouillet AM, Bell CMJ, Hiscox JG. Management of breast cancer in south east England. *BMJ* 1994;308:168-71. (15 January.)

4 Steele RJC, Forrest APM, Gibson T, Stewart HJ, Chetty U. The efficacy of lower axillary node sampling in obtaining lymph node status in breast cancer. *Br J Surg* 1985;72:368-9.

5 Rodger A. Comment on the role of axillary dissection in operable breast cancer. *Breast* 1992;1:48-9.

Attack on radiotherapy too simplistic

EDITOR,—Though I agree with Karol Sikora that patients who have been damaged by radiation need a sympathetic hearing,¹ I believe that several points need to be borne in mind. The recent publicity campaign by Radiotherapy Action Group Exposure has encouraged patients who have problems after radiotherapy to assume that these are due to radiation damage and to contact the organisation for a circular letter to send to their member of parliament, demanding an investigation and compensation. We therefore need an independent assessment of these patients to verify the diagnosis and exclude other conditions such as recurrent cancer, severe cervical spondylosis, or other neurological conditions causing similar symptoms. Probably some of the injuries to the brachial plexus are due to poor technique of delivery of radiotherapy and not simply a function of dose. A relatively high dose per fraction will be magnified by faulty technique, increasing the likelihood of injury.

Simplistic statements on television implying that half the radiotherapists in Britain give the wrong dose of radiation to their patients with breast cancer do nothing to maintain confidence in our ability to give effective treatment and fail to educate patients on the complexities of the issues. Sikora also gives the impression that he does not understand the concept of biological equivalence of radiation doses.

Much would be gained from an independent confidential inquiry (with power to insist on disclosure of medical records) examining details of dose and technique. Once the inquiry had made recommendations on preventing damage to the brachial plexus the allocation of resources would have to be reviewed to allow radiotherapy departments to meet the recommendations.