GPs' prescribing is irrational, says Audit Commission

Irrational and inconsistent prescribing by British general practitioners costs the NHS over £400m a year, says a report by the Audit Commission published this week. There are, it says, few limits to what and how much a general practitioner may prescribe, and although family health services authorities are supposed to keep general practice spending on drugs within preset limits, last year all English family health services authorities failed to meet spending targets.

The commission criticises the least cost conscious general practitioners for relying too heavily on drug treatment for minor complaints just to keep patients happy, for prescribing drugs that do not work, for prescribing expensive formulations that have no advantage over cheaper alternatives, and for being too quick to use new expensive drugs when older, cheaper ones are as effective.

Drugs being handed out to patients unnecessarily include antibiotics, drugs that heal ulcers, laxatives, tranquillisers, anti-inflammatory drugs, and drugs to lower serum cholesterol concentration. Inappropriate spending on these alone costs the health service nearly £300m each year, according to the report. A further £72m would be saved if doctors stopped giving patients drugs of questionable value such as cough suppressants, nasal decongestants, anti-inflammatory creams, appetite suppressants, and drugs to increase blood flow to the brain in patients with dementia.

The commission blames hurried consultations for some doctors' expensive prescribing; it urges them not to give in to patients who demand drug treatment when more time-consuming counselling is what they need. It warns that, although issuing a prescription may save time by terminating the consultation, it may also be dangerous. Benzodiazepines are known to cause addiction, yet over 500 million days' supply of these drugs was dispensed in England and Wales during 1992-3.

Ian Bogle, chairman of the British Medical Association's general practitioners' committee, said that overworked general practitioners would not be able to improve their prescribing habits unless they were given more time. He added, "It's important to recognise that GPs in this country still prescribe at a much lower level than their counterparts in other European countries, and it cannot be ignored that public expectations have risen in recent years with the introduction of the patient's charter."

The Audit Commission wants GPs and pharmacists to liaise more on prescribing

PIERS CAVENDISH/IMPACT

The report criticises doctors who prescribe new and more powerful drugs just to play safe. For example, doctors are increasingly giving patients newer, more expensive antidepressant drugs without waiting for conclusive evidence that they are safer. Newer antidepressants cost the health service an extra £30m last year. Some general practitioners are also prescribing a new tranquilliser which costs over 300 times as much as more traditional drugs. The manufacturer claims that the new drug is safer, but the Audit Commission says that the evidence is flimsy.

In a response to the report the Association of the British Pharmaceutical Industry warned that doctors should not be discouraged from using new drugs. "There are many examples of medicines, such as aspirin, steroids, and β blockers, whose full use has been recognised only after many years of clinical use. Much progress is made in medicines through small incremental steps, and to discourage such advances would be against the patients' interests," the association said.

The association's director, John Griffin, added that inadequate prescribing was as expensive as overprescribing. "The current cost of treating coronary heart disease and strokes is running at more than £1.5 billion a year . . . yet three quarters of the population suffering from raised blood pressure receive

no medication to reduce their risk of heart attack or stroke," he said.

General practitioners spent £3.6m on drugs during 1992-3, a tenth of all NHS costs. The Audit Commission estimates that nearly half these costs are a knock on effect from hospitals' prescribing habits. Drug companies sell drugs to hospitals at discount prices, encouraging hospital doctors to prescribe. Once a patient is discharged, the same drug prescribed by a general practitioner costs more, but a general practitioner must prescribe it or take responsibility for changing the specialist's treatment. Treatment with some drugs, such as the antiulcer drug ranitidine, costs twice as much in the community as in hospital. The report urges hospital pharmacies to consult general practitioners and family health services authorities before advising hospital doctors about prescribing.

The commission concludes that general practitioners should be more rational in the way they prescribe. They should take more time with patients, use the cheapest drugs that are effective and safe, agree with their colleagues locally about how to treat common conditions like asthma, and prevent serious—and expensive—disease by using more preventive drugs.—ALISON TONKS, *BMJ*

A Prescription for Improvement. Towards More Rational Prescribing in General Practice is published by HMSO, price £11.

Headlines

Americans turn against Clinton health plan: Eight out of 10 people surveyed in a recent poll thought that the quality of their health care would decline if President Clinton's health plan became law. The results may be affected by the advertising campaign mounted against the plan.

Europe falls behind in drugs research: The European Commission has warned that Europe's drug companies are not spending enough on research and development, that production is too fragmented, and that companies have not yet responded to the single market and have been left behind in the exploitation of biotechnology.

Ozone levels have declined: The levels of ozone, which acts as a filter against ultraviolet radiation, have declined by about 10% since 1969 but are expected to begin to rise soon because of the reduction in the use of ozone depleting chlorofluorocarbons, according to a report given at a meeting of the American Association for the Advancement of Science.

Intravenous immunoglobulin recalled: The maker of intravenous immunoglobulin, Baxter Healthcare Corporation, has stopped making Gammagard and Polygam and issued a worldwide recall of these products after three patients in Spain and two in Sweden were diagnosed as having hepatitis C after receiving Gammagard.

Mrs Bottomley admits London bed shortage: The British secretary of state for health told the Commons health committee that in some parts of London there might be "insufficient acute beds" for psychiatric patients. She announced that an extra 137 medium secure beds would be opened.

BMA calls for public inquiry into Allitt case: The BMA's council has supported the demand for a public inquiry into the case of Beverly Allitt, the nurse who was convicted of murdering four children in Grantham. The government inquiry heard evidence in private.

NHS should supply antismoking aids: The independent British think tank the Policy Studies Institute recommends that antismoking aids should be available on NHS prescription and that some of the money raised by tobacco tax should be used for community based health education.

Germany hails health reforms as a success

Germany is congratulating itself on having successfully reformed its public health insurance scheme after figures released last week showed a record surplus of DM10.2bn (£4bn) in 1993. This is in sharp contrast to 1992, when the scheme made a loss of DM9.4bn (£3.7bn). In western Germany the amount spent on health per person fell by 1.3% in 1993 while revenue from health insurance contributions rose by nearly 4%. In eastern Germany health expenditure rose by nearly 12% but revenue increased by nearly 17%.

Horst Seehofer, minister for health, said that the "impressive" savings were a result of the health reforms brought in at the beginning of 1993. He called the health reforms—the second within four years in Germany—a success. He insisted that they had not caused a reduction in the quality of medical care and denied that the system was providing "cheap medicine," as critics had predicted in 1992.

The bulk of the savings was made by cutting drug expenditure, which had risen by 9% in 1992. The latest figures show a reduction of nearly 20% in the amount spent on drugs last year. The reforms introduced fixed price lists for drugs, contributions by patients of up to DM10 (£4) per prescription, and generic prescribing for doctors. A total of DM10bn (£1.97bn) has been saved in drug costs alone.

The only health costs that increased in 1993 were those of patient transport and ambulance services, which went up by nearly 14%. This could be due to increases in fuel tax, but Seehofer said that he would take action to limit further rises in the cost of these services.

Seehofer has already urged health insurance companies to pass the savings on by lowering premiums. Germans spend 13% of their income on health insurance.

The main complaints have come from pharmacists: profits from sales of drugs have fallen by a third, and in Bavaria last year 21 chemist shops closed, most of their owners citing economic reasons.—HELMUT L. KARCHER, freelance journalist, Munich

France examines its poor record for using condoms

France lags behind most other countries in the European Union when it comes to using condoms. A survey by Agence Française de Lutte Contre le Sida—a French agency that fights AIDS—shows that the number of condoms used by Europeans varies from 1.7 to 3.8 per person per year and that the popularity of condoms is not related to cost.

Spain uses the most condoms even though they cost 60p on average. In 1992 about 150 million condoms were used in Spain—an average of 3.8 per person. In Britain, where the average cost of a condom is about 35p, a

total of 152 million were sold, representing 2.7 condoms per person per year. In France, where the cost averages 44p, two "préservatifs" are used per person per year. The cost of condoms is highest—71p—in the Netherlands, where their use is lowest—1.7 per person per year.

The study tries to explain why condoms are not popular in France. Legislation adopted at the beginning of the century made it illegal to advertise contraceptives, and this legislation was repealed only in 1987. Condoms had practically stopped being used in France by the end of the 1960s, having been replaced by oral contraceptives. In addition, the use of condoms in France became associated with prostitution and illicit sex. In 1970 condoms were used by 26% of people using contraception in Britain; in France the proportion of 22% was reached only last year. French health authorities are now trying to encourage the use of condoms to prevent HIV infection, and Philippe Douste-Blazy, minister delegate for health, has recently launched the so called "opération préservatif jeune" (contraceptives for the young), making the device—also known as "la capote anglaise" (the English coat)—available in pharmacies for Fr1 (about 15p).

Condoms are generally bought in pharmacies: 80% in Spain, 75% in Italy, 53% in Britain, and 48% in France. Only in the Netherlands and Germany do automatic coin machines account for sizeable sales.

The study notes that approaches to promote the use of condoms have changed in the past few years. The "dramatic approach" has shifted to a more humorous one. About 2.8 billion condoms are produced every year worldwide. According to the survey, Japan is the greatest consumer, using 58% of that total.—ALEXANDER DOROZYNSKI, medical journalist, Paris

Japan is the largest condom consumer but this anti-AIDS poster was rejected as offensive by transport authorities

ASSOCIATED PRESS

US attempts to stop its teenagers smoking

About one in eight teenagers—three million young people—smoke in the United States, and the surgeon general, Dr Joycelyn Elders, has called for a ban on advertisements that encourage teenagers to smoke. She has asked the Federal Trade Commission to stop the Joe Camel and Josephine Camel campaigns, which make smoking look glamorous to young people. American children are said to recognise Joe Camel almost as easily as Mickey Mouse.

There is a children's version of the surgeon general's report, which is intended to help children resist pressures to start smoking. And the American Medical Association has sent primary care physicians 240 000 copies of its guidelines for helping people to stop smoking.

About 3000 teenagers take up smoking every day. Between 1974 and 1984 the percentage of high school seniors (17 and 18 year olds) who smoked fell, but it is now increasing by about 1% a year, according to Dr Michael Eriksen, director of the Office on Smoking and Health at the Centers for Disease Control and Prevention.

An editorial in the *Journal of the American Medical Association's* annual antismoking issue said that one in four female high school seniors had smoked in the past month and 17% of both male and female seniors smoked daily. "If you make it to 20 without smoking chances are you won't start," the editor of *JAMA*, Dr George Lundberg, suggested.

Cigarette advertising has been banned on television and radio for many years, but sporting events seen on television commonly show evidence of tobacco sponsorship. Advertising in newspapers and magazines is permitted, and cigarette advertising is common in women's magazines. In the late 1960s and early 1970s advertising campaigns targeted at young women led to greatly increased sales of women's brands of cigarettes.

Now the manufacturer of Camel cigarettes, R J R Reynolds, has launched a major advertising campaign in women's magazines focusing on Josephine Camel, a sexy young woman pictured in exciting social situations. The vice president of the Robert Wood Johnson Foundation, Nancy Kaufman, pointed out in *JAMA* that the prevalence of smoking among women has risen since the late 1960s, although it has been falling among men since the late 1950s.

Dr Elizabeth Klonoff, professor of psychiatry at California State University in San Bernardino, reported in *JAMA's* antismoking issue that cigarettes are widely and illegally sold to children. All states ban the sale of cigarettes to people under 18. But in a survey of over 200 convenience stores and small groceries Dr Klonoff found that more than half were willing to sell single cigarettes to both adults and children. The stores charged 20-25 cents (14-16p) per cigarette, while the price of a packet of 20 was \$2-3 (£1.33-2).

The director of the Centers for Disease Control and Prevention, Dr David Satcher,

"Progress has stalled" on stopping Americans smoking

said, "Tobacco use is the single most preventable cause of death—420 000 deaths a year are due to smoking. In 1964, the year of the first surgeon general's report, 45% of Americans were smokers. Now only 25% are smokers, but there has been no notable decrease in recent years. Progress has stalled."

President Clinton's proposals for health reform initially included a \$2 (£1.33) tax for each packet of 20 cigarettes to discourage smoking. This has now been halved to the more politically acceptable \$1 (66p).—JANICE HOPKINS TANNE, contributing editor, *New York Magazine*

Memo suggests Upjohn concealed Halcion side effects

An investigation by the United States Food and Drug Administration (FDA) into possible misconduct by Upjohn, makers of the sleeping pill Halcion (triazolam), "strongly suggested" that the company knowingly concealed side effects in an early clinical trial, according to an FDA memorandum. Subjects in protocol 321, a study of 28 healthy prisoners for 42 days in 1972, experienced more adverse reactions than were shown in the summary that the pharmaceutical giant submitted to the FDA in its application for approval. The true incidence of side effects emerged only in 1991, when the original data came to light during a court case in the US over Halcion.

The memo of December 1992, obtainable under the US Freedom of Information Act, shows that the FDA considered referring the case to the US Department of Justice for an investigation by a grand jury. But it decided not to go ahead because the study was considered to be of marginal importance in Halcion's approval for marketing.

The FDA decided to investigate the circumstances surrounding the submission of the report on protocol 321 after Britain banned the drug in October 1991. The memo says that Halcion was banned in Britain "as a result of the information pertaining to study 321." In its letter to Upjohn announcing the suspension of its product licence the United Kingdom's Committee on Safety of Medicines said that the licence would never have been granted had adverse reactions been reported accurately.

The errors emerged when an American woman, Ilo Grundberg, sued Upjohn, claiming that she had shot her mother under the influence of Halcion. Ian Oswald, emeritus professor of psychiatry at Edinburgh and an expert witness for Mrs Grundberg, discovered the discrepancies when he compared the summary that the firm had submitted to the FDA with individual case reports.

Upjohn reported the errors to the FDA and drug regulators worldwide, blaming them on "transcription error." According to the FDA's memo the firm did not reanalyse the data, claiming that this would not change the frequency of side effects. In fact, the summary reported four events of nervousness or anxiety and two incidents of paranoia in the 28 prisoners, while case report forms show at least 70 events of nervousness or anxiety and seven prisoners with paranoia. Upjohn is suing Professor Oswald, a long time critic of Halcion, and the BBC for libel in the High Court in London. The trial, over a BBC *Panorama* programme and an article in the *New York Times* which quotes Professor Oswald, started at the end of January and is expected to last four months (22 January, p 221).

The memo, from George Prager, consumer safety officer in the FDA's division of scientific investigations, says: "FDA's investigation strongly suggests that the firm was aware that the report was inaccurate and incomplete but continued to withhold notification to the FDA that the report was flawed. Study 321 took place 20 years ago and because of the expired statute of limitations

any case developed would be predicated on evidence of a conspiracy to cover up the data from study 321. We were unable to develop such evidence from the current inspection. However, it was felt that case may be appropriate for an investigative grand jury."

But officials decided not to take further action because the facts withheld would not have been material to Halcion's application for approval. The memo cites the unusual nature of the study subjects—prisoners—and the high dosage, which was never seriously considered for marketing in the US.

The libel trial has focused attention again on protocol 321 and other studies of Halcion over which question marks hang. Documents which the company fought for years to keep secret are emerging in the trial and through lawsuits in the US.

Four of the investigators in clinical trials of the drug have been disqualified by the FDA. One, a Mississippi doctor, Samuel Fuerst, was disqualified in 1984 for professional misconduct. The FDA found no evidence that the people he had listed as patients had ever taken the drug. Upjohn agreed to strike the results of the study from its database in 1984, but the company's president, Ley Smith, admitted last month that it had submitted Dr Fuerst's results to British drug regulators in 1991 without mentioning the FDA's action seven years before.

Upjohn refused to comment until the end of the libel trial.—CLARE DYER, legal correspondent, *BMJ*

SALLY AND RICHARD GREENHILL

French researchers say riluzole may slow down motor neurone disease

Riluzole may help survival in motor neurone disease

A French research team has reported a small gain in the understanding of amyotrophic lateral sclerosis. But although there was widespread news coverage about it in Europe and the United States, scientists remain sceptical.

Amyotrophic lateral sclerosis affects upper motor neurones, and little more is known about it today than when it was first described by Charcot in 1865. It is a rare disease, striking about 1 in 100 000 adults a year, and is progressive, destroying neurones in the cortex, brain stem, and spinal cord. It is inevitably fatal, killing patients in middle age usually within four years of onset. It has claimed the lives of several well known people, such as the actor David Niven and the baseball player Lou Gehrig, whose name is the common eponym for the disease in the US. In Britain the disease is often called motor neurone disease and in France *maladie de Charcot*.

In the 3 March edition of the *New England Journal of Medicine* (*NEJM*) a group of French researchers reported that the drug riluzole, a glutamate antagonist, slowed the progression of disease in one subset of patients, those whose disease began in the brain as opposed to the limbs. Riluzole was chosen for the study because about 5% of

cases of amyotrophic lateral sclerosis are familial, and 40% of patients with the familial disease have high levels of glutamate, which suggests a toxic effect on neurones.

In a multicentre randomised clinical trial of 155 patients with the syndrome the researchers found no difference in survival at one year in 77 patients given riluzole compared with 78 given a placebo. The researchers reported "unexpectedly" that among patients with amyotrophic lateral sclerosis of bulbar onset survival at one year was 73% among the 15 treated patients and 35% among the 17 patients who received a placebo.

The finding was widely reported and was lauded by another researcher who is directing a larger study of riluzole, Dr Jeffrey Rothstein of Johns Hopkins University.

But others, including the writer of an editorial in *NEJM* (also 3 March), were less convinced. Dr Lewis P Rowland of Columbia University wrote that amyotrophic lateral sclerosis of bulbar onset "has not been previously postulated" as being a different disease from the disease of limb onset, and thus the finding "runs counter to clinical logic." He was worried that the report may make it "even more difficult to convince patients to enter studies with a placebo group."

Others say that the study may represent a case of "beating the data till they surrender." Statistician Ken MacCrea of Charing Cross and Westminster Medical Schools pointed out that too much "Subgroup analysis," especially when subgroups are small, represents a common problem in clinical trials.

Randomisation works well for big, long term studies because it minimises differences between groups, he said. But it does not ensure that no differences will occur, especially when groups are small. For example, it has been reported that age and duration of disease are predictors of survival in amyotrophic lateral sclerosis. Although the large

groups of patients given riluzole and a placebo were similar, when the 32 patients with disease of bulbar onset were compared the treated group was younger, had had the disease for a shorter time, had better scores of bulbar function, and had better scores on a "stiffness scale." These chance differences, not necessarily the drug, may have accounted for the better outcome.

The French study was funded by riluzole's manufacturer, Rhône-Poulenc Rorer. The drug remains a research agent and cannot be used clinically either in Europe or the US.—JOHN ROBERTS, *BMJ* North American editor

Doctors' incomes in Australia spark row

A controversial study showing that some specialist doctors received more than \$A700 000 (£280 000) a year from the public Medicare system has sparked a big row between doctors and the federal government. Senator Graham Richardson, the health minister, told parliament that some of the nation's surgeons were "greedy" and "exploited" their patients by charging excessive fees. He warned that any increases in the health budget would not be used to prop up the incomes of specialist doctors. "The great majority of doctors—more than two thirds—choose not to exploit their market power, but one third does. A minority of them exploit their position and do so shamelessly," he said.

The Australian Medical Association (AMA) accused the senator of indulging in the "politics of envy." The federal president, Dr Brendan Nelson, said that greed was not unique to doctors, who should not be measured by their earnings. He said that top specialists had the right to earn such amounts

considering the hours worked, personal sacrifices, and overheads.

The war of words comes as the two sides fight over the reforms of the private health system. The reforms are aimed at reducing the burden on the public health system by stabilising the number of Australians with private health cover at 40%. But Dr Nelson has branded Senator Richardson's reforms as "unworkable and discriminatory."

The study, conducted by the University of New South Wales, disclosed details of doctors' gross incomes from Medicare for the first time. The data came from evidence given to the New South Wales Industrial Commission by the New South Wales branch of the AMA during hearings on the pay of visiting medical officers in 1992. Thousands of general practitioners and hundreds of specialists gave evidence. The authors of the study found that while only one in 100 general practitioners received Medicare payments of over \$A300 000 (£120 000) in 1991-2, over 7% of specialists earned more than that. Most general practitioners had an annual income from Medicare of \$A40 000-120 000 (£16 000-£48 000) while specialists generally earned over twice that amount.

Over 40 doctors in New South Wales collected more than \$A700 000 (£280 000) each during 1991-2. One of the authors of the study, Professor James Lawson, called for doctors earning large amounts to be publicly accountable. He said that it was "very difficult for a skin specialist to justify earning \$A700 000 from the public purse." Most of the doctors receiving such amounts worked as cardiologists, ophthalmic surgeons, obstetricians, and gastroenterologists.

The day after the report's release Senator Richardson told parliament that one third of surgeons charged more than 30% above the Medicare benefits schedule, which patients had to meet out of their own pocket. He said it was "time for the AMA to acknowledge that greed has been the major factor in killing private health insurance," adding that 284 000 Australians had dropped their private cover last year alone.

The AMA claim that the medical benefits schedule has not kept up with inflation. "I defend the right of top doctors in Australia to earn that money, provided they are caring for their patients and working very long hours," said Dr Nelson.

The progressive Doctors' Reform Society supported the study's findings, saying that some specialists were guilty of overcharging. The society's president, Dr Con Costa, said that the situation existed because of the college's "restrictive trade practices" in limiting the number of specialists who were trained. The National Association of Specialist Obstetricians and Gynaecologists, however, defended its members against allegations of overcharging. "We're particularly sensitive about the suggestion that anyone who practises obstetrics is overpaid when we are dramatically underpaid," said the New South Wales chairman of the association, Douglas Reed. He said that over 40% of doctors in the specialty earned less than \$A120 000 (£48 000) a year from Medicare.—CHRISTOPHER ZINN, *Guardian* correspondent, Australia

Phase one trials can exploit terminally ill patients

Fears are emerging in the US that terminally ill patients who take part in phase one trials of chemotherapy are being exploited. A recent pilot study carried out by the department of haematology and oncology at Chicago University's centre for clinical medical ethics suggests that patients gave their informed consent because they believed in the possibility of therapeutic benefit. The 27 patients in the study were also asked whether, as is required, other options besides participation in the trial were discussed during the informed consent process; two thirds said no.

The purpose of phase one trials of chemotherapy—performed in the US mostly with solid tumours—is to test the toxicity of increasing doses of an experimental agent. Clearly, if the drug is sufficiently toxic or the dose sufficiently high, then some subjects will suffer. The drug is tested until the so called maximum tolerated dose is reached. If subjects fail to respond to an experimental agent after about two cycles of treatment at their assigned dose level they are usually released from the trial and sent back to their referring physicians.

Patients were asked in interviews to state the purpose of the phase one trial that they were entering. Only a third of the 27 patients stated that it was to assess the effects of dose escalation. The other two thirds said that they didn't know or that the purpose was therapeutic benefit. Clearly, despite attempts at explanations, patients' motivation to get some therapeutic benefit is so overwhelming that it obliterates the memories of much of what they have been told about the trial's purpose, said Chris Daugherty, an oncologist who carried out the study.

There is a view that since terminally ill people who take part in these trials give their informed consent and will probably die within months anyway, any resulting ill effects or hastening of their death may be unfortunate but do not preclude such trials. Others feel that there is no alternative. Dwight Kaufman, deputy director of the division of cancer prevention at the National Cancer Institute in the US, said: "To give an experimental, possibly toxic agent that might manifest such adverse effects as leukaemia years later to a healthy person, or one cured of cancer, is unthinkable." But Kaufman also said "terminally ill patients must be made to understand the facts: we must stress to such patients that it is unprecedented for a cure to occur—that the probability of cure approaches zero."

George Annas, director of the medical ethics and law programme at Boston University Schools of Medicine and Public Health, thinks that these patients need special protection. "These patients do have something to lose," he said. "They may end up dying a more miserable death than otherwise."

Annas believes that the dual roles of the phase one investigator as physician and scientist present a problem: "It's difficult to separate these roles, and they can't believe that as physicians they would harm patients."

In the US, in recent years, Annas continues, "AIDS activists have demanded experimental agents and now we all are equating research with treatment, the justification being the consent process."

Daugherty believes that "it's possible for all patients to understand the purpose of such trials and to give full informed consent. Certainly only the ones who understand what's involved should participate in these trials. This means that patients must become more involved in the process." One way to do this, which he says "may not be feasible ethically or clinically," is to allow patients to

Not all patients know they are taking part in phase one trials

GEOFF TOMPKINSON/SPL

select the doses they want to take. Essentially, this means that the lower doses would be skipped, saving both time and money in a drug trial."—GAIL MCBRIDE, medical journalist, Chicago

Egypt considers care in community

Appalling conditions at one of Egypt's largest psychiatric hospitals has prompted the Ministry of Health to set up a committee of psychiatrists to decide the future of the country's main psychiatric institutions. Abbasseya Psychiatric Hospital, built in the grounds of a former palace in Cairo, originally served as a hospital for the slaves of King Khedive in 1880. Over the past 40 years, doctors say, the hospital has been deprived of funds and has steadily deteriorated. "Many patients do not have treatment at all," said Dr Raouf Nabil, a young doctor at the hospital. "The standard of living for patients needs

many changes; they need better food, treatment, and clothing." The hospital is also accused of not being large enough to accommodate its 3000 patients.

"The situation is bad—it's inhuman for anyone to have to live there even for a day," said Dr Nabil. The 400 nurses are mostly unqualified, and there have been accusations of patients being maltreated. There are frequently shortages of basic drugs, and violence is common. Of the 40 doctors on the staff, 34 are young residents in training. Dr Ahmed El Hady, a consultant at the hospital for 15 years, said that the hospital could be considered to be "Egypt's fourth pyramid."

Dr Al Sayed Al Kott, general director of the Mental Health Department at the Ministry of Health, was appointed director of Abbasseya Hospital last December. He says that the hospital is not suitable for modern psychiatric treatment but believes that it should remain open. Other psychiatrists also fear that there is not enough funding to look after patients in the community.

The Ministry of Health has said that it intends to build smaller hospitals to replace the large institutions, but so far their con-

struction is not occurring at the expected rate. Two years ago the Ministry of Health released 2000 of 3000 patients from El Khanka Psychiatric Hospital, and most have remained in the community.

The problems in Abbasseya Hospital stem from a crisis in psychiatry in general. Egypt has only 300-400 psychiatrists, most of whom are based in large, overcrowded hospitals run by the Ministry of Health. The hospitals lack basic drugs and are staffed by a handful of psychiatrists. Teaching hospitals offer a high standard of care but have few psychiatric beds. Less than one in 10 people can afford private psychiatric care.—MARA GULENS, freelance journalist, Egypt

Correction

Guy's Hospital will stop acute care

An editorial error occurred in this news story (19 February, p 496). It was stated incorrectly that a decision on the application for trust status by the Royal National Throat, Nose, and Ear Hospital had been deferred. This hospital in fact became a trust in April 1991.

Focus: Brussels

Combating drug trafficking across the single market

Suggestions for tackling the growing drugs epidemic are as varied as the professional backgrounds and political beliefs of those who confront the problem. Doctors, social workers, customs officers, police, and politicians all have different, and sometimes conflicting, remedies.

But drug trafficking and drug misuse are international; despite the many standpoints, the fight against drugs will probably be more effective with international co-operation. So the European Union seems to think, as it is taking advantage of the new powers granted in the Maastricht Treaty to act against drug misuse. More opportunities for coordinated action among the 12 European Union members, and possibly among 15 or 16 countries from next year, exist now than six months ago.

That cooperation runs right through the three main pillars of the European Union constructed by the new treaty. It includes extending the supranational activity which the European Union has gradually developed over the past four decades to cover public health, and drug dependence in particular. It also extends to more classic intergovernmental operations, such as national police and customs authorities combining their efforts against traffickers, and to foreign ministers devising strategies for targeting the flow of drugs into the union.

The basic tools for this cooperation are gradually being put in place. The Europol drugs unit, sanctioned at last October's summit in Brussels, officially opened for business in the Hague last month. At the sharp end of the fight against drugs, it offers a rapid information exchange system on international shipments of drugs. The unit will not be fully operational until October, but already it can point to drug seizures achieved by the new cooperation.

The new European Monitoring Centre for Drugs and Drug Addiction in Lisbon (5 March, p 609) is also part of the same spirit of cooperation.

The monitoring centre's role is strictly limited to gathering, processing, and providing reliable comparable information on the drugs phenomenon, drug addiction, and their consequences. In this new task, designed to provide data on which medical and political decisions can later be made, it will cooperate with Europol, the Council of Europe's own antidrug efforts through the Pompidou group, and the United Nations.

Last month the European parliament too turned its attention to drugs policy when Italian Green MEP Marco Taradash tried to win support for a rethink of existing policies. Most controversy focused on his view that possession for personal use of small quantities of prohibited drugs should not in principle be a criminal offence. But the real thrust of his report was that the existing antidrugs policy, based largely on United Nations conventions laying down a

blanket ban on drugs, had failed and that a new approach was necessary.

In the event the call for a change of policy and review of the UN conventions was rejected, but by only six votes out of the 249 cast. The parliament has no legal power to determine drugs policy, but politically its views are yet another input into the current debate.

Other elements involve the European Commission's increasing emphasis on public health and social policies; pan-European programmes by interior and justice ministers to plug loopholes that emerge as internal frontiers disappear in the single market; and growing awareness that the union's foray into establishing a common foreign and security policy could encompass the international drugs trade.

It is these various strands that the social affairs commissioner, Pdraig Flynn, also responsible for public health, is now trying to draw together into a coherent package. His staff are preparing a report analysing the various initiatives now on the table. When it appears in June or July it is likely to contain specific recommendations for constructing a global policy.

Some governments may not take kindly to the European Commission playing such an active part in drugs policy. But that is unlikely to deter Flynn. Last month he unveiled a similar report, designed to weave together the many different programmes in the even more politically sensitive subject of immigration into the union. —RORY WATSON, *the European*