not the drug of choice for prophylaxis for aircrew. Doctors who care for pilots should keep in mind the uncertainties surrounding mefloquine before prescribing it in preference to other well tried prophylactic drugs. Certainly, vibramycin and combinations of chloroquine and proguanil may result in problems, but they have the advantage of having stood the test of time.

J C MERRITT

Cathay Pacific Airways,
Hong Kong International Airport,
Hong Kong

1 White NJ. Mefloquine. *BMJ* 1994;308:286-7. (29 January.)
2 Bradley D. Prophylaxis against malaria for travellers from the United Kingdom. *BMJ* 1993;306:1247-52.

Alcohol histories taken from elderly people

EDITOR,—P C Naik and R G Jones claim that elderly people are more likely to hide their drinking than younger people.¹ There is no evidence to support this. The Framingham study on alcohol consumption and hip fractures, which looked at 5209 patients, found that differential underreporting of consumption by older women did not occur.² Other studies have shown that older people drink less than younger ones,^{3,4} but this is unlikely to be the reason behind Naik and Jones's finding that doctors are less likely to record an alcohol history in elderly people. A failure to record any alcohol history in 48% of patients indicates a failure of basic medical education rather than an underestimation of the importance of ingestion of alcohol by elderly people.⁵ Fortunately, the number of patients in their study was small and so, presumably, was the number of doctors.

Even careful estimates of alcohol intake are inaccurate, and the estimated alcohol content of a hot toddy will be even more unreliable. The conclusions drawn from a study of such a small sample of patients should be regarded with caution.

D THOMPSON

Wharfedale General Hospital,
Otley,
West Yorkshire LS21 2LY

1 Naik PC, Jones RG. Alcohol histories taken from elderly people on admission. *BMJ* 1994;308:248. (22 January.)
2 Felson DT, Kiel DP, Anderson JJ, Kannel WB. Alcohol consumption and hip fractures: the Framingham study. *Am J Epidemiol* 1988;128:1102-10.
3 Glynn R, Bouchard GR, LoCastro JS, Laird NM. Ageing and generational effects on drinking behaviours in men: results from the normative ageing study. *Am J Public Health* 1985;75:1413-9.
4 Goodwin JS, Sanchez CJ, Thomas P, Hunt C, Garry PJ, Goodwin JM. Alcohol intake in a healthy elderly population. *Am J Public Health* 1987;77:173-7.
5 Warriss JP. Alcohol and old people. *Br J Psychiatry* 1983;143:306-7.

Blackening of the oesophagus

EDITOR,—Hamid A Khan and David Owens illustrate a circumscribed area of black pigmentation of the mid-oesophageal mucosa of an 84 year old former coal miner with a history of pneumoconiosis.¹ Histopathological studies led to the pigment being identified as coal dust, but the authors are puzzled about the mechanism of production of the lesion.

I suggest that the patient suffered from pulmonary anthracosis (or possibly extreme anthracosis) with or without associated pulmonary tuberculosis; that the lymph nodes draining his lungs became much enlarged by fibrotic anthracotic lesions similar to those that develop in the lungs; and that one or more nodes underwent central liquefactive necrosis with perforation of the capsule followed by tracking of the resultant black fluid into the oesophageal wall. A node of the

subcarinal group may have been the source. This rare complication has more often resulted in perforation into the trachea, a bronchus, or an adjacent blood vessel (usually a pulmonary vein).²

G A K MISSEN

London SE3 7PH

1 Khan HA, Owens D. Minerva. *BMJ* 1994;308:546. (19 February.)
2 Symmers WStC. The lymphoreticular system. In: Symmers WStC, ed. *Systemic pathology*. Vol 2. 2nd ed. Edinburgh: Churchill Livingstone, 1978.

Rewarding units that exceed performance targets

EDITOR,—In his editorial on the Labour party's proposals for an alternative health policy Chris Ham asks "without competition [under the proposed new system] where will be the incentive to reward providers that exceed their performance targets?"¹ This implies that exceeding performance targets is currently rewarded. Yet a news item published in the same issue of the journal says that a survey by the Royal College of Surgeons has found that 44% of surgical divisions had had their activity reduced²—presumably because they had reached their performance targets. I am sure that these surgeons (and anaesthetists and theatre staff, not to mention their patients) do not feel rewarded by the current system.

HILARY AITKEN

Alexandra Hospital,
Redditch B98 7UB

1 Ham C. An alternative for the NHS. *BMJ* 1994;308:485-6. (19 February.)
2 Headline. *BMJ* 1994;308:492. (19 February.)

Junior doctors' training

EDITOR,—P R Eldridge holds several of the more common misconceptions about training in other countries in the European Union.¹ Europe is not a single entity, and the quality of postgraduate medical training varies enormously. Eldridge's statement that "continental Europeans are not trained to a standard equivalent to that in Britain" is not only inaccurate but offensive. Some European countries have better health outcome measures than Britain. This must be due at least partly to superior training of health care professionals.

Eldridge believes that the processes in Britain to assess both trainees and training programmes result in high quality training. Eldridge may therefore be surprised to learn that several other European countries also have such systems. A recent survey by the Permanent Working Group of European Junior Hospital Doctors showed that in some countries in the European Union these systems are further advanced and use better developed curriculums and criteria than those in Britain (Binyon, paper in preparation).

Eldridge then moves on to the quantity of operations performed. Surely we should not mistake quantity for quality. One operation properly supervised must be of greater educational benefit than several unsupervised operations performed by a tired doctor in the early hours of the morning. We were particularly interested in the assertion that a trainee from France had operated on only 25 patients. This suggests that during a minimum of five years' training the trainee had operated on only five patients a year—something that no health care system is likely to allow. As mentioned above, however, if these 25 operations were of sufficient diversity and were used maximally for educational purposes this trainee might be better trained than Eldridge.

Eldridge's arguments to justify persisting with an average 83 hours a week ignores patients' safety and quality of care. It has been found in law that excessive working hours are associated with increased mortality and morbidity.²

Eldridge also suggests that the main determinant of the time currently spent in a training grade is the amount of experience required. We suggest that the greatest determining factor is the ratio of trainees to consultant posts and therefore the career opportunities.

Eldridge assumes that if the average working week is reduced then the total time in a training grade must be increased by the same proportion. This suggests that at present trainees are working for all of each duty period and that there are no inefficiencies of training (for example, certain operations being performed well in excess of the number of times required to acquire proficiency, etc). The statement that "evidence exists that long hours on call are necessary to train in a short period" is extraordinary. We are unaware of any such evidence.

Harmonisation of training in Europe should be the ultimate aim, but we should not be so arrogant as to believe that we cannot learn as much from other systems as they can from us.

SHARON E BINYON
GRANT J INGRAMS

Birmingham B29 4LS

1 Eldridge PR. Junior doctors' hours. *BMJ* 1994;308:417. (5 February.)
2 Taylor D. Primary care. In: Maxwell R, ed. *Reshaping the National Health Service*. Oxford: Policy Journals/Transaction Books, 1988.

Sharp advertising

EDITOR,—Charles Warlow asks how much Auckland Hospital gained from the huge illuminated advertisement for Sharp Electronic.¹ Soon after the advertisement was erected it was reported that the income received would allow one extra staff nurse to be employed for the next nine years.²

IAIN WAKEFIELD

Drury Lane Surgery,
PO Box 26, Drury,
Auckland,
New Zealand

1 Warlow C. Sharp advertising. *BMJ* 1994;308:472. (12 February.)
2 *New Zealand Herald* 1992 April 8:5.

ARTICLES — OR CLUSTERS OF ARTICLES — ATTRACTING THE MOST LETTERS FOR PUBLICATION IN 1993	
Subject of article (date)	Number of letters received (published)
Natural family planning (18 September)	50 (25)
Cardiopulmonary resuscitation (12 June)	32 (14)
Coronary bypass surgery in smokers (17 April)	27 (15)
Informed consent for clinical trials (6 November)	23 (15)
Low back pain (3 April)	21 (7)
Repetitive strain injury (20 November)	21 (15)
Management of patients in persistent vegetative state (12 June)	19 (10)
Risk of sepsis after splenectomy (27 November)	19 (5)
Consultant jobs for women (16 October)	15 (8)
Selective serotonin reuptake inhibitors (13 March)	14 (9)