bemused clinicians to assess the validity of recommendations that differ between the two sets of guidelines. This may require a longer reference document or tables of evidence, but these do not exist for either guideline. The longer version of the WHO-International Society of Hypertension's guidelines is no more explicit about how evidence was chosen or weighted.

What is the responsibility of a medical journal in assessing clinical guidelines submitted for publication? Is peer review sufficient? Perhaps journals should use one of the tools for appraising guidelines developed by the American guidelines industry or one of the shorter versions being developed in Britain. With the current profusion of clinical guidelines, the *BMJ* has a role in encouraging higher standards of development.

GENE FEDER

Hackney Collaborative Clinical Guidelines Project,
Department of General Practice,
St Bartholomew's Medical College,
London EC1M 6BQ

1 Sever P, Beevers G, Bulpitt C, Lever A, Ramsay L, Reid J, *et al*. Management guidelines in essential hypertension: report of the second working party of the British Hypertension Society. *BMJ* 1993;306:983-7.

2 Subcommittee of WHO/ISH Hypertension Liaison Committee. Summary of 1993 World Health Organisation-International Society of Hypertension guidelines for the management of mild hypertension. *BMJ* 1993;307:1541-6. (11 December.)

*Medical journals certainly have a responsibility to assess rigorously the clinical guidelines that are submitted to them. The International Committee of Medical Editors has this issue on the agenda for its meeting in Oslo in June, and we at the *BMJ* are trying to raise the quality of our assessment of guidelines. The most crucial issue is how the group producing the guidelines has assembled and assessed evidence in reaching its conclusions. Another important question is whether all the detail of how the evidence was assembled and assessed should be included in the published guidelines.—ED, *BMJ*

Prenatal screening for trisomy 18

Should not be contemplated

EDITOR,—K Spencer and colleagues' claims for prenatal detection of trisomy 18 by measurement of maternal serum α fetoprotein and free β human chorionic gonadotrophin concentrations are impressive.¹ Detection of 50% of cases for a false positive rate of only 1% seems to compare favourably with the detection rate for Down's syndrome when similar techniques are used, which is 70% for a false positive rate of 5%. Unfortunately, the authors fail to emphasise the importance of the relative incidence of the two conditions at birth before concluding that screening for trisomy 18 should be introduced.

The natural incidence of Down's syndrome at birth is approximately 12.6/10 000 births.² Among 10 000 pregnant women a 70% sensitivity would result in 8.8 cases being detected at the cost of 500 amniocenteses (5% of 10 000). This means that one case of Down's syndrome is detected for every 57 amniocenteses performed. The incidence of trisomy 18 at birth is 1.3/10 000 births. A sensitivity of 50% would detect 0.65 cases per 10 000 women tested at a cost of 100 amniocenteses (1% of 10 000). For each case of trisomy 18 detected, therefore, 154 women would have to have had amniocentesis. Thus a screening programme would cause the abortion of at least as many normal fetuses as it would detect cases of trisomy 18.

In many places it is still undecided whether screening for Down's syndrome is worth the disbenefits for the prospective parents. To my mind, the decision is clearer for screening for trisomy 18: screening should not be contemplated

until the predictive value of the test is considerably improved.

TIM DAVIES

Warwickshire Health Authority,
Warwick CV34 4DE

1 Spencer K, Mallard AS, Coombes EJ, Macri JN. Prenatal screening for trisomy 18 with free β human chorionic gonadotrophin as a marker. *BMJ* 1993;307:1455-8. (4 December.)

2 Cuckle H, Nanchahal K, Wald N. Birth prevalence of Down's syndrome in England and Wales. *Prenat Diagn* 1991;11:29-34.

Most die before or just after birth

EDITOR,—K Spencer and colleagues recommend biochemical screening for trisomy 18 on the basis of a detection rate of 60-65% for a false positive rate of 1%.¹ Although trisomy 18 is the second most common autosomal trisomy, its incidence at birth is between one in 6000 and one in 8000 (whereas that of trisomy 21 is one in 770).^{1,2} Consequently the number of karyotyping procedures necessary to identify a single case of trisomy 18 at term (the positive predictive value of an invasive prenatal diagnostic procedure) will be correspondingly greater. For example, assuming a detection rate of 60% for a false positive rate of 1% and an incidence at birth of one in 7000, in 70 000 pregnancies there will be 10 cases at term and 700 amniocenteses would be performed to detect six of them. Thus 117 (700/6) invasive procedures would be necessary to detect a single case of trisomy 18. The equivalent ratio for trisomy 21 at term is roughly 59 to one (adjusting the 40 to one ratio observed at 16 weeks³). In addition, most (70%) fetuses with trisomy 18 die between 16 weeks and term, compared with a minority (30%) of fetuses with trisomy 21.⁴ Liveborn infants with trisomy 18 rarely survive for longer than a year. Thus biochemical screening for trisomy 18 is clearly less efficient than that for trisomy 21 and would give rise to unnecessary anxiety in large numbers of women with normal pregnancies.

M C M MACINTOSH

T CHARD

Department of Reproductive Physiology,
St Bartholomew's Hospital,
London EC1A 7BE

1 Spencer K, Mallard AS, Coombes EJ, Macri JN. Prenatal screening for trisomy 18 with free β human chorionic gonadotrophin as a marker. *BMJ* 1993;307:1455-8. (4 December.)

2 Hook EB, Woodbury DF, Albright SG. Rates of trisomy 18 in livebirths, stillbirths, and at amniocentesis. *Birth Defects* 1978;15(5c):81-93.

3 Royal College of Obstetricians and Gynaecologists. *Report of the working party on biochemical markers and the detection of Down's syndrome*. London: RCOG, 1993.

4 Hook EB. Chromosome abnormalities and spontaneous fetal death following amniocentesis: further data and associations with maternal age. *Am J Hum Genet* 1983;35:110-6.

Ultrasound scanning would detect abnormalities

EDITOR,—K Spencer and colleagues' paper on biochemical markers for trisomy 18 prompts two comments.¹ Firstly, Twining reviewed the literature on markers of trisomy 18 evident on ultrasound scanning and concluded that 85% of fetuses with trisomy 18 have structural abnormalities that should be detected with careful scanning.² As most women have detailed scans at 19 weeks, over 80% of fetuses with trisomy 18 would be identified as compared with only 50% with biochemical screening. Secondly, for screening for both trisomy 18 and trisomy 21, it would be useful if ultrasound appearances, age, and biochemical markers could be combined in an algorithm to give a woman a personalised overall risk factor for trisomy rather than ever more conflicting advice.

KATE HARDING

Department of Obstetrics and Gynaecology,
Charing Cross and Westminster Medical School,
London W6 7DQ

1 Spencer K, Mallard AS, Coombes EJ, Macri JN. Prenatal screening for trisomy 18 with free β human chorionic gonadotrophin as a marker. *BMJ* 1993;307:1455-8. (4 December.)

2 Twining P. Choroid plexus cysts—a conservative approach. *British Medical Ultrasound Society Bulletin* 1993;Feb:43.

Conflict of interest

Sometimes consultancies cannot be disclosed

EDITOR,—Richard Smith rightly says that disclosure is a good remedy for conflict of interest. However, academics who choose to advise industry usually have to sign a secrecy agreement to ensure that commercially valuable information is not passed to a competitor. Sometimes the fact that firm A has employed an expert in field B as a consultant is itself commercially sensitive, so the secrecy agreement specifies that the existence of the consultancy should also be kept secret. For obvious reasons I do not know how often this happens, but I have twice had such an approach myself. I think it is important that academics who intend to contribute to the scientific literature as authors, editors, or referees should not accept consultancies which they cannot publicly disclose.

JOHN GARROW

European Journal of Clinical Nutrition,
London EC1M 6BQ

1 Smith R. Conflict of interest and the *BMJ*. *BMJ* 1994;308:4-5. (1 January.)

Speakers should declare financial support

EDITOR,—With regard to avoiding potential conflicts of interest,¹ I was impressed recently by a procedure used at a satellite meeting of the American Heart Association. The meeting was recognised for the continuing education that American doctors are required to undertake for reaccreditation. At the start of the meeting every participant was given a sheet of paper listing the speakers and any financial support they received that might lead to a conflict of interest. For example, it was possible to tell which speakers were receiving financial support directly or indirectly from particular pharmaceutical companies. Clearly, for meetings on contentious issues concerning prescribing it would be helpful to the audience to have such information and to be able to judge, for example, the balance of speakers chosen to talk on the topic. Surely such a procedure should be incorporated into the requirements for accreditation of postgraduate educational courses in Britain.

GRAHAM WATT

Department of Public Health,
University of Glasgow,
Glasgow G12 8RZ

1 Smith R. Conflict of interest and the *BMJ*. *BMJ* 1994;308:4-5. (1 January.)

Public funding should also be declared

EDITOR,—Though I understand the principle behind identifying the interests of the authors of published articles and that pharmaceutical and industrial interests are being distinguished, there are other factors.¹ The influence of governments must not be forgotten, one particular example of this being the current controversy regarding the use of the more expensive antidepressants rather than the older, cheaper tricyclic agents. Song *et al* did not make clear that their research had been funded by the Department of Health.² Though it is usually obvious when a pharmaceutical company is involved in a study, the involvement of some government funded bodies is much more difficult to ascertain.

It would also have been helpful on that occasion