especially in rural areas. Doctors are pressed to require compensation from their patients, who can hardly afford to pay it. The salary of a doctor amounts to \$US2 a month, so who can blame them? Obviously the state cannot sustain the 66 000 doctors, and many of them have become unemployed over the past two years. With one doctor per 170 inhabitants, Georgia has about the highest ratio of doctors to patients in the world.

The Ministry of Health has no hard currency to buy drugs: the shelves are filled with drugs from humanitarian relief agencies. Defining the quantities of drugs needed is difficult since previous consumption of drugs is not a good criterion for defining present needs in a system that is hospital based and where overprescription is the rule. Not all health staff are familiar with the generic names of pharmaceutical products used in the West, so distribution of drugs is not just a matter of supply. The recipients need information and training if they are to make correct decisions on drug treatment.

The national immunisation programme has collapsed. The supply of vaccine from Russia has stopped, and already small outbreaks of diphtheria and measles have been reported. To start up an immunisation programme again, antigens from Russia will have to be replaced with others and a new programme will have to be introduced, for which thorough training will be necessary.

There is no solution to Georgia's health problems without economic recovery and health reform.

PIM DE GRAAF

Médecins Sans Frontières,
PO Box 10014,
1001 EA Amsterdam,
Netherlands

Arbitration in medicolegal disputes

EDITOR,—Malcolm Forsythe's editorial on arbitration and the internal market is a useful contribution to the growing debate on the resolution of disputes in medicine.¹ Though arbitration undoubtedly offers a viable alternative to litigation in medicolegal disputes, however, it is not the only alternative but is part of a range of techniques encompassed by the term "alternative dispute resolution." The other main technique is mediation, which differs from arbitration in several important respects: the proceedings are without prejudice and confidential, and any resolution is arrived at by the disputants themselves and not imposed by the neutral mediator. If agreement is reached it is not legally binding on the parties, but this is not necessarily a weakness of the process. The advantages of mediation in medicolegal disputes are speed, low cost, a reduction of stress on both parties, and the likelihood of resolution satisfactory to both sides.

The use of alternative dispute resolution in commercial disputes is now well established in Britain, largely owing to the efforts of the Centre for Dispute Resolution. This centre was set up in 1990 with the support of the Confederation of British Industry. The first one day conference on alternative dispute resolution in medicine was held at the Royal Society of Medicine last December under the joint auspices of the Institute of Arbitrators and the BMA and was entitled "Disputes between patient and doctor. Is arbitration a solution?" In his keynote address the master of the rolls, Sir Thomas Bingham, suggested that "there must be a role for the skilled mediator at this early stage." The conference supported this view and explored the use of mediation and arbitration in resolving disputes between doctors and patients and, incidentally, partnership disputes between doctors.

With the review of complaints procedures in the

NHS due for publication soon and the universal dissatisfaction with the primacy of litigation in medical disputes, the time is now ripe for the medical and legal professions to put aside vested interests and to combine to produce a workable alternative to litigation.

JOHN QUARRIE

Orchard Medical Centre,
Maidstone,
Kent ME17 4PL

1 Forsythe M. Arbitration and the internal market. *BMJ* 1994; 308:151-2. (15 January.)

Screening for breast cancer

EDITOR,—On the basis of their interpretation of the results of the NHS breast screening programme for 1991-2¹ D J Watmough and K Kumar suggest that women aged over 50 should be screened for breast cancer by clinical examination by general practitioners rather than by mammography.² Their deductions from the data presented are, however, erroneous. They state that 1465 (22%) of the detected cancers were palpable and conclude that 78% could have been found by clinical examination. The original paper does not specify the number of palpable cancers. It says that there were 1468 invasive cancers ≤ 10 mm in size, which does not equate to the number of palpable tumours. Only 39% of the breast cancers that we detect by screening are palpable, even when the woman is examined by an experienced breast surgeon aware of the site of the mammographic abnormality. The Forrest report concluded that clinical examination is not an effective screening method when used alone. We know of no more recent data to contradict this conclusion.

The authors question whether compression of breast cancers during mammography might disseminate malignant cells. There is no evidence that this occurs. If it did the significant reduction in mortality shown in the randomised controlled trials of screening would not have occurred.³ The authors are also incorrect to suggest that mammography is unnecessary for palpable breast cancer; the treatment options may alter as a result of mammographic findings.

It is now established that mammographic screening performed to high standards can significantly reduce the mortality from breast cancer in women aged over (but not under) 50.⁴ The performance figures published by the NHS breast screening programme that we are achieving those standards.¹ This high quality of the breast screening programme has stimulated improved care for women with symptomatic breast disease, particularly the development of multidisciplinary specialist breast clinics, as suggested in Lesley Elliot's personal view.⁵ We run such a clinic, seeing women within 48 hours of referral by a general practitioner. Everything necessary for diagnosis is performed in one visit, and results are provided immediately if they are normal or within two further working days if a possible abnormality is detected.

Screening by mammography is the only intervention that has significantly reduced population mortality from breast cancer. We support continued research into all aspects of breast cancer, but the suggestion that this should be funded by the cessation of the national screening programme is misguided and ill informed.

L J YEOMAN
A R M WILSON
A J EVANS

Breast Screening Training Centre,
City Hospital,
Nottingham NG5 1PB

1 Chamberlain J, Moss SM, Kirkpatrick AE, Michell M, Lyons J. National Health Service breast screening programme results for 1991-2. *BMJ* 1993;307:353-6.

- 2 Watmough DJ, Kumar K. Screening for breast cancer. *BMJ* 1994;308:202. (15 January.)
- 3 Roebuck E. X-ray mammography and breast compression. *Lancet* 1992;340:366.
- 4 Royal College of Radiologists. *Guidelines for those involved in planning mammography in asymptomatic women*. London: RCR, 1994.
- 5 Elliot L. Breast cancer: a fighting chance. *BMJ* 1994;308:210. (15 January.)

General practitioners' role in commissioning

EDITOR,—Jonathan P Graffy and Juliet Williams are right to conclude that "health authorities must be prepared to share important decisions and offer representation at every level if a commissioning partnership is to be based on trust."¹ In Walsall this representation in the commissioning of services occurs at three levels.

Firstly, service review meetings with local providers take place regularly, at which one or often two general practitioners are part of the purchasing review team. Secondly, at the health authority level a local general practitioner is a non-executive member, able to report the view of his or her colleagues direct to the board. Thirdly, and, we think, innovatively, we have funded six local general practitioners to work on a sessional basis as "general practitioners in public health medicine." These general practitioners, who represent different localities in the borough, liaise with their colleagues to ascertain their views on health needs locally and to receive their impressions on the quality of care at local provider units. In addition, though these general practitioners may not be entirely representative of all local practitioners, they act as a sounding board for the health authority to test innovations or changes in practice and provision of services locally.

Information fed back by these general practitioners at a monthly meeting with the authority's executive officers is minuted and concerns are identified. The authority's proposed action in relation to these problems is then tabled at the next meeting for further discussion. To date, most of the feedback has concerned the acute services (which concurs with Graffy and Williams's comments), although several community unit and public health issues have been addressed. It is hoped that these latter two areas will become an increasing focus of the group's work, thus helping to re-establish links between public health and general practice, which the authors see as "the two branches of the profession . . . [that] . . . may have lost touch."

General practitioners are in a unique position to speak on behalf of their patients' health and health care needs. Like City and Hackney Health Authority, we in Walsall have recognised this and involve general practitioners in the commissioning process at every level.

PETER JAMES
SAM RAMAIAH

Walsall Health Authority,
Walsall WS1 1TE

1 Graffy JP, Williams J. Purchasing for all: an alternative to fundholding. *BMJ* 1994;308:391-4. (5 February.)

Invasive meningococcal infection after splenectomy

EDITOR,—Various articles have emphasised the lifelong risk of overwhelming sepsis after splenectomy,¹ and the importance of immunising asplenic patients with polyvalent pneumococcal vaccine.^{2,3} Mary McMullin and George Johnston point out that, while *Streptococcus pneumoniae* is the most common infecting organism, *Haemophilus influenzae*, *Neisseria meningitidis*, and *Escherichia*

coli are also important potential pathogens after splenectomy.¹ We report two recent deaths from fulminant meningococcal septicaemia in people in Western Australia who had undergone splenectomy.

The first patient was a 36 year old woman who had a splenectomy at the age of 30 for spontaneous rupture of the spleen during the third trimester of pregnancy. She had mild upper respiratory symptoms for four days and then collapsed suddenly; she died four hours later. Blood cultures subsequently grew *N meningitidis* serogroup W135.

The second patient was a 16 year old girl, who was not related to the previous patient. She had had a splenectomy during childhood for hereditary spherocytosis. She initially presented to her local hospital with fever and symptoms suggestive of tonsillitis with bacteraemia. Her condition deteriorated rapidly, and she was evacuated to Perth for intensive care but died the next day. *N meningitidis* was cultured from her blood and cerebrospinal fluid and yielded a positive result on testing with serogroup XYZW135 polyvalent antiserum.

The adult woman had received neither pneumococcal nor meningococcal vaccines, while the 16 year old girl had received one dose of pneumococcal vaccine since her splenectomy. Tetravalent ACYW meningococcal vaccine is now available in many countries. It is used in Britain for immunoprophylaxis of close contacts of an index patient with invasive disease caused by serogroup A, C, Y, or W135 of *N meningitidis*,⁴ and a similar policy is being considered by an advisory panel to Australia's National Health and Medical Research Council.

The two cases we report here emphasise the importance of offering tetravalent meningococcal vaccine (in addition to polyvalent pneumococcal vaccine) to patients who have had a splenectomy. Information on when to offer revaccination is scarce, but the duration of protection against disease provided by meningococcal vaccine⁵ suggests that an interval of five years would be appropriate, and this would coincide with revaccination with pneumococcal vaccine.⁴ Immunisation with one of the *H influenzae* type b conjugate vaccines should also be considered.

ROBERT J CONDON

Western Australian Research Institute for Child Health,
Subiaco,
Western Australia 6008

THOMAS V RILEY

Health Department of Western Australia,
East Perth,
Western Australia 6004

HEATH KELLY

Health Department of Western Australia,
Albany,
Western Australia 6330

- 1 McMullin M, Johnston G. Long term management of patients after splenectomy. *BMJ* 1993;307:1372-3. (27 November.)
- 2 Deodhar HA, Marshall RJ, Barnes JN. Increased risk of sepsis after splenectomy. *BMJ* 1993;307:1408-9. (27 November.)
- 3 Daly HM. Pneumococcal vaccine and splenectomy. *BMJ* 1993;307:1561. (11 December.)
- 4 Department of Health, Welsh Office, Scottish Office Home and Health Department. Department of Health and Social Security (Northern Ireland). *Immunisation against infectious disease*. London: HMSO, 1992.
- 5 Frasch CE. Vaccines for prevention of meningococcal disease. *Clin Microbiol Rev* 1989;2(suppl):S134-8.

Training in paediatric life support

EDITOR,—K F H Teale discusses paediatric resuscitation training¹; we wish to provide some detail on the two courses currently available. The paediatric advanced life support course is the European version of an international course originating in the United States in the late 1980s. Advanced paediatric life support has developed separately in Britain. Both are recognised by the

Resuscitation Council (UK) and, between them, are intended to offer comprehensive training in the immediate and early management of acutely ill children.

Children rarely suffer primary cardiac arrests, and therefore training has to include far more than cardiopulmonary resuscitation. The paediatric advanced life support course emphasises the importance of early recognition of children at risk and the theory and practice of the measures required immediately to reverse any deterioration and prevent cardiac arrest occurring. It should, we believe, be available to all doctors who care for children. The advanced paediatric life support course offers, in addition, details of the specific management of some of the conditions encountered and is more appropriate to those intending to make a career in emergency paediatrics.

Both courses started in Britain in March 1992 and both are expanding rapidly, with four centres running regular advanced paediatric life support courses and eight running regular paediatric advanced life support courses. Currently, however, demand for places on the courses exceeds supply. The main factor limiting expansion is the number of instructors. Because the courses encompass a broad range of acute paediatrics a high proportion of the instructors need to be experienced in treating, and continually exposed to, seriously ill children. They must also have taken and passed a provider and instructor's course. A high priority for directors of courses is therefore the training of those interested in instructing in the future. Nevertheless, the quality of instructors must be maintained, and pressure to increase numbers should not be allowed to jeopardise this.

Dealing with a child whose condition is deteriorating acutely can be frightening. Usually, however, simple measures are sufficient to reverse the process for long enough for expert help to arrive; these measures are within the capabilities of all doctors, given the right training. We endorse Teale's comment that it is vital that resources are available to allow this training to be carried out.

C GRANGE
R BINGHAM

Hospital for Sick Children,
Great Ormond Street,
London WC1N 3JH

- 1 Teale KFH. Training inadequate in paediatric resuscitation. *BMJ* 1994;308:275. (22 January.)

Health care in the US and Canada

EDITOR,—Minerva¹ refers to a paper in the *New England Journal of Medicine* that compares expenditure in the United States and Canada.² The authors note the greater expenditure on health care and poorer outcomes in the United States. Per capita expenditure is 40% higher in the United States despite the higher numbers of doctors, beds, and admissions (and longer lengths of stay) in Canada. There was also greater variation of expenditure within the United States.

Examination of admissions by diagnostic related groups suggests that American patients require slightly more complex care. Although systems used to grade complexity of care are similar, increased reimbursement for more complex cases in the United States may influence this. Expenditure per admission is 39% higher in the United States (22% higher after adjustment for the mix of diagnostic related groups) despite similar staff costs. Levels of clinical services are also comparable, although in the United States these are concentrated in a shorter stay.

Three explanations for the extra costs are suggested: administrative costs, other non-clinical services, and less intensive use of equipment and

beds. The authors refer to another study showing administrative costs to be 20% in California and 9% in Canada, which would account for half the difference in expenditure. Other non-clinical services are unlikely to explain the difference as staff costs are similar and length of stay is shorter in the United States. The authors suggest that the answer lies in the pattern of provision of services.

In Canada, specialised procedures are undertaken in a small number of large hospitals, while in the United States many community hospitals offer tertiary care services: examples include three times the number of centres providing cardiac surgery and 10 times the number of extracorporeal lithotripters. Equipment may therefore be used less efficiently and "occasional" operators may be less proficient in the United States. A shorter length of stay may increase the need for rapid access to equipment to facilitate early discharge.

Minerva's comment that administrative costs account for the difference is therefore misleading and ignores some important implications for British health care. The recently introduced market was intended to increase competition and responsiveness to customers. To continue in business provider units must supply what purchasers want. There is pressure to reduce waiting times and lengths of stay, with an expectation that shorter stays will be cheaper. The American evidence suggests that the opposite may be true as the need for "instant" access to resources increases. Also, action to reduce waiting times may lead to inefficient use of resources. Furthermore, if specialised treatment is provided locally, equipment may be used inefficiently and staff may be less expert: the protracted learning curve and fewer procedures performed annually may make maintaining skill more difficult. British purchasers of health care would be wise to consider carefully the lessons from the United States and Canada.

ALISON WATSON
JANE HALPIN

Department of Public Health Medicine,
Sunderland Health Commission,
Sunderland SR3 4AF

- 1 Minerva. *BMJ* 1994;308:214. (15 January.)
- 2 Redelmeier DA, Fuchs VR. Hospital expenditures in the United States and Canada. *N Engl J Med* 1993;328:772-8.

General practice fundholding

Fundholders not to blame for fast tracking

EDITOR,—More and more people who think that they know about fundholding seem to be trying to blame fundholders for creating a two tier system. They also complain that fundholders have too much money and are starving the district health authorities and commissions of funds to cater for the patients of non-fundholders. A recent article by Hart is a prime example of people talking without checking the facts.¹

In our area some practices are underfunded by up to 15% for inpatient and day case procedures. Fast tracking for them is potentially disastrous, but it is still happening. Practices in some parts of the region are having to institute contractual changes to combat this problem and prevent uncontrolled overspend.

This occurs because the providers are tagging the fundholders' patients and then processing them quickly, not because the practices are requesting a different level of treatment. Fast tracking by providers to preserve and even improve cash flow may well be the cause of the two tier system.

Angela Coulter's comments about patients in practices in Oxford show her lack of knowledge of the system.¹ Practices receive budgets based on historical levels of activity at NHS providers, except where the district health authority has a contract with a private agency for a particular