Clinical implications

- Systolic and diastolic blood pressure follow a circadian rhythm in normotensive people and those with hypertension, values being lower at night than during the day
- An inverse circadian rhythm in blood pressure has been reported in some forms of secondary hypertension
- This study found a blunted circadian blood pressure curve in all renal forms of hypertension and in all endocrine forms except primary hyperparathyroidism, but the circadian rhythm was normal in patients with unoperated coarctation of the aorta
- The fall in nocturnal blood pressure was about 15 mm Hg in systolic and diastolic readings (by 11% and 17% respectively) in normotensive subjects and those with primary hypertension
- The change in nocturnal blood pressure could aid the differential diagnosis of secondary hypertension

renoparenchymal and renovascular hypertension. In some hypertensive subjects with endocrine disorders nocturnal blood pressure even increased (phaeochromocytoma), whereas others had a significantly smaller fall in blood pressure (hyperthyroidism, primary hyperaldosteronism, and Cushing's syndrome). In patients with hypertension and primary hyperparathyroidism we surprisingly found a normal fall in nocturnal blood pressure. We also found a normal fall in three patients with unoperated coarctation of the aorta. This may indicate a purely mechanical mechanism for this form of hypertension. In endocrine

forms of secondary hypertension, however, as in primary aldosteronism, the blunted blood pressure curve could be due to increased hormonal activity over 24 hours.⁹

A lack of a fall in nocturnal blood pressure is not specific for secondary hypertension. If blood pressure does not sufficiently decrease during the night or even increases, however, further investigation may be warranted to diagnose secondary hypertension. A normal decrease in nocturnal blood pressure does not exclude secondary hypertension. The absence of an adequate fall in nocturnal blood pressure has important consequences for end organ damage and therefore for treatment.^{10 11}

We thank Bernard Doran for his linguistic advice.

- 1 Zadek I. Die Messung des Blutdrucks am Menschen mittels des Basch'schen Apparates. *Zeitschrift für Klinische Medizin* 1881;2:509-51.
- 2 Howell WHA. Contribution to the physiology of sleep, based on plethysmographic experiments. *J Exp Med* 1897;2:313.
- 3 Katsch G, Pansdorf H. Die Schlafbewegung des Blutdrucks. *Münch med Wschr* 1992;69:1715-8.
- 4 Hany S, Baumgart P, Frielingsdorf J, Vetter H, Vetter W. Circadian blood pressure variability in secondary and essential hypertension. *J Hypertens* 1987;5(suppl 5):487-9S.
- 5 Imai Y, Abe K, Sasaki S, Minami N, Nihei M, Munakata M. Altered circadian blood pressure rhythm in patients with Cushing's syndrome. *Hypertension* 1988;12:11.
- 6 Middeke M, Mika E, Schreiber MA, Beck B, Wächter B, Holzgreve H. Ambulante indirekte Blutdrucklangzeitmessung bei primärer und sekundärer Hypertonie. *Klin Wochenschr* 1989;67:713-6.
- 7 Middeke M, Klüglic M, Holzgreve H. Circadian blood pressure rhythm in primary and secondary hypertension. *Chronobiol Int* 1991;8:451-9.
- 8 Schrader J, Person C, Pferner U, Buhr-Schinner H, Schoel G, Warneke G, et al. Fehlender nächtlicher Blutdruckabfall in der 24-Stunden Blutdruckmessung: Hinweis auf eine sekundäre Hypertonie. *Klin Wochenschr* 1989;67:659-65.
- 9 Kern DC, Weinberger MH, Gomez-Sanchez C. Circadian rhythm of plasma aldosterone concentration in patients with primary aldosteronism. *J Clin Invest* 1973;52:2272-7.
- 10 O'Brien E, Sheridan J, O'Malley K. Dippers and non-dippers. *Lancet* 1988;i:397.
- 11 Schrader J, Schoel G. Benefits of noninvasive ambulatory blood pressure monitoring during antihypertensive therapy. *Journal of Ambulatory Monitoring* 1990;3:203-14.

(Accepted 2 November 1993)

Effect of advertising on awareness of symptoms of diabetes among the general public: the British Diabetic Association Study

Baldev M Singh, Jeremy J W Prescott, Roland Guy, Simon Walford, Moira Murphy, Peter H Wise

Abstract

Objective—To determine the impact of posters advertising symptoms of diabetes on public knowledge of these symptoms.

Design—Structured street interviews of members of the general public before, at the end of, and 10 weeks after a campaign advertising the main symptoms of diabetes.

Setting—Basingstoke and Wolverhampton.

Subjects—Three samples of 1000 members of the general public were interviewed. Samples were selected randomly but stratified to match the local population's age (20-75), sex, social class, and racial characteristics.

Main outcome measures—Knowledge of symptoms of diabetes; perceived seriousness of diabetes; and induction of anxiety about symptoms in the target population.

Results—Advertising significantly raised knowledge (without prompting) of symptoms: thirst, 245 before v 411 at end of campaign ($P < 0.0001$) v 341 after ($P = 0.0012$ v before); polyuria, 72 v 101 ($P = 0.0211$) v 92 ($P = 0.5169$); lethargy, 180 v 373 ($P < 0.0001$) v 298 ($P < 0.0001$); knowledge of weight loss and visual disturbance was unaffected. The number of subjects lacking knowledge of any symp-

toms was reduced from 550 to 388 ($P < 0.0001$). The perceived seriousness of diabetes was unaffected (mean 7.6 in each phase on a scale of 1 (not) to 10 (very). Before advertising, 449 (45%) claimed to have one or more symptoms of diabetes, but this number fell at the end of the campaign (403; $P = 0.0419$) and 10 weeks afterwards (278; $P < 0.0001$).

Conclusions—An advertising campaign raised public knowledge of diabetes symptoms without inducing fear of diabetes or anxiety about symptoms. Its potential for achieving earlier detection of non-insulin dependent diabetes should be evaluated.

Introduction

In patients with non-insulin dependent diabetes mellitus, microvascular disease at diagnosis is attributed to prolonged asymptomatic hyperglycaemia.¹⁻³ We have recently shown, however, that non-insulin dependent diabetes is infrequently asymptomatic. When systematically questioned, 93% of newly diagnosed patients reported classic symptoms, often previously ignored; 40% had had these symptoms for 12 months or more.⁴ This failure to recognise symptoms may reflect the general public's lack of knowledge of the symptoms of diabetes—half are unable to name any

Department of
Endocrinology, Charing
Cross Hospital, London
W6 8RF

Baldev M Singh, senior
registrar

Peter H Wise, consultant

Kilmartin-Baker Ltd,
London WC2E 9OS
Jeremy J W Prescott,
managing director

Basingstoke District
Hospital, Basingstoke
RG24 9NA
Roland Guy, consultant
physician

New Cross Hospital,
Wolverhampton
WV10 0QP
Simon Walford, consultant
physician

British Diabetic
Association, London
W1M 0BD
Moira Murphy, research
director

Correspondence to:
Dr Singh.

BMJ 1994;308:632-6

symptoms and only 4% could volunteer the important combination of thirst and polyuria.³ In view of the increased morbidity, and possibly mortality, associated with delayed diagnosis,⁶ reducing the time from onset of non-insulin dependent diabetes to formal diagnosis is likely to be important and might be achieved by increasing general awareness of symptoms of diabetes, aided by public health education programmes. Therefore the aim of the present study was to determine the ability of a campaign advertising diabetes symptoms to raise knowledge of the symptoms of diabetes.

Methods

The study was conducted simultaneously in two English towns, Basingstoke and Wolverhampton, between May and November 1992. It consisted of three consecutive 10 week phases: before advertising, the advertising, and after advertising had been removed.

During the advertising phase, posters (91 cm × 152 cm) advertising symptoms were displayed in a quarter of four-sheet sites in all shopping precincts and other pedestrian areas in each town. Advertisements also appeared in the local press twice weekly. The six classic symptoms selected (thirst, polyuria ("passing excessive urine"), weight loss, lethargy ("tiredness"), genital irritation, visual disturbance) are those most commonly given by newly diagnosed patients with non-insulin dependent diabetes.^{4,6} Two posters (fig 1) were specifically designed (by Kilmartin-Baker Ltd, London) to be educational and optimistic in tone without being alarmist or threatening. They were text based, in contrast with the highly visual posters of the British Diabetic Association depicting a mother injecting her child, for example (fig 2).

Structured street interviews were conducted in the final week of each phase of the study by a professional market research organisation (Business Planning and Research International Ltd, London). Altogether 3000 subjects were questioned—500 in each phase in each town. Subjects were randomly selected in the age range 20–75 years to meet quotas that proportionately reflected the age, sex, social class, and racial mix of the

This mother has to hurt the one she loves 730 times a year.

A baby with diabetes doesn't just depend on his mother for nourishment and love. The child is also dependent on two insulin injections a day, every day, just to stay alive. Sadly, thousands of parents in this country have to face this distressing task. Please help us to find a cure by sending a donation to:

BRITISH DIABETIC ASSOCIATION
10 Queen Anne Street, London W1M 0BD. Telephone: 071-323 1531

Advisory: helping people with diabetes and supporting diabetes research. Reg. Charity No. 215199

FIG 2—Example of British Diabetic Association fund raising poster advertising

populations in the two local authority areas. The samples were statistically matched for these characteristics between the phases. No subject was questioned twice, either within or between phases. People with diabetes and health professionals (doctors, nurses, and pharmacists) were excluded. Of the 3000 subjects, 1493 (50%) were male; 700 (23%), 1146 (38%), and 1154 (39%) were in social classes A-B, C1-C2, D-E respectively; ethnic origin was determined as white, Asian, Afro-Caribbean, or other in 2689 (89.5%), 181 (6%), 114 (4%), and 16 (0.5%) respectively; age distribution was 747 (25%) 20–29 years, 609 (20%) 30–39 years, 594 (20%) 40–49 years, 467 (16%) 50–59 years,

If you suffer from the following symptoms,

☐ thirst	☐ passing excessive urine
☐ tiredness	☐ weight loss
☐ blurred vision	☐ genital irritation

don't

You could be suffering from a form of diabetes. Ask your own doctor for a test – diagnosis and simple treatment could quickly restore your old sparkle.

BRITISH DIABETIC ASSOCIATION
10 QUEEN ANNE STREET, LONDON W1M 0BD. REG. CHARITY NO. 215199.

FIG 1—Posters advertising diabetes symptoms displayed in Basingstoke and Wolverhampton for 10 weeks

If you're tired, rundown, thirsty—it may not be a tonic you need.

You could be suffering from a form of diabetes. Ask your own doctor for a test – diagnosis and simple treatment could quickly restore your old sparkle.

BRITISH DIABETIC ASSOCIATION
10 QUEEN ANNE STREET, LONDON W1M 0BD. REG. CHARITY NO. 215199.

434 (14%) 60-69 years, and 149 (5%) 70-75 years. Overall, 765 (26%) claimed to have a family member with diabetes and 783 (26%) had a friend with diabetes; there was no significant difference in these characteristics between phases.

The question regarding knowledge of symptoms was first asked unprompted: "Can you tell me any of the symptoms of diabetes? . . . Anything else?" and all volunteered responses were recorded. Knowledge of symptoms was subsequently determined, prompted by the use of a letter coded, randomly ordered list of symptoms. Whether subjects themselves had any of the advertised symptoms was assessed in a similar way. When awareness of advertising was assessed an unprompted question, "Have you seen any advertising on the subject of diabetes?" was followed by a question prompted by pictures. Awareness of certain concurrent commercial advertisements was also determined by using picture prompts. Questions regarding the perceived seriousness of diabetes were asked with the aid of cards bearing combined numerical and text scales.

To determine the impact of the advertising either on new presentations of patients with non-insulin dependent diabetes or on the induction of anxiety about symptoms among the target populations, general practices were recruited to monitor the number of patients who attended specifically requesting tests for diabetes and the number of these who were confirmed as having diabetes. Other newly diagnosed patients who presented routinely with symptoms or who were found at routine screening were documented separately. Twenty one practices in Basingstoke and 22 in Wolverhampton volunteered to collect these data; they collected data during the first and second phases of the study only. New presentations of non-insulin dependent diabetes were also monitored in the local hospital diabetic clinics (Basingstoke District Hospital and New Cross Hospital, Wolverhampton).

Statistical analysis was by the χ^2 test for differences between proportions. Results were considered significant at $P < 0.05$.

Results

AWARENESS OF ADVERTISING

At the end of the advertising phase, the number of subjects who, without prompting, described the diabetes symptoms posters (fig 1) when asked to recall diabetes related advertising was 324, with 235 recalling the poster with text only and 142 the poster with pictorial material ($P < 0.0001$); the combined total fell to 215 ($P < 0.0001$) at the follow up 10 weeks after advertising was withdrawn. With prompting, 445 people recognised the poster with text only and 372 the other poster ($P = 0.0009$) at the end of the advertising phase. Twenty five people remembered the standard British Diabetic Association posters without prompting in the pre-advertising phase; the number remained low in the next two phases (35 ($P = 0.2381$) and 14 ($P = 0.1059$)), while with prompting 158, 240 ($P = < 0.0001$), and 214 ($P = 0.0016$) remembered these posters in each of the three phases. The prompted recall of concurrent poster advertising of a well known washing powder, chocolate bar, and proprietary analgesic—121, 84, and 72 respectively—indicated the greater success of both the diabetes related campaigns.

KNOWLEDGE OF SYMPTOMS

In the pre-advertising phase, unprompted knowledge of individual symptoms (table 1) was low. The combination of thirst and polyuria was given by only 51 of the 1000 people interviewed. Advertising led to a considerable increase in the awareness of thirst and

lethargy, which was sustained after the campaign. There were also significant but small improvements in the awareness of polyuria and genital irritation, these were not maintained, and there was no change in awareness of weight loss or visual disturbance. The rise in awareness of thirst and polyuria combined was small ($P = 0.0506$).

In the pre-advertising phase, lack of knowledge (unprompted) of any symptoms of diabetes (table 1) was significantly associated with male sex ($P < 0.0001$), younger age ($P = 0.0002$), lower social class ($P = 0.0382$), not having a friend or family member with diabetes (44% (227/515) with *v* 67% (323/485) without; $P < 0.0001$), but not with race. Table I shows unprompted knowledge of 0, 1, 2, or 3 or more symptoms and table II shows knowledge of symptoms prompted from a symptoms list.

PERCEIVED SERIOUSNESS OF DIABETES

Respondents were asked to rate the seriousness of diabetes on a scale of 1 to 10 ("not at all serious" to "very serious"). The mean score in each of the three phases was 7.6 ($P = 0.55$). The perceived seriousness of diabetes relative to other diseases was determined on a scale of 1 to 5 (1 = much less serious, 3 = about the same, 5 = much more serious). Small but significant movements occurred between categories; table III shows the mean scores in the pre-advertising, advertising, and post-advertising phases. The order of seriousness of these diseases relative to diabetes, ranked by mean scores, was constant between phases.

THE WORRIED WELL

Among the sampled population that was presumably non-diabetic, the number of claiming to have symptoms of diabetes (499) was high in the pre-advertising phase but fell significantly both during and after advertising (table IV). The prevalence of each symptom also fell, as did the numbers of subjects volunteering 1, 2, or 3 or more of these symptoms.

IMPACT ON NEW PRESENTATION OF PATIENTS WITH DIABETES

During the 10 weeks before and during advertising, 43 general practices monitored the numbers of their

TABLE I—Number of non-diabetic members of the general public knowing symptoms of diabetes (unprompted) before, during, and after a campaign advertising symptoms of diabetes

Symptom	Before advertising (n=1000)	At end of campaign (n=1000)	At follow up (n=1000)
Thirst	245	411***	341***
Polyuria	72	101*	92
Thirst and polyuria	51	72	57
Lethargy	180	373***	298***
Weight loss	79	90	82
Genital irritation	6	16*	14
Visual disturbance	147	136	118
No knowing no symptoms	550	388***	444***
No knowing 1 symptom	255	244	263
No knowing 2 symptoms	130	241***	212***
No knowing > 3 symptoms	65	127***	87

* $P < 0.05$, *** $P < 0.001$ (χ^2 test) compared with before advertising.

TABLE II—Prompted knowledge of symptoms of diabetes among non-diabetic members of the general population before, during and after a campaign advertising symptoms of diabetes

Symptom	Before advertising (n=1000)	At end of campaign (n=1000)	At follow up (n=1000)
Thirst	673	783***	732***
Polyuria	497	579***	520
Thirst and polyuria	418	534***	456
Lethargy	708	822***	791***
Weight loss	484	523***	519
Visual disturbance	649	683	655
Genital irritation	167	227***	194

** $P = < 0.01$, *** $P < 0.001$ (χ^2 test) compared with before advertising.

TABLE III—Perceived serious of diabetes relative to other diseases

	Before advertising (n=1000)	At end of campaign (n=1000)	At follow up (n=1000)
AIDS	1.6	1.5	1.4**
Cancer	1.8	1.8	1.8*
Heart disease	2.2	2.3	2.2
Chronic bronchitis	2.6	2.8***	2.7
Tuberculosis	2.9	3.0*	2.7*
Arthritis	3.0	3.2***	3.0***

*P<0.05, **P<0.01, ***P<0.001 (χ^2 test) compared with before advertising.

TABLE IV—Prevalence of symptoms of diabetes in presumed non-diabetic members of the general population before, during and after a campaign advertising symptoms of diabetes

Symptom	Before advertising (n=1000)	After the campaign (n=1000)	At follow up (n=1000)
Thirst	173	96***	59***
Polyuria	157	129	77***
Thirst and polyuria	65	50	24***
Lethargy	302	264	182***
Weight loss	40	26	17**
Visual disturbance	115	113	63***
Genital irritation	84	77	47***
No with any symptom	449	403*	278***
No with 1 symptom	204	232	164*
No with 2 symptoms	136	86***	78***
No with ≥ 3 symptoms	109	85	36***

*P<0.05, **P<0.01, ***P<0.001 (χ^2 test) compared with before advertising.

patients who specifically requested tests to exclude diabetes. Compared with the pre-advertising phase, there was a 71% increase in such presentations during the advertising phase (99 v 58). Since the proportion confirmed as having diabetes remained constant between the two phases (17%), the numbers of new patients with non-insulin dependent diabetes who presented themselves requesting tests similarly increased by 70% (17 v 10). This occurred while the number diagnosed as having diabetes after a routine presentation either in general practices (64 v 70) or in the local hospital diabetic clinics (75 v 90) fell by 9% and 17% respectively.

Discussion

Our hypothesis that health education about diabetes symptoms might lead to earlier diagnosis of non-insulin dependent diabetes is founded on our observation that patients often have symptoms for prolonged periods before they present,^{4,6} together with the general population's poor knowledge of the symptoms of diabetes.⁵ In our initial exploration of this hypothesis, we have determined the ability of advertising to influence the public's knowledge of diabetes symptoms.

In advertising terms this short duration, moderate intensity campaign achieved an exceptionally high level of penetration,⁷ compared not only with advertisements for commercial products but also with diabetes advertisements more oriented towards fund raising. The reasons for this are not clear, but the study advertisements were deliberately designed to educate by factually and clearly presenting the symptoms of diabetes, using a low key approach with a positive tone in order to avoid inducing anxiety and consequently denial.

The advertising campaign proved capable of raising the public's awareness of diabetes symptoms, although the effect varied between symptoms. Significant and sustained improvements occurred in the unprompted knowledge of thirst and lethargy, but the rise in knowledge of polyuria was of little practical significance and resulted in a disappointing lack of effect on the important combination of osmotic symptoms, thirst and polyuria. The prompted recall of the combined osmotic symptoms did, however, show

significant improvement. Neither unprompted nor prompted knowledge of weight loss or visual disturbance was affected by advertising. Future diabetes symptoms advertising research will focus on lethargy and the osmotic symptoms and will be designed to have greater impact on polyuria.

Advertising to promote symptoms raised the spectre of inducing anxiety about symptoms in the general population, but we found no evidence of this. There was no alteration in the perceived seriousness of diabetes, and a significant decrease occurred in those claiming to suffer from these symptoms, perhaps indicating the converse—a positive health benefit in reduced anxiety about symptoms as a result of health education.

Prompting non-specific symptoms opened the potential for a significantly increased workload in primary care but, in the event, this was small. Nevertheless, the number of patients specifically requesting tests for diabetes was increased by 70% during the 10 week advertising period and among these there remained a high diagnosis rate for diabetes. The proposal that health education about diabetes symptoms may lead to earlier diagnosis of non-insulin dependent diabetes is supported. Although the impact of advertising on numbers of cases of new diabetes was not large, the short advertising phase and the possibility of improving the design of the poster make this a likely underestimate of the potential value of such a campaign.

Cost effectiveness is difficult to determine. The total advertising costs in this study were £21 000. An improved, high profile, six month campaign throughout the United Kingdom using posters and advertising in national newspapers would cost about £350 000. If our results are extrapolated to all general practices in the two areas surveyed, 22 additional patients would have presented during the advertising phase (a cost of £1000 per case), but considering we reached 1% of the target United Kingdom population aged 20-75, the potential nationwide cost per case over six months (2.6×10 weeks; $£350\,000 / (22 \times 2.6 \times 100)$) would be £61, easily competitive with more conventional screening methods.⁸

In conclusion, a poster based public health education advertising campaign proved capable of improving knowledge of symptoms of diabetes among the general population without creating anxiety about symptoms or fear of diabetes. This approach will now be refined and its impact on achieving earlier diagnosis of non-insulin dependent diabetes further evaluated.

We gratefully acknowledge the help of general practitioners and practice nurses in Basingstoke and Wolverhampton who collected data for this study.

Clinical implications

- Patients with non-insulin dependent diabetes present late, despite prolonged symptoms before diagnosis, with a high prevalence of established micromacular and macrovascular disease
- This may be because the knowledge base of symptoms of diabetes in the general population is poor
- Poster based advertising of diabetes symptoms can be effective in educating the public without inducing fear of diabetes or anxiety about symptoms
- Refining this approach may be a cost effective way of encouraging non-insulin dependent diabetic patients to present earlier

- 1 Soler NG, Fitzgerald MG, Malins JM. Retinopathy at diagnosis of diabetes with special reference to patients under 40 years of age. *BMJ* 1969;iii:567-9.
- 2 Mincin I. Micro and macroangiopathies and other chronic degenerative complications in newly diagnosed diabetes mellitus. *Med Internat* 1970;304:1331-4.
- 3 Harris MI, Klein R, Wellborn T, Knudman MW. Onset of NIDDM occurs at least 4-7 years before diagnosis. *Diabetes Care* 1992;15:815-8.
- 4 Singh BM, Jackson DMA, Davies J, Wills R, Wise PH. Delayed diagnosis in non-insulin dependent diabetes mellitus. *BMJ* 1992;304:1154-5.
- 5 Jackson DMA, Davies J, Wills R, Pegg A, Fitzgerald F, Singh BM, et al. Public awareness of symptoms of diabetes mellitus. *Diabetic Med* 1991;8:971-2.

- 6 Hillson RM, Hockaday TDR, Newton DJ, Pim B. Delayed diagnosis of non-insulin dependent diabetes is associated with greater metabolic and clinical abnormality. *Diabetic Med* 1985;2:383-6.
- 7 Institute of Practitioners in Advertising. Advertising effectiveness awards, 1990. In: Feldurek P, ed. *Advertising works*. Henley on Thames: NTC Publications, 1991.
- 8 Forrest RD, Jackson CA, Yudkins JS. The glycohaemoglobin assay as a screening tests for diabetes mellitus: the Islington diabetes survey. *Diabetic Med* 1987;4:254-9.

(Accepted 30 December 1993)

Prevalence of HIV infection in patients attending an inner city accident and emergency department

M C Poznansky, J Torkington, G Turner, M J K Banks, J V Parry, J A Connell, R Touquet, J Weber

Department of Accident and Emergency Medicine, St Mary's Hospital, London W2 1NY

M C Poznansky, *honorary research fellow*

J Torkington, *senior house officer*

G Turner, *senior house officer*

M J K Banks, *senior house officer*

R Touquet, *consultant in charge*

Central Public Health Laboratory, Virus Reference Division, London NW9 5HT

J V Parry, *head of retrovirus unit*

J A Connell, *clinical scientist B*

Department of Genitourinary Medicine and Communicable Diseases, St Mary's Hospital Medical School, London W2 1NY
J Weber, *professor*

Correspondence to: Professor Weber.

BMJ 1994;308:636

The prevalence of HIV-1 infection among patients attending accident and emergency departments in hospitals in the United States is consistently higher than that among other groups of patients served by these hospitals.¹ A study at the Johns Hopkins University Hospital showed that 1 in 17 (6%) patients attending its accident and emergency departments were positive for HIV-1 antibodies, compared with 1 in 166 (0.6%) military recruits from the area served by the hospital.² Until the present study, no such comparison could be made in Britain as few data were available on the prevalence of HIV-1 antibodies in patients attending accident and emergency departments in Britain. This study sought to determine the prevalence of HIV infection in adult patients attending the accident and emergency department of St Mary's Hospital, London, which is in an inner city area and is attended by 56 000 new patients a year.

Patients, methods, and results

Anonymised urine samples were obtained from 875 consecutive patients seen in the accident and emergency department; blood samples were taken from 47 patients who could not produce a urine sample. No repeat samples were taken from patients who attended again later. The collection and numbering of samples conformed to the requirements of Parkside Research Ethics Committee. Each patient's sex, age, postal district of residence, general practitioner, and triage group were noted, as were the presence or absence of open wounds and whether the patient was a foreign visitor. These details were also obtained for patients who did not agree to participate in the study and did not give samples.

All but four of the 875 urine samples had adequate concentrations of IgG (>0.1 mg/l) for testing for antibodies to HIV-1 and HIV-2 with IgG antibody capture enzyme linked immunosorbent assay (ELISA) (Murex Diagnostics, Dartford). Urine samples that were reactive in this assay were titrated in an IgG antibody capture particle assay and tested by western blotting.^{3,4} Blood samples were screened for antibodies to HIV-1 and HIV-2, and if they were reactive the result was confirmed with conventional ELISAs. Of the 918 urine and blood samples tested, 12 urine samples were positive for HIV-1 antibodies, giving a prevalence of 1 in 77 (1.3%).

The patients who were positive for HIV-1 antibody differed significantly from those who were negative in that they were more likely to be male, foreign visitors, and aged 26-45 (table). All of the eight patients who

reported themselves as infected with HIV-1 when they presented at the accident and emergency department were among the 12 patients whose urine was positive for HIV-1 antibody. The patients who provided blood or urine samples were representative of the overall intake of patients in the accident and emergency department with regard to age, sex, triage group, and proportion of foreign visitors; the 69 patients who did not give samples did not differ significantly in these respects from the patients who did participate in the study. None of the 918 patients was positive for HIV-2 antibodies.

Details of 918 adults attending accident and emergency at St Mary's Hospital, according to whether they were negative or positive for HIV-1 antibody

	Negative (n=906)	Positive (n=12)	P value*
No who provided urine specimen (IgG > 0.1 mg/l)	859	12	
No who provided blood specimen	47		
No (%) of men	451 (50)	9 (75)	<0.001
No (%) of aged:			
16-25	163 (18)	1 (8)	NS
26-45	335 (37)	11 (92)	<0.001
46-95	408 (45)		<0.001
No (%) whose general practitioner was known	453 (50)	8 (67)	<0.05
No (%) who were foreign visitors	254 (28)	9 (75)	<0.001
No (%) with open wound	154 (17)	2 (17)	NS

* χ^2 test, df=1.

Comment

This study shows a relatively high prevalence of HIV-1 infection (2.4%) in patients aged 16-45 attending the accident and emergency department of St Mary's Hospital, London. This prevalence was significantly higher than that (1 in 167; 0.6%) found in a study of HIV-1 infection in hospital patients aged 16-49 in the four Thames regions (P<0.001, df=1).⁵ Our findings should alert authorities in inner cities to the prevalence of HIV infection in patients attending accident and emergency departments, even in a hospital with a large HIV clinic. There is a continuing risk of HIV-1 infection to health care workers, particularly inexperienced senior house officers practising emergency medicine in an environment that is often frenetic.

We thank Drs Philip Mortimer and Jane Fothergill for their help with the study and their critical review of the manuscript, David Parker and Ursula Beckford at Murex Diagnostics for donating the IgG antibody capture ELISA kits, and Mrs J Gandhi for technical help.

- 1 Kelen GD, Fritz S, Qaqish B, Brookmeyer R, Baker JL, Kline RL, et al. Unrecognized human immunodeficiency virus infection in emergency department patients. *N Engl J Med* 1988;318:1645-50.
- 2 Kelen GD, Fritz S, Qaqish B, Floccare D, DiGiovanna T, Baker JL, et al. Substantial increase in human immunodeficiency virus (HIV-1) infection in critically ill emergency patients: 1986 and 1987 compared. *Ann Emerg Med* 1989;18:378-82.
- 3 Connell JA, Parry JV, Mortimer PP, Duncan J. Novel assay for the detection of immunoglobulin G anti-human immunodeficiency virus in untreated saliva and urine. *J Med Virol* (in press).
- 4 Connell JA, Parry JV, Mortimer PP, Duncan RJS, McLean KA, Johnson AM, et al. Preliminary report: accurate assays for anti-HIV urine. *Lancet* 1990;335:1366-9.
- 5 Public Health Laboratory Service. Unlinked anonymous monitoring of HIV prevalence in England and Wales: 1990-92. *GDR Review* 1993;3 (1):1-11.

(Accepted 15 October 1993)