present trainees should attack this problem by insisting that vocational training is strengthened with at least two years' training in general practice—preferably by vocational training being extended to four years or, alternatively, by the present split of two years' training in hospital and one year as a trainee in general practice being reversed²—and by insisting that there is national summative assessment to the highest standards. I believe that the logical tool for that is the membership examination of the Royal College of General Practitioners. It may not be perfect, but it is a rigorous, tested assessment tool; none other exists.

Present trainees should be able to say, "I believe passionately in general practice; I have been properly trained for general practice through a new lengthened and strengthened vocational training scheme; and I have the qualifications to prove it." Only when general practice sets its own standards high will others see it in the same light.

T A CARNEY
General practitioner

Burn Brae Medical Group,
Hencotes, Hexham NE46 2ED

- 1 Beecham L. GPs want a brake on end point assessment of trainees. *BMJ* 1994;308:661-2. (5 March.)
- 2 Carney T. A national standard for entry into general practice. *BMJ* 1992;305:1449-50.

Community care: the first year

EDITOR,—Trish Groves reports on mental health care in Bassetlaw one year into the Community Care Act, referring to people with enduring mental health problems and the importance of rehabilitation.¹ There is no mention of finding community placement for the most disabled group of long stay chronic psychotic patients.

Latimer House is a nursing home in West Birmingham, set up under the joint auspices of the local health authority and the National Schizophrenia Fellowship. Twenty four residents were transferred from the long stay wards of the local hospital. At one year there was evidence of continued slow deterioration in both mental state and social functioning, similar to that found in the community resettlement of equally disabled patients elsewhere.² There was also a steady deterioration in physical health, requiring drug treatment and nursing care. Both these outcomes seem to be common in such patients, whether in hospital or community setting.³

We wish to draw attention to the practical difficulties that the community placement of this group presents. Their complex problems require interagency working—in the case of Latimer House this includes voluntary agencies, the purchaser, the general practitioner, and the hospital team. Before patients move the issues of responsibility must be established. Regular medical input, for both physical and mental problems, is vital. In Bassetlaw there were few permanent medical staff; for a group as disabled as ours continuity of care, including daytime activity, is vital. Such care is not cheap, requiring high staffing levels, large drug bills, and regular replacement of fixtures and fittings (at Kneesworth House this occurs every three months (J Taylor, lecture at Royal Society of Medicine)).

The move to community care develops the interface between primary and secondary care. Such nursing homes could provide invaluable experience for general practitioner trainees, who as principals in practice may find such a nursing home in their area.

A K STANLEY
Senior registrar

Reaside Clinic,
Birmingham B45 9BE

J F MACMILLAN
Senior lecturer

University of Birmingham,
All Saints Hospital,
Birmingham B18 5SD

- 1 Groves T. Community care in Bassetlaw. *BMJ* 1994;308:708-11. (12 March.)
- 2 O'Driscoll C, Wills W, Leff J, Margolius O. The TAPS project 10. The long-stay populations of Friern and Claybury Hospitals. The baseline survey. *Br J Psychiatry* 1993;162(suppl 19):30-5.
- 3 Cunningham Owens DG, Johnstone EC. The disabilities of chronic schizophrenia—their nature and the factors contributing to their development. *Br J Psychiatry* 1980;136:384-95.

BMA's guidelines for private fees

EDITOR,—I suggest that the Monopolies and Mergers Commission has itself made a monopolistic and unilateral declaration about the BMA's guidelines on private fees.¹ The word "guidelines" is important because guidelines are not mandatory but only advisory.

I recently needed the services of an architect and was shown a brochure from the Royal Institute of British Architects, which suggested that fees should be a fixed percentage of the total building costs. When I asked the Monopolies and Mergers Commission and the Office of Fair Trading why this did not represent a complex monopoly I was told that after an investigation a few years ago the architects had made this advice advisory.

If anything quotes fixed prices it is the British United Provident Association's manual; twice as many specialists adhere to that as to the BMA's booklet. Make no bones about it—we have been stitched up.

P C MAY
Consultant orthopaedic surgeon

Princess Royal Hospital,
Telford TF6 6TF

- 1 Beecham L. British government bans BMA's private fees guide. *BMJ* 1994;308:492-3. (19 February.)

Election of members of GMC

EDITOR,—It was interesting to see comments by several doctors in the medical press about the scant publicity given to the forthcoming elections of members of the General Medical Council.¹ The single advertisement about the elections in the *BMJ* was buried halfway through a scientific paper.² Many doctors, not unnaturally, interpreted this as indicating a desire by the council to keep a closed shop for a cosy coterie of members—this at a most critical time for the profession, when recent enforced changes have proved to be so divisive and destructive of morale. These changes have succeeded in politicising a new generation of doctors (as the only alternative to acute depression or early retirement), who now wish their voices to be heard in a proper forum.

This perceived lack of openness by the General Medical Council has succeeded in increasing the pervading cynicism of many doctors towards their political leaders and representatives. It has been suggested that the council's period for nomination for this year should have been extended,¹ and I agree. I also hope that before future elections the council will ensure that all doctors are circularised with the appropriate information in good time. Perhaps, too, elections should be held every three years and tenure should be limited to a maximum of 10 years to limit complacency and encourage fresh thinking.

I M PEEK
General practitioner

London NW3 3JX

- 1 Cooper C. GPs "frozen out" of GMC election. *GP* 1994; Mar 18:3.
- 2 *BMJ* 1994;308: facing p 239 (clinical research and general practice editions).

Health policies in British medical schools

EDITOR,—Under the influence of the World Health Organisation,¹ the development of health promotion policy and strategy is increasing.² This approach requires widespread support, particularly from doctors. Newcastle medical and dental schools are developing a corporate health policy to improve the health of staff and students; exercise leadership in health promotion; contribute to national, regional, and local health strategies; and educate students and staff. A document that was disseminated to students and staff for consultation last year proposed policy on issues including alcohol, diet, exercise, stress, sexual behaviour, drugs, environmental protection, and occupational health and safety (Newcastle University already has a no smoking policy).³ The consultation has been positive, and students and staff want to help develop the policy.

We also wrote to the deans of the 29 medical schools in the United Kingdom about health policies. Twenty medical schools responded, and 15 informed us about policy (some written, some not) on the following issues: smoking (eight medical schools), alcohol (eight), occupational health (two), hepatitis B (two), drugs (two), safety in the workplace (two), and travel during student electives (two). Several initiatives to develop policy were under way, two in response to the "health at work in the NHS" initiative.⁴ Most medical schools were reacting to policy initiatives in their university or the NHS or to legislation (for example, the Control of Substances Hazardous to Health regulations or Health and Safety at Work Act). The target group for policy was usually students rather than staff (for example, travel during student electives). Policies were unlinked and concerned single issues.

Ought medical schools to lead in developing and implementing health policies? In *Health of the Nation* "health settings" were identified as a focus for health promotion; surely the medical schools should be on a list which already includes schools and hospitals. The close relation between most medical schools and the NHS should facilitate the development of policies compatible with those in the NHS.² Our initiative has links with the Newcastle healthy city project, which is working within a health for all context. Health policy in a medical school also provides opportunities for enhancing the teaching of health promotion and public health, as recommended by the General Medical Council.⁵

The replies we received from the medical schools indicate that debate on this subject is welcome. We thank the medical schools that supplied information and would like to hear about similar initiatives from other medical schools around the world.

RAJ BHOPAL
Professor of epidemiology and public health

MARTIN WHITE
Senior lecturer in public health medicine

A L CROMBIE
Dean of medicine

Department of Epidemiology and Public Health,
School of Health Care Sciences,
Medical School,
University of Newcastle,
Newcastle upon Tyne NE2 4HH

- 1 World Health Organisation. *Global strategy for health for all by the year 2000*. Geneva: WHO, 1980.
- 2 Health Education Authority. *Health at work in the NHS*. London: HEA, 1992.
- 3 Medical School Health Policy Working Group. *A health policy for the medical and dental schools: a consultation document*. Newcastle: University of Newcastle upon Tyne, 1993.
- 4 Secretary of State for Health. *The health of the nation. A strategy for health in England*. London: HMSO, 1992. (Cm 1986.)
- 5 General Medical Council Education Committee. *Recommendations on undergraduate medical education*. London: GMC, 1993.