Secondly, all hospitals should establish clear and coordinated managerial responsibility for the transfusion process. Thirdly, transfusion laboratories should be required to have a procedure for recording all transfusion errors and the corrective action taken and to report regularly to the proposed national scheme. Fourthly, statistics on the number of blood units transfused and the number of patients who receive transfusions should be collected and reported nationally to provide denominators for any reporting scheme. Finally, pilot projects should be set up to identify and report cost effective ways of improving the safety of the clinical transfusion process.

We thank the organisers of the national external quality assurance scheme in haematology, which distributed the questionnaire and the final report to all hospital transfusion laboratories in Britain.

- 1 Roberts B, ed. *Standard haematology practice*. Oxford: Blackwell Scientific, 1991.
- 2 Szazama K. Reports of 355 transfusion-associated deaths: 1976 through 1985. *Transfusion* 1990;30:583-90.
- 3 Linden JV, Paul B, Dressler KP. A report of 104 transfusion errors in New York State. *Transfusion* 1992;32:601-6.
- 4 Murphy WG, McClelland DBL. Deceptively low morbidity from failure to practise safe blood transfusion: an analysis of serious blood transfusion errors. *Vox Sang* 1989;57:59-62.

(Accepted 17 February 1994)

***Pneumocystis carinii* in bronchoalveolar lavage fluid and bronchial washings**

Simon P Matusiewicz, Ronald J Fergusson, Andrew P Greening, Graham K Crompton, Sheila M Burns

Respiratory Unit, Western General Hospital, Edinburgh EH4 2XU
Simon P Matusiewicz, registrar
Ronald J Fergusson, senior registrar
Andrew P Greening, consultant physician
Graham K Crompton, consultant physician

Regional Virus Laboratory, City Hospital, Edinburgh EH10 5SB
Sheila M Burns, consultant virologist

Correspondence to:
Dr Matusiewicz.

BMJ 1994;308:1206-7

Antibodies to *Pneumocystis carinii* develop in early childhood,¹ although no associated illness has been identified. The classic concept is that *P carinii* pneumonia in immunocompromised patients represents a reactivation of dormant childhood infection. Studies with DNA amplification, however, have failed to detect pneumocystis in sputum² or specimens of lung tissue taken at necropsy³ from immunocompetent patients. Reinfection rather than reactivation may therefore be more relevant. To see whether *P carinii* exists as a commensal organism of the lower respiratory tract we examined bronchoalveolar lavage fluid and bronchial washings in apparently immunocompetent patients undergoing routine diagnostic (or research) bronchoscopy by using an immunofluorescent antibody test.

Subjects, methods, and results

We studied 220 patients with various clinical diagnoses (table). Patients were considered to have no

Details of 220 patients examined for presence of Pneumocystis carinii in bronchial washing or bronchoalveolar lavage specimens

	Bronchial washings	Bronchoalveolar lavage specimens
No of patients	104	116
Mean (range) age (years)	65 (22-83)	59 (24-84)
Diagnoses:		
No active disease	18	22
Asthma or chronic obstructive lung disease	6	12
Bacterial infection	10	8
Interstitial lung disease	3	13
Lung cancer*	67	61

*Cell types: squamous, 37; adenocarcinoma, 10; large cell, 16; small cell, 27; bronchiolo-alveolar, 1; carcinoid adenoma, 2; lymphoma, 1; metastatic, 12; histology unknown, 22.

active disease if investigations, including bronchoscopy, failed to show any lung disease. Patients with known HIV infection or suspected pneumocystis pneumonia or those taking immunosuppressive drugs were excluded. In all, 116 patients had bronchoalveolar lavage, specimens being obtained in aliquots of 10 ml from 240 ml lavages (8 × 30 ml) of the lingula or middle lobe; 104 patients had bronchial washing, saline being introduced into and aspirated from the lung with disease diagnosed radiologically or seen to be abnormal at bronchoscopy. All specimens were examined for

pneumocystis by a virologist (SB) without access to clinical data.

Cytospin preparations of samples were fixed (50% methanol-acetone) and stained with murine monoclonal antibodies labelled with fluorescein that react with cysts and trophozoites (Genetic Systems Corporation, SYVA). Samples were considered positive if cysts or trophozoites were seen in three or more high power fields.

Of 220 specimens examined, only five (three lavage specimens and two washings) were positive for *P carinii* by the fluorescent antibody test. The five positive samples all had cysts and trophozoites in more than three high power fields. All were from patients with small cell lung cancer. No cysts or trophozoites were seen in any fields of samples defined as negative. No patients were thought to have clinical or radiological evidence of *P carinii* pneumonia at the time of bronchoscopy. Two positive specimens were available for staining by an alternative less sensitive method (toluidine blue O), and both were positive. Four of the five patients were clinically well. Two patients had evidence of macroscopic extrathoracic disease at presentation. Two patients had additional causes for impaired immunity—namely, previous cytotoxic treatment and ectopic secretion of adrenocorticotrophic hormone.

Comment

We were unable to detect *P carinii* by fluorescent antibody testing of bronchoalveolar lavage fluid or bronchial washings in 215 of the 220 patients studied and therefore consider that it may not be a commensal organism of the lower respiratory tract. Although the antibody test requires interpretation by a skilled pathologist, it is more sensitive and specific than silver staining. Theoretically, DNA amplification techniques are able to detect one or two cells but have failed to detect organisms in sputum² or postmortem samples of lung³ in normal subjects. Not all studies, however, have shown any increase in diagnostic sensitivity of DNA amplification over fluorescent antibody testing in bronchoalveolar lavage samples.⁴ Our findings support the concept that reinfection rather than reactivation leads to disease in susceptible patients. No environmental source has been identified, although infection rates show a seasonal variation. Infection was first identified in debilitated young and elderly subjects. These groups, in addition to patients infected with HIV, may act as a reservoir for pneumocystis.

Subclinical *P carinii* infection was found in five out of 27 patients with histologically proved small cell lung cancer. This may reflect a local defect in immune function associated with small cell lung cancer, which may have prognostic implications. Previous colonisation with pneumocystis may be clinically relevant for patients who subsequently receive cancer chemotherapy as pneumocystis pneumonia is a recognised

and perhaps underreported terminal event. The importance of subclinical colonisation of the lower respiratory tract needs to be studied.

We thank Dr Margaret McIntyre and staff of the Department of Pathology, Western General Hospital, Edinburgh.

1 Pifer LL, Hughes WT, Stagno S, Woods D. *Pneumocystis carinii* infection: evidence for high prevalence in normal and immunosuppressed children. *Pediatrics* 1978;61:35-41.

- 2 Leigh TR, Wakefield AE, Peters SE, Hopkin JM, Jacoub MH, Cattell WR, *et al.* The detection of subclinical *Pneumocystis carinii* infection using the polymerase chain reaction in asymptomatic renal and heart/lung recipients receiving no *Pneumocystis carinii* prophylaxis. *Thorax* 1991;46:743.
- 3 Peters SE, Wakefield AE, Sinclair K, Millard PR, Hopkin JM. A search for *Pneumocystis carinii* in post mortem lungs by DNA amplification. *J Pathol* 1992;166:195-8.
- 4 Lipschik GY, Gill VJ, Lundgren JD, Andrawis VA, Nelson NA, Neilson JO, *et al.* Improved diagnosis of *Pneumocystis carinii* infection by polymerase chain reaction on induced sputum and blood. *Lancet* 1992;340:203-6.

(Accepted 17 December 1993)

Predicting the adult height of short children

Berit L Heitmann, Thorkild I A Sørensen, Niels Keiding, Niels E Skakkebaek

Institute of Preventive Medicine, Copenhagen Health Services, Copenhagen Municipal Hospital, DK-1399 Copenhagen K, Denmark
Berit L Heitmann, *senior resident*
Thorkild I A Sørensen, *MRC professor*

Statistical Research Unit, University of Copenhagen, DK-2200 Copenhagen N, Denmark
Niels Keiding, *professor*

Department of Growth and Reproduction, Rigshospitalet, University Hospital, DK-2100 Copenhagen, Denmark
Niels E Skakkebaek, *professor*

Correspondence and requests for reprints to: Dr Heitmann.

BMJ 1994;308:1207

Some recent studies have suggested that treatment with growth hormones can stimulate growth in very short children who do not have growth hormone insufficiency,¹ but others suggest that growth hormone treatment may modulate normal puberty and shorten the pubertal growth spurt.² It is not known whether growth hormone treatment in children with short stature will make them taller than if they had not been treated. Most of the ongoing research uses changes in standard deviation scores of height before and after treatment as the measure of effect.³ Evidence supporting the common assumption that children who are short before puberty will also be short in adulthood is lacking. The present study therefore sought to describe the adult height of subjects who were short in childhood.

Methods and results

The data represent a subset of 93 800 Danish conscripts born 1930-1956 who attended the schools in the central municipalities of Copenhagen.⁴

Height (without shoes) was measured in all subjects except for those who were unquestionably unfit for service and did not appear before the board.⁴ Short stature was, by itself, not an acceptable reason for exemption from examination. In a random sample of 529 conscripts exempted, one was a dwarf owing to chondrodystrophia. One per cent of the total

population was randomly selected for the study. This sample comprised 938 draftees, of whom 912 had had their height measured. The median age at this examination was 19 (range 18-27) years.

In all schools of the central municipalities of Copenhagen, annual systematic health examinations, including measurements of height, were carried out. Every year at ages 7 through 13, about 80 000 of the 93 800 subjects had been examined.

Height was measured to the nearest centimetre and given as integers. Boys with heights below the height closest to the third centile are denoted boys with short stature.

Mean height of boys aged 7 increased from 124 (SD 6) cm to 176 (SD 7) cm in adulthood. For each age from 7-13 years the eventual adult height of boys with short stature ($n=18-26$) was 1.68-1.91 SD below the mean, and 30-43% of these boys remained below the third centile in adulthood. None of the boys with short stature had adult heights above average, and 90% had adult heights of more than 1.0 SD below the mean.

The figure shows the cumulative distribution of adult height for boys with short stature at 7 and 13 years, together with adult height in the total sample.

Comment

Data on childhood and adult heights were collected over a period of 26 years, during which the height of Danish men increased appreciably. This secular trend affected all height classes equally.⁵ As we compared heights of the same individuals as boys and adults, this trend will not affect predictions of adult shortness from childhood shortness. All men may not have achieved final height when measured at the draft board. In 313 subjects measured 4-34 years later height increased 0.5 (1.4) cm; height of subjects who were short in childhood increased 1.1 (1.4) cm.

The finding that short boys will, if untreated, remain short through adulthood, is important for evaluation of treatment with growth hormone in short children without growth hormone deficiency. The data may serve as references for final height in healthy boys small for age.

We thank Stig Sonne-Holm, Karen Margrethe Fensbo, and Anna-Lise Petersen for their contribution to the collection and sampling of the data and Katharina Main for helpful comments.

- 1 Wit JM, Rietveld DHF, Drop SLS, Oostdijk W, Gons M, Otten BJ, *et al.* A controlled trial of methionyl growth hormone therapy in prepubertal children with short stature, subnormal growth rate and normal growth hormone response to secretagogues. *Acta Paediatr Scand* 1989;78:426-35.
- 2 Stanhope R, Albanese A, Hindmarsh P, Brook CGD. The effects of growth hormone therapy on spontaneous sexual development. *Horm Res* 1992;38 (suppl):9-13.
- 3 Lesage C, Walker J, Landier F, Chatelain P, Chaussain JL, Bouganères PF. Near normalization of adolescent height with growth hormone therapy in very short children without growth hormone deficiency. *J Pediatr* 1991;119: 29-34.
- 4 Sørensen TIA, Sonne-Holm S. Risk in childhood of development of severe adult obesity: retrospective, population-based case-cohort study. *Am J Epidemiol* 1988;127:104-13.
- 5 Christensen U, Sonne-Holm S, Sørensen TIA. Constant median body mass index of Danish young men, 1943-1977. *Hum Biol* 1981;53:403-10.

(Accepted 17 December 1993)

Cumulative distribution curves of adult height in boys who had short stature (<3rd centile) as children and for all adults in sample