Asymptomatic hepatitis C infection

Be cautious with liver biopsy

EDITOR.—In her review of hepatitis C Carol A Seymour arrives at a confusing conclusion about the value of liver biopsy in infected asymptomatic blood donors.¹ Liver biopsy "seems essential" once the condition has been diagnosed because patients respond to antiviral treatment. Seymour later admits that therapeutic trials are required in this group of patients but then advocates a nationwide policy to identify, follow up, and treat asymptomatic people infected with hepatitis C virus. The two important issues she fails to address are sampling bias in the estimation of the frequency of liver disease in asymptomatic donors and the lack of evidence that treatment affects the long term outlook in hepatitis C.

It is wrong to make statements about the prevalence of disease in selected populations. The careful documentation of the North London Blood Transfusion Centre shows how selected the group who undergo liver biopsy can become.² In contrast, the investigators in Trent region gave no indication of how many patients did not have a biopsy or what the indications for biopsy were.³ Only studies of unselected or randomly sampled populations will give reliable estimates of the prevalence of histological disease in asymptomatic subjects that are generally applicable. Sampling bias has probably led to an overestimation of the frequency of liver disease associated with hepatitis C in Trent.

The consensus reached both in a recent issue of *Drug and Therapeutics Bulletin*⁴ and at a conference on hepatitis C in Edinburgh late last year was that antiviral treatment can normalise hepatic enzymes and improve the appearances on liver biopsy. There is no evidence, however, that it affects long term prognosis.

With so much uncertainty about the prevalence of liver disease related to hepatitis C and the long term benefits of treatment it is irresponsible to advocate liver biopsy in all asymptomatic infected patients.

ROLAND VALORI
Consultant physician

JOHN CHRISTIE
Registrar in general medicine

Gloucestershire Royal Hospital,
Gloucester GL1 3NN

- 1 Seymour CA. Asymptomatic infection with hepatitis C virus. *BMJ* 1994;308:670-1. (12 March.)
- 2 Ryan KE, MacLennan S, Barbara JAJ, Hewitt PE. Follow up of blood donors positive for antibodies to hepatitis C virus. *BMJ* 1994;308:696-7. (12 March.)
- 3 Irving WL, Neal KR, Underwood JCE, Simmonds PN, James V. Chronic hepatitis in British blood donors infected with hepatitis C virus. *BMJ* 1994;308:695-6. (12 March.)
- 4 Managing patients with hepatitis C. *Drug Ther Bull* 1993;31:61-2.

More questions, few answers

EDITOR.—Unfortunately, the editorial¹ and papers on asymptomatic infection with hepatitis C virus²⁻⁴ created more questions than answers for me. I work in an inner city practice that has many intravenous drug users on its list. Over the past few years a proportion of these patients have requested HIV tests. The virology laboratory has been telling us these patients' hepatitis B and C status even

Priority will be given to letters that are less than 400 words long and are typed with double spacing. All authors should sign the letter. Please enclose a stamped addressed envelope for acknowledgment.

though we have generally not asked to know this. I now realise that we are in the middle of an epidemic. About 80% of the intravenous drug users on our list who have been tested are positive for hepatitis C virus. So far, none of these patients has shown any abnormal results of liver function tests.

Many of the early patients were not counselled before having the test. When I received the positive results nobody seemed to know what I should do with the information. As knowledge of hepatitis C virus infection has grown among intravenous drug users more patients have come forward requesting to have their hepatitis C status checked. Before the test is performed I have tried to explain the current thinking on monitoring and available treatment. Few of these patients have chosen to go for further investigations in hospital, and none have chosen to have liver biopsies or to receive interferon.

Although it may be reasonable to suggest that a liver biopsy is essential, what are we hoping to achieve by this and what do we have to offer these patients after the biopsy?

CHRIS FORD
General practitioner

London NW6 6RY

- 1 Seymour C. Asymptomatic infection with hepatitis C virus. *BMJ* 1994;308:670-1. (12 March.)
- 2 Irving W, Neal K, Underwood J, Simmonds P, James V, on behalf of Trent Regional Hepatitis C Virus Study Group. Chronic hepatitis in United Kingdom blood donors infected with hepatitis C virus. *BMJ* 1994;308:695-6. (12 March.)
- 3 Ryan K, MacLennan S, Barbara J, Hewitt P. Follow up of blood donors positive for antibodies to hepatitis C virus. *BMJ* 1994;308:696-7. (12 March.)
- 4 Bruno S, Rossi S, Petroni M, Villa E, Zuin M, Podda M. Normal aminotransferase concentrations in patients with antibodies to hepatitis C virus. *BMJ* 1994;308:697. (12 March.)

Hepatitis C can be sexually transmitted

EDITOR.—It was disappointing that Carol Seymour and Kate Ryan and colleagues failed to mention the sexual transmission of hepatitis C virus.^{1,2}

For some time there has been epidemiological evidence that hepatitis C can be sexually transmitted.³ Other studies have not shown a risk,⁴ indicating that the risk is low. This overall view is supported by other large scale studies suggesting that there is a cumulative risk with time.⁵ Epidemiological evidence of sexual transmission is supported by case reports of horizontal transmission in couples with no other known transmission risk, in whom nucleotide sequencing has shown concordance of virus type.⁶ Osmond *et al* concluded in their paper that although they found the likely risk of sexual transmission in an individual case to be low, because of the large pool of asymptomatic carriers within the population sexual transmission could account for thousands of new infections in the United States each year.⁷

This has two important implications. Firstly, Seymour supports the view that wider screening should occur. Tedder *et al*⁸ found that 2.2% of homosexual men attending a sexually transmitted disease clinic in central London had positive results for hepatitis C antibodies on second generation antibody tests. As this figure is considerably higher than that in the blood donor population, if a policy of wider screening is to be introduced then those attending such clinics represent a group at least deserving further study.

Secondly, with respect to counselling of patients positive for hepatitis C, sufficient evidence exists for it to be negligent not to counsel patients regarding possible sexual transmission. Sexual partners of patients with hepatitis C virus should be screened, and if they are discordant they should be advised to have protected intercourse as there is a small risk of sexual transmission. If long term partners are not willing to do this, they should be aware that there is a cumulative risk with time, so that they can make an informed decision. This should form part of the nationwide policy to prevent the spread of hepatitis C in the population that Seymour has called for.

SIMON CURTIS
Senior registrar

Harrison Department of Genitourinary Medicine,
Radcliffe Infirmary,
Oxford OX2 6HE

- 1 Seymour CA. Asymptomatic infection with hepatitis C virus. *BMJ* 1994;308:670-1. (12 March.)
- 2 Ryan KE, MacLennan S, Barbara JAJ, Hewitt PE. Follow up of anti-HCV positive blood donors. *BMJ* 1994;308:696-7. (12 March.)
- 3 Tedder RS, Gilson RJ, Briggs M, Loveday C, Cameron CH, Garson JA. Hepatitis C virus. Evidence for sexual transmission. *BMJ* 1991;302:1299-302.
- 4 Bresters D, Mauser-Bunschoten EP, Reesink HW, Roosendaal G, van der Poel CL. Sexual transmission of hepatitis C. *Lancet* 1993;342:210-1.
- 5 Osmond D, Padian N, Hayes W, Sheppard E, Glass S. Risk factors for hepatitis C positivity in heterosexual couples. *JAMA* 1993;269:361-5.
- 6 Rice PS, Smith DB, Simmonds P, Holmes E. Heterosexual transmission of hepatitis C virus. *Lancet* 1993;342:1052-3.

Failure in community care

Inpatient care is part of the integrated approach

EDITOR.—Jeremy Coid's editorial¹ contains sentiments about supervision registers and the government's response to proposals for community supervision orders, which will find broad agreement within the profession. It also contains misrepresentations of the research findings and aims of community psychiatry.

Despite Coid's protestations about "professional dogma," not one of the research studies he quotes suggests that community services can exist without admission beds or that "mental hospitals can close and that seriously mentally ill people can be conveniently and easily managed in the community." All of these studies report inpatient spells as part of an integrated approach to their patients' needs. What they do show is that admission is less often needed and discharge more prompt when community services are adequately developed.

Coid suggests that community services research has sinned by either excluding too many patients²

or by including too many.³ Early studies excluded patients judged at severe risk when these approaches were new and untested. With increasing confidence these exclusion criteria have been dropped.⁴ Our study indicated major benefits for moderately to severely ill patients, rather than those with milder, transient disorders. Coid suggests that community developments rely on unrealistic staffing levels obtained through "generous development grants," often at the cost of hospital services. This comment sits uncomfortably directly after work is quoted that clearly states that no extra resources were made available.⁵ Indeed, community care research has generally been characterised by a remarkable detail and frankness about resource implications.

Research into community psychiatry has been more extensive than that on any other shift in psychiatric service delivery.⁶ How many carefully controlled studies were there of the establishment of day hospitals or the move to district general hospitals? Resistance within the profession has ensured that community psychiatry has been carefully evaluated (as well it should be).

Coid is quite wrong to ask "whether community based services can safely replace inpatient beds in deprived . . . catchment areas." Community psychiatry practice aims to ease the pressure on these much needed beds by more appropriate provision for patients who can be supported out of hospital. He might find purchasers and managers more receptive to demands for adequate inpatient resources (which we all support) by showing that despite best practice they are still needed. Caricaturing community approaches as a stand alone alternative to hospital psychiatry is as damaging to sensible mental health planning as any "simplistic criteria" used by purchasers.

The difficulty of providing comprehensive, community based services in the inner city does not prove that such approaches are wrong. Rather it demonstrates, once again, the disproportionate disadvantage suffered by severely mentally ill people in these areas and the need to sustain the pressure for them to get services as good as those in more settled areas.

TOM BURNS
Professor
ANDY KENT
Senior lecturer

Department of Mental Health Sciences,
St George's Hospital Medical School,
London SW17 0RE

- 1 Coid J. Failure in community care: psychiatry's dilemma. *BMJ* 1994;308:805-6. (26 March.)
- 2 Stein LI, Test MA. Alternative to mental hospital treatment. Conceptual model, treatment program, and clinical evaluation. *Arch Gen Psychiatry* 1980;37:392-7.
- 3 Burns T, Beadmoore A, Bhat AV, Oliver A, Mathers C. A controlled trial of home-based acute psychiatric services. I. Clinical and social outcomes. *Br J Psychiatry* 1993;163:49-54.
- 4 Muijen M, Marks I, Connolly J, Audini B. Home based care and standard hospital care for patients with severe mental illness: a randomised controlled trial. *BMJ* 1992;304:749-54.
- 5 Braun P, Kochansky G, Shapiro R, Greenberg S, Gudeman JE, Johnson S, et al. Overview: deinstitutionalisation of psychiatric patients: a critical review of outcome studies. *Am J Psychiatry* 1981;138:736-49.
- 6 Solomon P. The efficacy of case management services for severely mentally disabled clients. *Community Ment Health J* 1992;28:163-80.

Psychiatrists fear care in the community

EDITOR,—Far from making a case for failure in community care, the editorial by Jeremy Coid confirms the continuing failure of psychiatrists to come to terms with the fundamental changes that are taking place in mental health care and the increasing research evidence that points to the feasibility and success of alternatives to institutional care in the treatment of severely mentally ill people.¹ Any unbiased and dispassionate appraisal of the available literature would have confirmed this—instead, Coid's editorial is a rather predict-

able response to the current moral panic created around "madness in the community," and it reasserts psychiatry's historical preoccupation with segregating mentally ill people and the continuing plea for more resources to achieve that.

None of the studies of home treatment for severe mental disorders cited in the editorial has come out in favour of the traditional method of institutional care, while all of them attest to the safety, efficacy, and—most importantly—acceptability to both clients and relatives of home treatment compared with hospital treatment.² Without providing any substantive evidence, Coid asserts that the results of these studies should not be accepted at face value, and he implies that the researchers may not have been objective in their evaluation because they stand to gain most from community based approaches. Such a claim, like the rest of the editorial, reveals the anxiety experienced by many psychiatrists when faced with any challenge to the "traditional hegemony" of hospital based psychiatry,³ such as the advantages of domiciliary care, which is not exclusively dependent on traditional psychiatric skills or knowledge. Evaluations of acute domiciliary care have shown consistently positive results; why have they not been fully integrated into existing services?

The failure of institutional psychiatry, with its reliance on custody, compulsion, and control of patients, is what is most evident in contemporary psychiatric practice, especially in our inner city areas. Coid's plea for "more of the same," either in the form of more beds (a convenient euphemism for locking up more people) or greater control of mentally ill people in the community, ignores the available evidence that such conventional methods of psychiatric care have failed over the years to address the needs of substantial client groups—namely, black people, women, homeless people, and socially disadvantaged people. It is this crisis in relation to hospital psychiatry, especially in inner city areas, that has been most evident from the cases that have been highlighted recently; the issues here cannot be addressed without recognising that in many instances the nature of psychiatric treatment continues to be determined by notions of control and custody rather than the need for cure or care. By seeking alternatives to institutional models, we are also engaged in shifting the ideological balance away from custodial aspects of psychiatry, something that may not be acceptable to many psychiatrists, especially forensic psychiatrists.

We have been providing a home treatment service in inner city Birmingham for nearly four years, and what has emerged in the context of this development in our local area is a greater recognition of the need for alternatives to hospital care in the local community and an increasing acceptance by psychiatrists that the inpatient service can be managed more efficiently and flexibly when there is easy access to acute domiciliary care. Unless we are prepared to consider such alternatives (and rigorously evaluate them, especially against criteria other than those derived from failed institutional models), most of our inpatient units will become indistinguishable from forensic secure units, something which even the most ardent advocates of hospital care would consider unacceptable.

S P SASHIDHARAN
Senior lecturer

Academic Unit,
All Saints Hospital,
Birmingham B18 5SD

- 1 Coid J. Failure in community care: psychiatry's dilemma. *BMJ* 1994;308:805-6. (26 March.)
- 2 Reynolds I, Hoult JE. The relatives of the mentally ill: a comparative trial of community oriented and hospital oriented psychiatric care. *J Nerv Ment Dis* 1984;172:480-9.
- 3 Tantam D. Alternatives to psychiatric hospitalisation. *Br J Psychiatry* 1985;146:1-4.
- 4 Mosher LR. Alternatives to psychiatric hospitalization: why has research failed to be translated into practice? *N Engl J Med* 1983;309:1579-80.

Community environment is stressful

EDITOR,—Jeremy Coid's timely editorial¹ about community care is a welcome contribution at a time when rapid changes are taking place in mental health care provision. Many questions still remain unanswered. How do we balance the civil liberties of individuals against the rights of society to protect itself? How far should society go in protecting mentally ill people from the consequences of their own actions committed under the influence of their illness? What should take precedence: an individual's right to freedom or the right to effective treatment?

The notion of "normalisation" disregards the fact that living in the community is not inherently therapeutic; people become ill in the first place while living in the community. It also underestimates the profound loneliness and alienation of mentally ill people in the community, and society's deep rooted prejudice against them. Experience from the United States shows that many "new chronic patients," who have never been institutionalised, are receiving inadequate care and pose the most difficult management problems in the community.² Many have severe mental illness, associated substance abuse, and constitute a large proportion of homeless and mentally ill people in jail. For this group the loss of asylum function of the old institutions has meant a loss of care, support, and protection. They are constantly exposed to the stresses of living alone in a sometimes uncaring and hostile environment. Denial of mental illness, misinformation about psychotropic drugs, and scare stories about the exploitative or abusive nature of psychiatry further alienate them from mental health services. A similar picture has emerged about homeless mentally ill people in Britain.³

Despite evidence of a bedrock of mental illness that requires inpatient care regardless of increases in community provision,⁴ the number of psychiatric beds is being reduced drastically. The division of opinion among psychiatrists that Coid mentions and the muted responses of the royal college are therefore worrying. We shouldn't have to wait for an increasing number of well publicised tragedies before these issues are discussed; that would swing the pendulum back, with demands for a harsh and punitive institutional care. An uncritical commitment to community care that does not address these questions may bring home the truth behind the saying "to err is human, to really cock things up requires an ideology."

SWARAN P SINGH
Senior registrar in psychiatry

Nottingham Healthcare Unit,
Mapperley Hospital,
Nottingham NG3 6AA

- 1 Coid J. Failure in community care: psychiatry's dilemma. *BMJ* 1994;308:805-6. (26 March.)
- 2 Lamb HR. Lessons learned from deinstitutionalisation in the US. *Br J Psychiatry* 1993;162:587-92.
- 3 Scott J. Homelessness and mental illness. *Br J Psychiatry* 1993;162:314-24.
- 4 Lawrence RE, Copas JB, Cooper PW. Community care: does it reduce the need for psychiatric beds? *Br J Psychiatry* 1991;159:334-40.

Supervision register needs rethinking

EDITOR,—Jeremy Coid highlights many of the difficulties in psychiatric care that we are experiencing in Bristol.¹ The inner city area with its large homeless population (the second largest in England), has high psychiatric morbidity,² and the number of compulsory admissions under the Mental Health Act has risen in the past few years. Inpatient units are full, and we also have a shortage of forensic beds. Where community developments continue, we are concerned these may uncover further hidden morbidity.

In this climate we have been directed to implement the supervision register with no extra powers