We report an unusual complication after thrombolysis was given to a patient in whom the diagnosis was wrong.

A 34 year old man was transferred from a district hospital to our invasive care unit because of haemodynamic instability after thrombolytic treatment for acute myocardial infarction.

Acute myocardial infarction had been diagnosed on the basis of 1 mm horizontal elevation of the ST segment in leads II and III and AVF and decreased pain after sublingual nitrates.

On arrival in the intensive care unit the patient was complaining of epigastric and left upper abdominal pain that radiated to his shoulder. He was pale and sweating, with blood pressure 120/80 mm Hg and pulse rate 130/min. The lungs were clear, and, except for the tachycardia, an examination of the heart yielded normal findings. Curiously, deep abdominal palpation was associated with moderate pain in the left upper abdomen. An electrocardiogram showed a sinus tachycardia of 120 beats/min. The patient was given intravenous morphine, and his pain disappeared almost completely.

Results of laboratory tests showed that the haemoglobin concentration had fallen to 8.4 g/l. Simultaneously, the patient went into shock. Transfusions and generous fluid treatment were started. Ultrasonographic examination of the abdomen showed splenic laceration and a large haemoperitoneum.

As his coagulation status was incompatible with an immediate operation the patient was given fresh frozen plasma and fibrinogen. Two hours later splenectomy was performed under general anaesthesia. The operation was uneventful, and the patient left hospital a week later. On close questioning he remembered having fallen while skiing four days before admission.

Many signs in this patient's history were not consistent with the diagnosis of acute myocardial infarction; sinus tachycardia is unusual in this setting. Nevertheless, this case is a good example of the risks and pitfalls of overzealous thrombolytic treatment. We suggest that a history of participation in violent sports or possible accidents or falls should be sought before thrombolytic treatment is given.

N MILLIET C DIEM
K HAYOZ F SADRY
E HERRERA V PHILLIPS

Hôpital Cantonal,
1708 Fribourg,
Switzerland

1 Weston CFM, Penny WJ, Julian DG, on behalf of the British Heart Foundation Working Group. Guidelines for the early management of patients with acute myocardial infarction. *BMJ* 1994;308:787-71. (19 March.)

Train relatives to intervene

EDITOR,—In his editorial on myocardial infarction Peter Herbert mentions training in basic life support for the families of patients with heart disease and that this training apparently causes families less anxiety than had been feared.¹ I wish to emphasise the value of targeting such training and to state more positively that anxiety need not be increased (and can even be decreased) in both the patient and his or her relatives. Training in basic life support, including how to call for help, for patients with heart disease and their families makes excellent sense as cardiac death generally occurs out of hospital, with over 60% of fatal attacks occurring in the home.²

Colleagues and I studied the effects of anxiety on training in basic life support for patients with heart disease and their relatives.³ We found that the 49 trainees in the study already had some anxieties; when a coping strategy in the form of training was provided those anxieties were reduced or at least not increased.³ This applied to both the patients and their relatives. We thought it important to

involve the patient as well as the relatives so that the issues were in the open and could be discussed by all parties. (In another study only family members, not the patient, were trained and anxiety in the patients was increased.)⁴

Single session training courses in basic life support are not yet widely available in Britain, but voluntary schemes such as the Bart City Life Saver and Frenchay City Life Saver provide training for the general public that would be suitable for patients with cardiac disease and their families. The British Heart Foundation will help and support local "Heartstart" schemes. Support for the families from general practitioners and hospital doctors is vital so that any anxieties or questions can be dealt with and debriefing is available should a family member attempt resuscitation.

If more cardiac deaths are to be prevented the general public must be trained in what to do in the vital first four minutes and until paramedics arrive with a defibrillator. The suspicion that training in basic life support increases anxiety for patients with heart disease or their family is wrong.

CHRISTOPHER A J MCLAUGHLAN
Accident and emergency consultant

Frenchay Hospital,
Bristol BS16 1LE

- Herbert P. Suspected myocardial infarction and the GP. *BMJ* 1994;308:734-5. (19 March.)
- Tunstall-Pedoe H. A coronary heart register in east London [dissertation]. Ann Arbor: Universal Microfilm International: 78-70,028.
- McLaughlan CAJ, Ward A, Murphy NM, Griffin MJ, Skinner DV, Camm AJ. Resuscitation training for cardiac patients and their relatives—its effect on anxiety. *Resuscitation* 1992;24: 7-11.
- Dracup K, Guzy PM, Taylor SE, Barry J. Cardiopulmonary resuscitation (CPR) training: consequences for family members of high-risk cardiac patients. *Arch Intern Med* 1986;146: 1758-61.

Lack of knowledge is not the problem

EDITOR,—The publication of three short papers¹⁻³ and an editorial⁴ relating to the pre-hospital initiation of aspirin by general practitioners in patients with suspected acute myocardial infarction prompts me to report similar results from a recent study carried out in a relatively rural community in Scotland.⁵ A case series of 107 patients with confirmed myocardial infarction of distinct onset, admitted to a single district general hospital, was reviewed retrospectively. Unlike in urban settings, 97% of the study population had been seen by a general practitioner before admission.

On the basis of hospital case records (which invariably included a referral letter from the attending general practitioner) only 12 patients (11%) were recorded as having definitely been given aspirin by their general practitioner. This was at variance with the findings of an unpublished cross sectional postal survey of 74 local general practitioners (response rate 74%), of which 42 of the 55 responders (76%) indicated that it was their policy always to give aspirin to patients with suspected acute myocardial infarction, in the absence of contraindications. Only three (5%) practitioners indicated that they never gave aspirin in these circumstances. Reasons cited for only sometimes giving aspirin were "forgot to give it"; "patient too ill"; "not always carried in emergency bag"; "patient definitely being admitted to hospital anyway"; "patient already on aspirin"; "refusal by patient"; and "too busy with other, more pressing priorities." Some of these reasons are, of course, perfectly legitimate, but they seem unlikely to be operating in most cases.

From a simplistic point of view, therefore, the low rate of administration of aspirin by general practitioners seems to relate more to attitudes and practices than merely to lack of knowledge. Unfortunately this serves only to underline the complexity of the task facing those who are trying

to promote the widespread implementation of broadly based clinical guidelines.

DAVID BREWSTER
Senior registrar in public health medicine

Common Services Agency for the
Scottish Health Service,
Edinburgh EH5 3SQ

- Moher M, Johnson N. Use of aspirin by general practitioners in suspected acute myocardial infarction. *BMJ* 1994;308:760. (19 March.)
- Wyllie HR, Dunn FG. Prehospital opiate and aspirin administration in patients with suspected acute myocardial infarction. *BMJ* 1994;308:760-1. (19 March.)
- Moher M, Moher D, Havelock P. Survey of whether general practitioners carry aspirin in their doctor's bag. *BMJ* 1994; 308:761-2. (19 March.)
- Herbert P. Suspected myocardial infarction and the general practitioner. *BMJ* 1994;308:734-5. (19 March.)
- Brewster D. Initial management of acute myocardial infarction in a rural area. *Health Bulletin (Edinburgh)* 1993;51:240-50.

Screening for diabetes

EDITOR,—Baldev Singh and colleagues reported the results of the British Diabetic Association's study on the effects of advertising on the general public's awareness of diabetes, and they suggested that this approach should be further evaluated as a means of achieving earlier diagnosis of non-insulin dependent diabetes mellitus.¹ We have doubts about this conclusion.

One of the reasons for these conclusions was that the approach could be cost effective (£1000 per new case of diabetes). This method has a high false positive rate and a low specificity, as of those presenting to the general practitioner, only 17 (17%) were found to have diabetes. The cost does not compare well with other methods and presumably does not include the cost of performing diagnostic tests on the 82 patients without diabetes.²

In terms of achieving an earlier diagnosis of diabetes, the 10 week campaign revealed an additional 17 cases of diabetes. The total population aged 15-75 years subjected to the campaign was approximately 300 500 according to the 1991 population census.³ It has often been assumed that there are as many undiagnosed cases of diabetes as there are diagnosed, and if we assume an overall prevalence of diabetes of 1% (probably an underestimate in this age group) then we would expect that over 3000 cases of diabetes remain undetected. The ability of this campaign to identify only a predicted 22 (0.7%) of these subjects must be disappointing.

In contrast, we have shown in a large study based in general practice that a postal request system with self testing for postprandial glycosuria is an effective method of screening for diabetes.² In a target population of 13 795 subjects aged 45-70 years, 99 new cases of diabetes were identified by this method at a cost of £81 per new case (1990 prices). We have proposed ways in which this cost could be reduced.² We have since shown that this method can be effectively repeated after 30 months, suggesting that public cooperation in repeated testing for diabetes can be maintained.⁴

The current study, as a means of raising public awareness regarding diabetes, seemed to achieve its aim; it may also have been beneficial in terms of reducing anxiety about symptoms of diabetes. However, it does not seem effective in making an early diagnosis of—that is, screening for—non-insulin dependent diabetes. When other more effective and inexpensive methods based in general practice have already been shown to have promising results, stretched resources should firstly be channelled into these approaches.⁵

MELANIE DAVIES
Senior registrar

Leicester Royal Infirmary,
Leicester LE1 5WW

JOHN DAY
Consultant physician

Ipswich Hospital,
Ipswich IP4 5PD

- 1 Singh BV, Prescott JJW, Guy R, Walford SW, Murphy M, Wise PH. Effect of advertising on awareness of symptoms of diabetes among the general public: the British Diabetic Association study. *BMJ* 1994;308:632-6. (5 March.)
- 2 Davies MJ, Metcalfe J, Williams DDR, Day JL. Community screening for non-insulin-dependent diabetes mellitus (NIDDM). *Q J Med* 1993;86:677-84.
- 3 Office of Population Censuses and Surveys. *Key population and vital statistics: local and health authority areas*. London: HMSO, 1991:6-16. (Series VS 18, PP1, No 14.)
- 4 Davies MJ, Day JL. Screening for non-insulin-dependent diabetes (NIDDM): how often does it need to be repeated? *Diabetic Med* 1993;10(suppl 3):S29.
- 5 Patterson KR, Professional Advisory Committee of the British Diabetic Association. Population screening for diabetes mellitus. *Diabetic Med* 1993;10:777-81.

Hepatitis B and admission to medical school

New policy unclear and unreasonable

EDITOR,—We agree with A M L Lever's editorial on hepatitis B and medical students.¹ While some of our views on this subject have already been described,² we should like to make several additional points.

We strongly support existing practices whereby medical students are screened for hepatitis B virus and vaccinated early after admission to medical school. The medical school and each student share responsibility to complete the vaccination protocol. After this students should be investigated for seroconversion, told of their antibody titre, and advised about boosters. Students who fail to seroconvert but are shown to be uninfected should repeat the vaccination. Students who are found to be infectious carriers of hepatitis B virus (e antigen positive) should be counselled and receive interferon treatment. While they are infectious their courses and preregistration house officer year should be modified to exclude "exposure prone procedures," as described by the Department of Health's guidelines.³ All students should have access to occupational health and careers advice.

The new policy being adopted by the Council of Deans of UK Medical Schools, the Committee of Vice-Chancellors and College Principals, and the Universities and Colleges Admission Service is both unclear and unreasonable. It is unclear in not specifically addressing the issue of students who fail to seroconvert but are shown to be uninfected. We see no justification for such students being denied entry to medical school or having their admission delayed. With regular testing of their antigen status these students do not pose any risk to patients.

The policy is unreasonable in being an alarmist overreaction to a problem with a manageable solution, where risk of infection can be reduced to bizarre theoretical levels. With regard to the future careers of affected students, there are already adequate guidelines on pre-employment screening for specialties in which "exposure prone procedures" are performed. In addition, we would not be opposed to restrictions on registration for people who are infectious chronic carriers, limiting the areas of practice open to them. The necessary statutory amendments could be included during the present review of General Medical Council health procedures.

In view of our objections and those described elsewhere, we recommend that the new policy on hepatitis B and medical students should be withdrawn and attention switched to promoting current practices, which, when efficiently implemented, protect both students and patients from this infection.

RUPERT GAUNTLETT
Medical student representative

MARK BAILEY
Chairman

BMA medical students committee,
London WC1H 9JP

- 1 Lever AML. Hepatitis B and medical school admission. *BMJ* 1994;308:870-1.
- 2 Kingman S. Hepatitis B status must be known for medical school. *BMJ* 1994;308:876.
- 3 NHS Management Executive. *Protecting health care workers and patients from hepatitis B*. Lancashire: NHSME, 1993. (HSG(93)40.)

Test medical students while preclinical

EDITOR,—A M L Lever may be right in declaring that routine testing for hepatitis B e antigen before students enter medical school sets a dangerous precedent but is wrong to imply that infected students present no risk to patients.¹ Medical students are likely to help in theatre during their clinical years and during the preregistration year as house surgeons. Unless universities alter the curriculums to abolish such clinical experience or the General Medical Council changes the mandatory requirements for registration, all students who are positive for hepatitis B e antigen will at some stage be an infective risk in theatre. When students have graduated and start surgical house jobs their employing hospital will, as a matter of course, ensure that they have been immunised against hepatitis B and, if they have not responded to the vaccine, screen them for markers of infection. So why not do this at medical school before they start their clinical years?

It is misleading to suggest that this is the thin end of the wedge with regard to multiple serological testing. In any assessment of the risk of transmission of bloodborne infection from doctor to patient hepatitis B virus is by far the most important pathogen; there have been numerous examples of carriers of hepatitis B e antigen causing outbreaks through surgical procedures. HIV and hepatitis C virus do not have such an established track record. Unfortunately, even since an effective vaccine was introduced the number of outbreaks of hepatitis B associated with surgery or dentistry has not fallen—hence the need of guidelines from the Advisory Group on Hepatitis.²

Personally, I would not want anyone who might be positive for hepatitis B e antigen to have his or her hands in my abdominal cavity near sharp instruments. All students should be immunised in their preclinical years and should know their response to the vaccine. It is then simple to test the 7% or so who are non-responders for the presence of hepatitis B surface and e antigens. They should be referred for clinical management, have career counselling, and have their clinical studies and preregistration house jobs adapted to cater for their infective status.

A G ELDER
Senior registrar in occupational medicine
Glasgow Occupational Health,
Glasgow G1 1JA

- 1 Lever AML. Hepatitis B and medical student admission. *BMJ* 1994;308:870-1. (2 April.)
- 2 UK Health Departments. *Protecting health care workers and patients from hepatitis B: recommendations of the Advisory Group on Hepatitis*. London: HMSO, 1993.

Patients must come first

EDITOR,—One body of evidence and two different conclusions is nothing new in medicine. One reason is the difference between, on the one hand, the luxury of opinion without responsibility and, on the other, the necessity of making publicly accountable decisions. The Council of Deans of UK Medical Schools and Faculties repeatedly debated the potential risk to patients from students infected with hepatitis B virus and the responsible policy to adopt. We took impeccable expert advice. After two years of deliberation we arrived at the recommended policy challenged by A M L Lever,¹ a policy that vice chancellors were free to accept or reject and chose to accept.

The situation was changing, and the matter had

to be addressed: concerns included new awareness of potential risks, which were small but real enough to worry our consciences and interest lawyers; growing public concern, acknowledged by the General Medical Council, that doctors may be a health risk to their patients; the widespread development of medical curriculums that no longer separate preclinical from clinical work and bring students into contact with patients from their earliest days at medical school; and greater emphasis on competence in practical skills before qualification and the continued requirement for hands on medical and surgical responsibility as a house officer before full registration with the General Medical Council. Applicants to medical school want to become doctors, which almost invariably means that they wish to look after patients. If for reasons of their health, as it concerns either themselves or others, they are absolutely or relatively unsuited to the task (in a highly competitive field) they need to know this at the outset. Further, it is important that they are told about the many non-medical alternative pathways to major contributions to health care through, for example, laboratory science, epidemiology, economics, or management. They may also, of course, need advice about their own treatment.

Deans are familiar with the occasional but unenviable task of balancing the interests of needs be patients against the aspirations of would be doctors with problems of health or habits. The task falls morally to the dean, while legally it seems to be nobody's responsibility. The public can reasonably expect the balance of doubt to fall in its favour. If hepatitis B is only the tip of an iceberg of ensuring as far as possible the safety of students from their patients and patients from their students, so be it: that is the cost and responsibility of our calling. That patients should come first is not "totalitarian" but uncompromisingly our professional duty. In these circumstances, for the *BMJ* to put the weight of its authority behind the subtitle "No reason to exclude applicants who are e antigen positive from medical school" is unfortunate if not irresponsible.

PETER RICHARDS
Chairman

FRANK HARRIS
Executive secretary

Council of Deans of UK Medical Schools and Faculties,
PO Box 138,
Leicester LE1 9HN

- 1 Lever AML. Hepatitis B and medical student admission. *BMJ* 1994;308:870-1. (2 April.)

Safer use of traditional remedies

Don't generalise about traditional remedies

EDITOR,—David J Atherton's article on traditional remedies makes repeated generalisations about these remedies, their practitioners, and patients.¹ He states that "when people from less developed countries emigrate they continue to seek medical advice from traditional practitioners" and that "large amounts of traditional medicine are imported into Britain," but he offers no references to substantiate these claims. It would be interesting to know if there is indeed any evidence to support his statements.

Atherton also credits Indian practitioners of traditional medicine with training, understanding, and beneficial effects. It then almost seems a disservice to speculate about the hazards of their treatments, particularly when orthodox or Western medicine has a long history of progress through empiricism. Perhaps some traditional therapies have a rather better holistic approach, which compensates for their lack of "science."

Many of David Atherton's points strike us as sensible, and there does seem to be little published