flexibility, less confrontation, and more openness, as well as striking an appropriate balance between professional self regulatory and employer based mechanisms for dealing with problem doctors. Moreover, more varied retirement and other exit options must be created to enable doctors with problems to depart with dignity when the circumstances are appropriate. Without all this patients and the quality of their care will suffer as they have done in the past because of a reluctance to face up to difficult issues.

- 1 NHS and Community Care Act 1990. London: HMSO, 1990.
- 2 National Health Service. *Hospital medical and dental staff (England and Wales). Terms and conditions of service April 1986.* (Revised.) London: HMSO, 1986.
- 3 Department of Health. *Disciplinary procedures for hospital and community medical and dental staff.* London: DoH, 1990. (HC(90)9.)
- 4 Ministry of Health. *Disciplinary proceedings in cases relating to hospital medical and dental staff.* London: MoH, 1961. (HM(61)112.)

- 5 Department of Health and Social Security. *Prevention of harm to patients resulting from physical or mental disability of hospital or community medical or dental staff.* London: DHSS, 1986. (HC(82)13.)
- 6 Secretaries of State for Health, Wales, Northern Ireland, and Scotland. *Working for patients. NHS consultants: appointments, contracts and distinction awards.* Working paper 7. London: HMSO, 1989.
- 7 UK Health Departments, Joint Consultants Committee, and chairmen of Regional Health Authorities. *Hospital medical staffing—achieving a balance.* London: HMSO, 1986.
- 8 Savage W. Disciplinary procedures and secrecy. *BMJ* 1991;303:52.
- 9 Smith R. Profile of the GMC: the hordes at the gates. *BMJ* 1989;298:1502-5.
- 10 General Medical Council. *Professional conduct and discipline: fitness to practice.* London: GMC, 1993.
- 11 General Medical Council. *Annual report of the General Medical Council 1992.* London: GMC, 1993.
- 12 *Medical Act 1983.* London: HMSO, 1983.
- 13 General Medical Council. *Proposals for new performance procedures: a consultation paper.* London: GMC, 1992.
- 14 Smith R. The GMC on performance: self regulation on the line. *BMJ* 1992;304:1257-8.

(Accepted 25 January 1994)

Treating childhood asthma in Singapore: when West meets East

G J Connett, B W Lee

Though Western medicines and ideas about asthma have become popular in many Asian nations, local beliefs about treatment prevail. The multiracial society of Singapore shows a variety of beliefs about causes of asthma attacks (for example, the balance of yin and yang) and types of treatment—herbal remedies, inhaled versus eaten medicines, the influence of Ramadan. Many of the cultural practices mentioned are probably preserved among south east Asian minorities residing in the United Kingdom. Eastern treatments typically take a holistic approach to asthma and do not ignore the psychosomatic component of the disorder.

As in most Western countries, asthma is increasingly common in many Asian nations. Its prevalence during childhood (5-17 year olds) in Singapore is 12% among Chinese communities, 19% among Malays, and 15% among Indians (unpublished data). In recent years Western medicines and ideas about asthma have become popular, but despite this trend local beliefs about treatment prevail. Some of these sound implaus-

ible, but others hint at underappreciated differences in the pathogenesis of asthma between Asians and white people.

Singapore is a multiracial society, consisting mainly of Chinese (76%), Malays (15%), and Indians (6%). These main racial groups are culturally diverse, as are their beliefs in traditional medicine. The Chinese in Singapore originate from southern China and belong mainly to the Cantonese, Fukien, and Teochew dialect groups. These dialect groups share the same written language and have common traditional beliefs about illnesses and their cures. The Malays in Singapore are also a heterogeneous community, with main origins from Javan and Bawean provinces of Indonesia. Like the Chinese, they share distinct beliefs about diseases and their cures.

This article reviews some of the local beliefs about asthma reported by the parents of children attending the respiratory clinics of the National University Hospital of Singapore. Although this report is based on our experience with childhood asthma, these local beliefs apply to asthma as a disease entity and are not unique to childhood asthma.

Department of Paediatrics,
National University of
Singapore, Singapore 0511
G J Connett, overseas fellow
B W Lee, associate professor

Correspondence to:
Dr Lee.

BMJ 1994;308:1282-4

Inter-racial differences in food induced symptoms of asthma suggest unexplored processes that cause changes in bronchial reactivity and the expression of atopy

SPRAGUE/PANOS

Causes of asthma attacks

Central to understanding on how to treat asthma in our cosmopolitan community is to realise that eating is the most important national preoccupation. Beliefs about what foods should or should not be taken during illness are often held with great conviction and are considered just as important for recovery as the doctor's prescription. Although some families' concern with food becomes obsessional, it would be counterproductive to dismiss such ideas on the grounds that medical evidence is lacking or because they are difficult to explain.

Whereas many Western people with asthma identify pollens, grasses, and moulds as important precipitants of symptoms, aeroallergens are less frequently implicated in equatorial climates where exposure is often perennial. Although viral infections are often recognised as precipitants of wheezing, the most common answers to "what makes your child's asthma worse?" are cold fruits, cold drinks, ice cream, and chocolate from the fridge. It is difficult to understand the basis for this purported effect of all things cold on asthmatic

airways, but it is consistently reported by all Asian races. Although airway cooling has been implicated in the pathogenesis of asthma induced by exercise, this effect occurs in conjunction with airway drying, probably as a result of increased ventilation.¹ It is unlikely that the ingestion of "things from the fridge" would cause significant changes in airway temperature. The effect of cold foods is usually reported to start several hours after ingestion and to continue for days at a time. Exercise induced symptoms are typically of rapid onset without a late reaction to the airway stimulus.

The concept of cold ingestion extends beyond that of temperature. The Chinese believe that for the body to function optimally there must be a balance of yin and yang elements. Associations of these polar opposites include yin coolness and yang warmth. All foods are characterised by their relative balance of these two elements. Cough, wheeze, and too much phlegm are attributed to an excess of yin and so yin types of foods should be avoided by children prone to asthma symptoms. These foods include fruits such as apples, grapes, oranges, and melon. Heaty (yang) fruits, including papaya, banana, and durian, are acceptable during respiratory illnesses, although not in large quantities.

The pathophysiology of these effects is unknown. Scepticism is common, and food related asthma has been attributed to the psychological effects that might occur in a family when a food taboo is transgressed.² However, there are some scientific data in support of food related asthma. Controlled challenge tests have confirmed increased bronchial reactivity after ingestion of ice among asthmatic Asians.³ Cold fruits include citrus fruits, and there is evidence suggesting that changes in oesophageal pH mediate vagal reflexes on bronchial reactivity.^{4,5} Allergenicity to orange seeds has also been shown among Asians.⁶ It is difficult to explain the effects of other hot and cold foods as stated in local folklore. For example, chicken is a "cold" meat and should be avoided during asthma, whereas "hot" meats such as crocodile and dog are held to be of particular benefit. Asthma symptoms are commonly attributed to fried foods in Indian families.

Ideas about asthma from Western medicine have done little to dispel these notions. Radioallergosorbent and skin tests for food sensitivities suggest that Asians produce IgE antibodies to the same spectrum of food allergens as white people and not to the foods commonly identified as problematic by parents. Local diets are usually based around rice and noodles, and dairy products are rarely implicated as causing asthma. Seafoods such as crabs, prawns, lobster, and shellfish emerge as the occasional cause of immediate hypersensitivity reactions, as do peanuts, but these reactions are usually cutaneous. Taken together, these data suggest that food allergies in the Asian population are similar to those documented in white people. On the other hand, Asians' sensitivities to ice ingestion and to acidic fruits such as oranges suggest possible differences in the triggers and pathophysiology of bronchial reactivity between races.

Treating asthma

The restoration of the body's balance of yin and yang as part of Chinese medicine's approach to treating illness has important implications for the use of medication for asthma. Tonics for the lungs are the preferred form of therapy, and Chinese herbal remedies such as the root of *Ephedra sinica* remain popular. There are no published clinical trials validating the use of Chinese herbs to treat asthma, although recent studies have shown them to have benefits in treating eczema⁷ and bronchiolitis.⁸

The concept of inhaling medicines directly into the airways, without reaching the rest of the body, seems nonsensical to a culture where the traditional approach is to treat the whole person. Some of the older members of Chinese families harbour concerns that the regular use of inhaled medicines can lead to addiction. Some children might become psychologically dependent on excessive puffs of bronchodilator inhalers, but apprehensions of grandparents may be caused by the connection between the use of inhaled medicines and memories of the ruinous habit of smoking opium, a practice prevalent in this part of the world at the turn of the century.

The Chinese prefer to eat their medicines; they prefer a syrup or a potion rather than a pill. It is no surprise therefore that ketotifen remains one of the most popular treatments for asthma in this part of the world. Not only is it consumed by mouth, but its stimulation of appetite and the resultant weight gain are perceived as effective therapeutic properties. A good body weight is extremely important, and updating weight on the centile chart is considered as important as recording the latest peak flow measurement. Ketotifen is so effective in making children sit up and eat their dinners that it is one of the few drugs that parents come and demand from the outpatient clinic. The lack of validated effects on respiratory function does not detract from its perceived efficacy in promoting good health. Unfortunately, the similar properties of oral steroids have led to some practitioners prescribing them in circumstances when inhaled anti-inflammatory drugs would be a safer alternative.

The Malay culture presents a seasonal problem in treating many illnesses. During the period of Hari Raya (Ramadan) food and drink are forbidden during daylight hours. Although fasting is excused for those with illnesses such as diabetes, many people with asthma undertake their religious obligations. The exclusion of food includes the taking of medicines. This seasonal lapse of compliance has been recognised as a cause of deterioration in asthma symptoms. Daytime smoking is permitted during Hari Raya and so (it can be argued) is the taking of inhaled medicines. Hari Raya presents a good opportunity to make the switch to using inhaled asthma therapy.

Traditional remedies

In countries with such a large preoccupation with food it is inevitable that some rather obscure oral remedies have evolved. Some Chinese believe that swallowing live, newborn mice is particularly effective. Several parents have testified to this, but we have yet to meet anyone who has given this treatment to their child. The Malays often cook up their remedies in a coconut shell. The remedy preferred for asthma is a skinned frog (species unspecified) poached gently in a fresh green coconut. An Indonesian parent reported two tried and tested remedies: bat's lung fried whole in a dry pan until well blackened, and dried gecko. Geckos are lizards that enjoy a symbiotic relationship in Singaporean houses, where they feast on winged insects around light fittings. Their effects on asthma are achieved by sun drying a killed lizard, grinding it to a fine powder, and then ingesting it in small quantities over a period of a few days. The taste is quite bitter but can be disguised by mixing the powder with mashed banana.

Proprietary cough medicines are commonly prescribed for asthmatic cough. The dubious medical benefits of these preparations are supposedly enhanced by the use of the kris. This double edged dagger, with its characteristic serpentine blade, is a traditional Malay weapon endowed with far reaching mystical

powers. These include the ability to cure cough—the point of the dagger is dipped into a cough syrup before ingestion. The kris has not, to our knowledge, been used on a metered dose inhaler. Pieces of jade are commonly invested with healing properties by many Chinese families, and some Indian families “disguise” their asthmatic children with make up so that avenging gods do not recognise them and cast illness on them.

Conclusions

Many of the above practices are anecdotal and hint at superstitious quackery, but it is clear that the social consequences of asthma among children in Asian communities is very different from those in their Western counterparts. Pity the small child deprived of chilled drinks and ice cream in a tropical climate. It is easy to be scathing of non-scientific traditional practices, but something can be learnt for the benefit of all patients. Psychological factors are extremely important in determining the expression of asthma, and these have received little attention during the recent push towards a greater use of drug treatment in the Western world. Eastern treatments typically take a holistic approach to the problem of asthma and do not ignore this psychosomatic component of the disorder. Interracial differences in food induced symptoms suggest unexplored, genetically determined, pathophysiological processes that cause changes in bronchial

reactivity and the expression of atopy. Commonly used herbal remedies might yet stand the rigours of double blind, placebo controlled studies and become safe, acceptable alternatives to the anti-inflammatory drugs that are currently used by Western doctors.

Illnesses affect cultures, and cultural practices have a considerable effect on the consequences of illness. The rich diversity of culture in south east Asia cannot be matched anywhere else in the world. Discovering what Asian families believe about illness is not only fascinating but is essential in reconciling the differences between what doctors want them to do, and what they think they should do, about treating their children's asthma.

- 1 Fitch KD, Morton AR. Specificity of exercise-induced asthma. *BMJ* 1971;iv: 577-81.
- 2 De San Larazo C. Food related asthma: a difference between two ethnic groups. *Arch Dis Child* 1986;61:97.
- 3 Wilson NM, Dixon C, Silverman M. Increased bronchial responsiveness caused by ingestion of ice. *Eur J Resp Dis* 1985;66:25-30.
- 4 Wilson NM, Charette L. Gastro-oesophageal reflux and childhood asthma: the acid test. *Thorax* 1985;40:592-7.
- 5 Rauscher H, Popp W, Ritscha L. Effect of short oesophageal acidification on airway hyperreactivity. *Respiration* 1989;55:11-5.
- 6 Zhu SL, Ye ST, Yu Y. Allergenicity to orange juice and orange seeds: a clinical study. *Asian Pac J Allergy Immunol* 1989;7:5-8.
- 7 Sheehan MP, Atherton DJ. A controlled trial of traditional Chinese medicinal plants in widespread non-exudative atopic eczema. *Br J Dermatol* 1992;126: 179-84.
- 8 Kong XT, Fang HT, Jiang GQ, Zhai SZ, O'Connell DL, Brewster DR. Treatment of acute bronchitis with Chinese herbs. *Arch Dis Child* 1993;68:468-71.

(Accepted 22 February 1994)

Equity issues in the NHS

Who cares about equity in the NHS?

Margaret Whitehead

This is the first of a series of five articles about equity in the health services. Margaret Whitehead will be speaking next week at a conference on equity in the NHS organised by the Society for Social Medicine and the Public Health Alliance. The conference will be in Oxford on Thursday 19 May and the fee is £25. Further information is available on 021 643 7628.

The Old School, Ash
Magna, Whitchurch,
Shropshire SY13 4DR
Margaret Whitehead,
independent health policy
analyst

BMJ 1994;308:1284-7

The concept of equity in relation to the National Health Service in Britain encompasses not one but at least eight distinct principles. Until the 1980s the NHS had a good record of incorporating these principles into practice. Throughout the 1980s, however, there has been a pronounced change, with the gradual introduction of business values into the service, culminating in the market based reforms of the 1990s. Several recent policies seem to be taking the NHS away from the goal of an equitable system—for example, the new arrangements for community care and the incentives within contracting to select patients on financial grounds. To restore equity as a value demands priority for ethical values, monitoring of policies for their effects on equity, some national planning, and a new debate about the entitlement to services such as continuing care.

Over the past decade successive reorganisations of the NHS have been presented as technical adjustments, to improve efficiency, quality, and patient choice but without affecting the basic equitable foundations of the service. The impression is of noble intentions, but if we dig beneath the rhetoric is the principle of equity being upheld and protected, or is it being quietly abandoned?

Why the NHS was established

In the midst of continual reorganisation it is easy to forget why the NHS was set up in the first place. The political consensus for change grew out of a widespread realisation that the prewar system was inequitable,

inefficient, and near to financial collapse.^{1,2} In the 1930s, for instance, only 43% of the population were covered by the national insurance scheme, mainly men in manual and low paid occupations and only for general practitioner services.¹ That left 21 million people, predominantly the wives and children of employed men, not covered by the scheme. The sick carried much of the burden of payment and could be faced with devastating bills.

Across Britain services were spread unevenly, with deprived areas poorly served both in quantity and quality of service. This arose because doctors needed to make a living from private practice and only in more prosperous districts could local authorities raise enough local taxes to provide a range of high quality public services. In some places the same practitioner provided a highly visible two tier service:

Sometimes middle class patients go to the doctor's front door and working class patients to the surgery door. One class of patient comes by appointment, the other is expected to take his turn and may have a considerable time to wait.³

A highly fragmented and unplanned service added to the haphazard and inefficient use of resources.⁴ For example, 1000 voluntary hospitals and 2000 municipal hospitals were all administered separately and often competed with one another. In 1945 the National Hospital Survey reported that in some places increasing competition for dwindling charitable donations had led to wasteful duplication of equipment and high technology processes in desperate ploys to attract funds.^{5,6} Competing hospitals were also reluctant to transfer fee