ultimate power, and capital punishment is not a mandatory sentence for the expression of non-violent options. The death penalty is only suggested as the maximum punishment for several crimes, and it is the judge's ruling that sets the sentence; appeal is possible. If the appeal fails a parliamentary commission reviews the sentence, and only if the commission agrees with the sentence will parliament vote on it. No such sentence has been voted on in the past 10 years, before which the country was under military rule.

Turkey is a developing country desperately trying to improve its human rights record and relations with the West. At a time when the extreme right and religious fundamentalism are gaining power in most parts of the world, developed countries, especially those in Europe, should consider supporting democratic governments in their struggle for humanitarian ideals and condemning all acts of terrorism.

MEHMET RAFET GAZVANI

Registrar
Obstetrics and Gynaecology Department,
Singleton Hospital,
Swansea SA2 8QA

1 Mansour P. Doctors and execution in Turkey. *BMJ* 1994;308:656-7. (5 March.)

First aid for snake bite

EDITOR,—I was angered by the covert messages in Drs Craig Gerrard and Janice Adam's piece on *Minerva's* page.¹ I would very much like to know what they would have done to help a child bitten by a poisonous snake if they had been in the foothills of the Himalayas in the 1930s. It certainly would not have been possible to flag down a passing bullock cart and nip to the nearest accident and emergency department for a quick antivenom jab.

K TINKER

Southampton SO14 6UW

1 *Minerva*. *BMJ* 1994;308:1050. (16 April.)

Increasing the number of organ donations

EDITOR,—Once again the shortage of donors for organ transplantation has been highlighted.¹ Various suggestions have been made to try to improve the current system of using organ donor cards.² Earlier this year I surveyed 231 consecutive patients attending health promotion and disease management clinics and asked about their views on organ donation. Altogether 217 (94%) patients returned a completed questionnaire. Seventy two were male (average age 45 (range 18-72)) and 145 were female (average age 35 (16-76)).

Fifty patients normally carried an organ donor card, but only 35 had the card with them that day. One hundred and fifteen said that they were willing to be an organ donor when they die, 32 said that they did not want to be an organ donor, and 70 did not know. One hundred and forty six were willing for their decision to be recorded in their medical notes, nineteen did not want any record made in their notes, and 52 did not know. Those who were undecided about possible organ donation accounted for the majority (43) who did not want any record made in their notes.

The proportion of patients who normally carried an organ donor card was similar in this study to that reported in previous studies. The study shows the inadequacy of the current donor card system, with 30% of patients not carrying their card on the day in question. It also shows that half of patients would be willing to be an organ donor when they died, this was more than double the number who had organ donor cards. Recording this information

in the notes was acceptable to almost all those who wished to be an organ donor.

General practitioners are encouraged to record information about patients that is of dubious value. If, however, a practice was known to record its patients' views about organ donation the practice could be contracted at the time of death of a potential organ donor to find out his or her views on organ donation. This simple and cost effective system could potentially double the number of organ donations.

RICHARD VAUTREY
Trainee general practitioner

Spotland Group Practice,
Rochdale OL12 6PH

- 1 Court C. Shortage of donors hinders organ transplantation. *BMJ* 1994;308:938. (9 April.)
- 2 Riad HN, Banks RA. Recording patients' view on organ donation: when to ask and how to record the answer. *BMJ* 1990;301:155.

Hidden danger in water melons

EDITOR,—D A T Southgate's answer to a question concerning the safety of eating water melons is appropriate to Britain,¹ but in parts of India and Africa there are two additional risks. One is the

practice in some places of injecting water—often river or pond water—into melons to increase their weight and consequently their price. The other is the practice of selling sliced melon on open stalls by the roadside with no protection against flies or other contamination.

FRANK I TOVEY
Medical adviser

Interhealth,
Mildmay Mission Hospital,
London E2 7NA

1 Southgate DAT. Any questions. *BMJ* 1994;308:777. (19 March.)

Textbook on endocrinology

EDITOR,—The *BMJ's* book reviews generally provide succinct and pertinent comments that encourage—or otherwise—browsing or buying. One difficulty that authors and editors have is in targeting their readership. We were therefore surprised to read F J Woodroffe's comments about *Diagnostic Tests in Endocrinology and Diabetes*.¹ The boundaries between endocrinology and metabolism are non-existent physiologically and are arbitrary in the medical and literary worlds. Woodroffe's suggestion that hyperlipidaemia, hyponatraemia, and glycogen storage disease do

not fit into such a textbook seems strange when they are relevant to diabetes and other endocrine disorders.

We recently reviewed 25 adult patients, aged 17-35, with the hepatic glycogen storage diseases who had attended the paediatric departments of King's College Hospital and the Hospital for Sick Children, London, over the past 30 years. The table shows the specialties of their current physicians. The most remarkable findings are the number of patients not being actively followed up—despite the increasing number of long term complications including polycystic ovaries,² hepatic tumours, renal disease, hyperlipidaemia, and osteoporosis³—and the number still being managed by paediatricians.

These results emphasise that there is an expanding cohort of patients with glycogen storage diseases surviving into adulthood, and adult physicians need to be aware of this. Some of these patients are indeed managed by endocrinologists, and thus these diseases probably warrant a place, albeit a small one, in textbooks that may be read by such specialists.

PHILIP LEE
Clinical research fellow
JAMES LEONARD
Professor of metabolic disease

Institute of Child Health,
London WC1N 1EH

- 1 Woodroffe FJ. Diagnostic tests in endocrinology and diabetes [book review]. *BMJ* 1994;308:1049. (16 April.)
- 2 Lee PJ, Patel A, Hindmarsh PC, Brook CGD, Leonard JV. Polycystic ovaries and glucose tolerance in hepatic glycogen storage disease [abstract]. *Pediatr Res* 1993;33:S14.
- 3 Moses SW. Pathophysiology and dietary treatment of the glycogen storage diseases. *J Pediatr Gastroenterol Nutr* 1991;11:155-74.

Poets' corner

EDITOR,—James Owen Drife writes that he "flew to the bookshop and embraced it [*The Virago Book of Birth Poetry*] triumphantly."¹ Had he embraced it a little less and opened his hands (and his mind) a little more, he would not have suffered from the "welling inside" that was caused by the perceived absence of obstetric contributions to *The Virago Book of Birth Poetry*. He would have seen that two physicians do appear in the book: John Stone, professor at the Emory University School of Medicine, and Sally Sange, a practising physician in Merritt Island, Florida. Stone's poem, "To a 14 Year Old Girl in Labor and Delivery," is the "obstetric contribution" that Drife insists is not there.

The two poems (purportedly written by two obstetrician colleagues) that Drife submits as examples of "empathic verse missed by the *Virago* anthologist" smell of the rubber apron that he wore to Waterstone's. As editor of *The Virago Book of Birth Poetry*, I feel disappointment welling up inside me when I read "Night feed" and "Obstetric extraction." I fail to detect the "throbs" of the "heart of the poet" that Drife hears. But, then, we use different instruments.

CHARLOTTE F OTTEN
Professor of English

Calvin College,
Grand Rapids, MI 49546,
USA

1 Drife JO. Poet's corner. *BMJ* 1994;308:796. (19 March.)

Specialty of physicians managing 25 adult patients with glycogen storage diseases type I and type III. Figures are numbers of patients

Diagnosis	Specialty					
	No	Metabolic medicine	Endocrinology	Hepatology	Paediatrics	None
Type I disease	13	2	2	2	3	4
Type III disease	12	2	1		2	7
Total	25	4	3	2	5	11