Teaching laparoscopic surgery

Practise on live animals is illegal

EDITOR,—There is increasing pressure on the royal colleges to introduce some form of accreditation and assessment of practical skills in minimally invasive surgery. This is partly driven by attention in the media to cases that have had an unfavourable outcome. There has never, however, been a demand for such accreditation in general surgical skills. Theoretically, a doctor who has never performed surgery can obtain a fellowship of the Royal College of Surgeons or membership of the Royal College of Obstetricians and Gynaecologists. Yet there are no calls for assessment of the technical competence of general surgeons or gynaecologists. Perhaps it would be more appropriate if assessment of technical skill in surgical specialties generally, rather than just in innovations such as minimally invasive surgery, was addressed.

C M S Royston and colleagues discuss different modes of training in laparoscopic surgery and suggest that the porcine model is useful.¹ They do not make clear that performing laparoscopic procedures on living models is illegal in Britain. Thousands of novice laparoscopists have attended hands on courses in Britain and have carried out advanced procedures on patients, albeit under the supervision of an experienced tutor. It is ironic that the law is less protective of patients than animals when it comes to learning new surgical skills.

PAUL BYRNE

Consultant obstetrician and gynaecologist

Royal College of Surgeons in Ireland,
Department of Obstetrics and Gynaecology,
Rotunda Hospitals,
Dublin

1 Royston CMS, Lansdown MRJ, Brough WA. Teaching laparoscopic surgery: the need for guidelines. *BMJ* 1994;308:1023-5. (16 April.)

Training courses are popular and valuable

EDITOR,—The proved advantages of laparoscopic cholecystectomy have put considerable pressure on the surgical community to learn new techniques.¹ Reports of high complication rates,^{2,3} however, have suggested that the training and evaluation of laparoscopic general surgeons need urgent attention.⁴ To address these important issues a practical training course was devised, based on laparoscopic repair of a hernia, and participants were invited to evaluate it. We also invited suggestions on ways of ensuring adequate training of future laparoscopic surgeons. Repair of an inguinal hernia was selected as the ideal model for teaching advanced procedures to experienced laparoscopic surgeons both because standard laparoscopic techniques are used and because anatomy as seen through an endoscope, stapling techniques, and the handling of prostheses must be learnt.

Twenty three surgeons attended the two day training courses, which comprised comprehensive lectures, a review of the literature, video demonstrations, and practice on artificial simulators. Each surgeon assisted in one repair and then performed one under supervision. A questionnaire was sent to each participant.

Twenty two questionnaires were returned (96%

Letters should be no more than 400 words long, should contain a maximum of five references including one to the *BMJ* article to which they refer, and must be typed with double spacing. All authors need to sign the letter and provide their current appointment and address. Please enclose a stamped addressed envelope if you require an acknowledgment. First reports of original research are not published in the letters section.

response rate). Seventeen respondents had previously performed at least 50 laparoscopic cholecystectomies and could therefore be deemed experienced laparoscopic surgeons. All rated the format of the course, the quality of the speakers, and the technical and practical content as good to excellent. Nineteen respondents thought that the artificial laparoscopy simulators were adequate. Four of the five surgeons who had previously been taught on live animals thought that our course would not have been improved if pigs had been used as a training model.

Nineteen participants thought it most important that surgeons should attend a course such as ours before attempting a first case. Twenty and 16 respondents advocated similar courses for surgeons of senior registrar and registrar grade respectively. Eighteen said that they would attend another such course—for example, on colorectal laparoscopy. Suggested ideal training requirements for unsupervised practice were on the job training with attendance of a recognised course (n=22), completion of a specified number of procedures (14), and a probationary period with an audit of morbidity (12).

This survey shows, firstly, that, once the basic skills of laparoscopy have been attained and the means are available for supervised practice, animal training models are not needed; secondly, that surgeons are interested to learn advanced laparoscopic techniques; and, thirdly, that a high proportion of the surgeons questioned think that training courses are important and should be available to junior surgeons. Many courses on laparoscopy are available, varying in both content and quality. We hope that the recent establishment of the three national minimally invasive therapy training units by the Royal College of Surgeons may go some way to solving the problem of providing standardised training. Questions concerning the accreditation of laparoscopic surgeons have yet to be answered.

CHARLES C NDUKA
Research assistant

ARA DARZI
Consultant surgeon

Department of Minimally Invasive Surgery,
Central Middlesex Hospital,
London NW10 7NS

- 1 Wastell C. Laparoscopic cholecystectomy. *BMJ* 1991;302:304-5.
- 2 Kingman S. Private hospital ban laparoscopic surgery. *BMJ* 1993;306:1227.
- 3 Tompkins RK. Laparoscopic cholecystectomy: threat or opportunity? *Arch Surg* 1990;125:1245.
- 4 Cuschieri A. The laparoscopic revolution—walk carefully before we run. *J R Coll Surg Edinb* 1989;34:295.
- 5 Royston CMS, Lansdown MRJ, Brough WA. Teaching laparoscopic surgery: the need for guidelines. *BMJ* 1994;308:1023-5.

See all patients preoperatively

EDITOR,—C M S Royston and colleagues' article on teaching laparoscopic skills contains a disturbing sentence.¹ The suggestion that patients could "be operated on by a surgeon who they have never met and probably never will meet" is unwise and probably unethical. As an itinerant teacher of transthoracic endoscopic sympathectomy I have gone to other hospitals and always make a point of seeing the patients preoperatively. It is precisely because the local surgeons are unfamiliar with the procedure that careful preoperative counselling by the visitor is essential.

ALAN CAMERON
Consultant surgeon

Ipswich Hospital NHS Trust,
Ipswich IP4 5PD

1 Royston CMS, Lansdown MRJ, Brough WA. Teaching laparoscopic surgery: the need for guidelines. *BMJ* 1994;308:1023-5. (16 April.)

Vascular surgery in smokers

Doctors are well qualified to decide on treatment . . .

EDITOR,—Julie Dawson makes many valid points in her letter on whether the medical profession has the right to decide whether smokers should be entitled to vascular surgery,¹ but she overlooks two issues. She asks how much longer society is prepared to let members of the medical profession take it on themselves to decide who gets treatment. Doctors have always had to carry out this duty because society has never put forward any group better educated or qualified to do so. I find it difficult to believe that there is another body of people who are so qualified, but this should certainly not be a political issue. What is clear is that society in general and the medical profession in particular welcome any guidance on the difficult problems of what society should both be expected to pay for and have a right to expect as treatment.

The second point raised by Dawson relates to outcomes. When we, as a medical body, decide what treatment is offered to patients we inevitably make judgments on whether such treatment will benefit patients—or, in other words, whether there will be a favourable outcome. We want to offer treatments that have good outcomes since those treatments are most valuable. If a patient diminishes the probability of a good outcome of an intervention then the value of that intervention diminishes, maybe to the extent that the same resource could be better deployed elsewhere.

What we have not agreed as a society is how much blame should be incorporated as part of the allocation of priorities. I believe that ascribing blame would be a dangerous course to adopt since so many ills that affect us are in one way or another self inflicted. The blameworthiness of smoking and of falling over when drunk are relatively easy to assess, but what about the blameworthiness of a meniscus being torn during a game of football? We as a medical profession encourage people to play sports to keep fit, but any sport has its associated dangers. Are sports then to be blameworthy or exempt? If football is acceptable, what about skydiving?

These are difficult questions that need public debate by all concerned with ethics in medicine.