address the issue of selection bias in their study population. Hypertension in black people may be diagnosed at a more advanced stage than in the general population, and referrals to a specialist clinic may also differ by ethnic group. This would mean that the true degree of hypertension was greater in the black people studied.

These selection biases can be overcome by a population based study. In such a study we showed that average resting systolic blood pressure is 6 mm Hg higher in men of black African descent and 17 mm Hg higher in women of black African descent than in the general population.² Further, we showed that the nocturnal fall in blood pressure is less in black than white people, even when resting blood pressure is taken into account. The overall burden of hypertension for a similar blood pressure would therefore be greater in black than in white people.

People of black African descent have a greater mortality from stroke,³ more left ventricular hypertrophy,⁴ and more renal damage⁵ than white people. Racial differences in cardiac structure alone cannot account for this. A more plausible explanation is that ethnic differences in hypertensive load are inadequately characterised by current methods of blood pressure measurement.

N CHATURVEDI
Research fellow
M G MARMOT
Professor

Department of Epidemiology and Public Health,
University College London,
London WC1E 6EA

P M McKEIGUE
Senior lecturer

Department of Epidemiology and Population Sciences,
London School of Hygiene and Tropical Medicine,
London WC1E 7HT

- Mayet J, Shabi M, Foale RA, Poulter NR, Sever PS, Thom SAMC. Racial differences in cardiac structure and function in essential hypertension. *BMJ* 1994;308:1011-4. (16 April.)
- Chaturvedi N, McKeigue PM, Marmot MG. Resting and ambulatory blood pressure differences in Afro-Caribbeans and Europeans. *Hypertension* 1993;22:90-6.
- Office of Population Censuses and Surveys. Mortality and geography: a review in the mid-1980s. *The Registrar-General's decennial supplement for England and Wales*. London: HMSO, 1990. (Series DS No 9.)
- Hammon IW, Devereux RB, Alderman NH, Lutas EM, Spitzer MC, Crowley JS, et al. The prevalence and correlates of echocardiographic left ventricular hypertrophy among employed patients with uncomplicated hypertension. *J Am Coll Cardiol* 1986;7:639-50.
- McClellan W, Tuttle E, Issa A. Racial differences in the incidence of hypertensive end-stage renal disease (ESRD) are not entirely explained by differences in the prevalence of hypertension. *Am J Kidney Dis* 1988;12:285-90.

(3750 g) 260 g higher than previous British birth-weight standards, which are based on menstrual dates and show considerable flattening at term.²

We have since examined our accuracy of gestational dating by ultrasound measurement of the fetal biparietal diameter using Campbell and Newman's dating tables.⁴ We studied 19 consecutive singleton, normal term pregnancies resulting from assisted conception by in vitro fertilisation or intrauterine insemination booked at our hospital. All had the usual detailed ultrasound examination at 18-19 weeks by ultrasonographers who were not aware of the precise gestational age at the time of measurement. Gestational age derived from the biparietal diameter⁴ was compared with the true gestational age as determined from the date of fertilisation. The mean error (age estimated on ultrasonography minus true gestational age) was -0.57 days, with a standard deviation of 2.96 days and normal distribution.

We then looked at the possible effect of constitutional variation shown by the measurement obtained from the dating scan on the subsequent birth weight. There was no significant correlation between biparietal diameter centiles, adjusted for gestational age, and birth weight for gestation centiles ($r=0.256$; $P=0.37$). Furthermore, there was no significant difference in average birth weight for gestation centile between the two groups of babies that had the largest and smallest biparietal diameters (average of centiles, 47 v 40).

Thus the 95% confidence interval for ultrasound dating by biparietal diameter is -5.9 to 5.9 days in our unit. In contrast, as we previously reported,⁵ "certain" menstrual dates (routine ultrasound dates being used as a reference) have a heavily skewed distribution of error of -9 to 27 days (95% confidence interval). This inclination for menstrual dates to overestimate the true dates can lead to birth weights being plotted against gestational ages that are too advanced. We believe this to be the reason for an artificial flattening of the birthweight curve, emphasizing the need for dates of determined on the basis of ultrasound scans to be used to derive accurate birthweight standards.

JASON GARDOSI
Director
THEO MUL
Research fellow
MAX MONGELLI
Research fellow

Perinatal Research and Monitoring Unit,
Department of Obstetrics and Gynaecology,
Queen's Medical Centre,
Nottingham NG7 2UH

MARK WILCOX
Visiting lecturer

Department of Obstetrics and Gynaecology,
Prince of Wales Hospital,
Shatin NT, Hong Kong

Ultrasound dating and birth weight at term

EDITOR,—Tine Brink Henriksen and Allen Wilcox' comment on our finding that the birthweight curve in our study was steeper at term than that observed in other studies.² We ascribed this to our use of dates based on routine ultrasound scans instead of menstrual dates to derive our weight for gestational age standard.² They suggest that a constitutionally large fetus's gestational age is overestimated by the dating scan and that therefore bigger babies, with presumably larger birth weights at term, would selectively be plotted against later gestations, thus creating bias.

While it is true that the study they refer to showed variation in biparietal diameter between small (≤ 10 th centile) and large (≥ 90 th centile) babies,³ the difference between these extremes amounted to only about 1.5 mm at 18 weeks (figure 2 in reference 3). This difference is equivalent to less than one day's variation in gestational age⁴ and would translate at term to a difference in birth weight of under 20 g. In contrast, our mean birth weight at 40 weeks (3522 g) is about 120 g higher, at 41 weeks (3652 g) 167 g higher, and at 42 weeks

- Henriksen TB, Wilcox A. Ultrasound dating subject to bias. *BMJ* 1994;308:201. (15 January.)
- Wilcox M, Gardosi J, Mongelli M, Ray C, Johnson I. Birth weight from pregnancies dated by ultrasonography in a multicultural British population. *BMJ* 1993;307:588-91. (4 September.)
- Persson PH, Grennert L, Gennser G. Impact of fetal and maternal factors on the normal growth of the biparietal diameter. *Acta Obstet Gynecol Scand* 1978;suppl 78:21-7.
- Campbell S, Newman GB. Growth of the fetal biparietal diameter during normal pregnancy. *Journal of Obstetrics and Gynaecology of the British Commonwealth*. 1971;78:513-9.
- Gardosi J, Mongelli. Risk assessment adjusted for gestational age in maternal serum screening for Down's syndrome. *BMJ* 1993;306:1509-11.

Vitamin K for neonates

EDITOR,—Whether vitamin K predisposes children to cancer certainly needs investigation. However, this issue was not central to my article,¹ as Gerald Draper and Andrew McNinch seem to suggest.² My question is whether any serious side effects exist for this drug and whether it is possible to justify treatment of 25 000 babies at low risk of

haemorrhage to avoid one major haemorrhage without regard to the possibility that we might endanger many more than one child. Because of the difficulty of investigating small risks of unspecified type, the treatment decision will eventually rely on prejudice concerning the likelihood of risks more injurious than the benefits. The point I made was that grosser risks can, and should, be excluded by clinical trial.

It is difficult to comment on doubts about the ethics of a randomised, controlled trial when it is not clear what treatment strategy is being proposed as more ethical. The options for infants at low risk of early haemorrhage seem to be to recommend treating all; to recommend treating none; and to recommend joining a clinical trial. Probably, with our present very limited state of knowledge, any of these would be ethical provided a clear statement of our uncertainty is made to the parents. The major advantage of the clinical trial is that it helps to reduce uncertainty in a way that none of the other options can. It is this property which makes me believe that it is the most ethical policy.

Doctors should be cautious about endorsing a policy of treatment for all. The argument that we know for sure that one very rare disease is eliminated by treatment is weak. If a child suffers a haemorrhage when untreated, both doctors and parents must suffer the pain of knowing that, with hindsight, this could have been avoided. However, avoidance of such personal pain is not a good reason for following a treatment strategy which may cause more suffering overall but allows us to remain ignorant of our causal role.

If non-treatment is recommended then Draper and McNinch's suggestion of making carers aware of the importance of warning bleeds should be followed, and it probably makes sense to suggest to breast feeding mothers who experience major difficulties during the early weeks that they consult a health visitor concerning supplementary feeding without delay.

JIM SLATTERY
Medical statistician

Department of Clinical Neurosciences,
University of Edinburgh,
Edinburgh EH4 2XU

- Slattery J. Why we need a clinical trial for vitamin K. *BMJ* 1994;308:908-10. (2 April.)
- Draper G, McNinch A. Vitamin K for neonates: the controversy. *BMJ* 1994;308:867-8. (2 April.)

Thalidomide may be a mutagen

EDITOR,—The birth of two malformed children in England raises the question whether thalidomide, α -phthalimidoglutarimide, is a human mutagen as well as a potent teratogen. The fathers of both children are thalidomide victims.

CASE 1

In July last year a girl was born in Peterborough with no thumbs and only two digits on both hands. She has severe malformations of both legs, and the left leg is much shorter than the right. Both feet taper to one toe, neither of which has nails. Her father was born in 1960 with malformations of both hands and both legs. He was treated at the Hospital for Sick Children, Great Ormond Street, and his legs were amputated below the knee because of the severe malformations of both feet. He was assessed by the Thalidomide Victims' Compensation Panel and was awarded substantial damages and an annual pension for his severe disabilities. His mother, who is now dead, said that she had taken six to eight thalidomide tablets.

The affected child has two siblings, both boys, who are normal. Her mother was well during the pregnancy, had no bleeding in the early weeks of pregnancy, and took no medication. She had three ultrasound scans.

The x ray report of the child's limbs showed