mineral density in the women with distal forearm fractures was lower after correction for age and weight than that in the population based controls (mean Z score -0.46 (95% confidence interval of the difference -0.63 to -0.29)). Bone mineral density in younger women with distal forearm fractures was lower than that in older women (ages 50 to 59, mean Z score -0.72 (-1.0 to -0.45); ages 60 to 69, -0.48 (-0.78 to -0.19); ages ≥ 70 , -0.11 (-0.41 to 0.9)). The figure shows the relation between bone mineral density and age in women with distal forearm fractures and the population based group. The intercepts ($P=0.001$) and slopes ($P=0.006$) of the regression lines were different.

Relation between bone mineral density of lumbar spine and age in women with fractures of distal forearm (■, ---; $r=0.04$, NS) and population based controls (○, —; $r=-0.24$, $P<0.0001$)

Comment

We found decreased bone mineral density of the lumbar spine in a group of women with fracture of the distal forearm, indicating an increased risk of vertebral fracture. This decrease was significant only in the younger women with distal forearm fracture. It was

notable that in the women with distal forearm fractures bone mineral density of the lumbar spine did not decrease with aging. This cross sectional design does not allow us to determine whether these subjects had low peak bone mass or had undergone accelerated bone loss.

The type I-type II classification of osteoporosis proposes that distal forearm fracture is a type I fracture. This may be an oversimplification. We found that in women under 60 distal forearm fracture is strongly associated with low bone mineral density of the lumbar spine, and in this case it may be appropriate to consider it a type I fracture. Owen *et al* found no association between distal forearm fracture in women under 60 and subsequent hip fracture, a classic type II fracture.⁴ We found that in older women, however, distal forearm fracture did not predict low bone mineral density of the lumbar spine, and Owen *et al* showed that the risk of hip fracture is doubled in women who sustain a distal forearm fracture after the age of 70. In older women distal forearm fracture may therefore be considered a type II fracture.

We concluded that it may be worth while to measure bone density in women with distal forearm fracture in the sixth decade since they seem to be at high risk of vertebral fracture.

This study was supported by a project grant from the Arthritis and Rheumatism Council (R44); NFAP is supported by the council as a clinical research fellow. We thank Miss M Cooke and Ms D Greenfield for recruiting patients, and Mrs A Johnson, Mrs S Bowles, and Mr A Milne for measuring bone mineral densities.

1 Riggs BL, Melton LJ III. Involutional osteoporosis. *N Engl J Med* 1986;314:1676-86.

2 Ross PD, Davis JW, Epstein RS, Wasnich RD. Pre-existing fractures and bone mass predict vertebral fracture incidence in women. *Ann Intern Med* 1991;114:919-23.

3 Kleinbaum G, Kupper LL. *Applied regression analysis and other multivariate methods*. North Scituate, MA: Duxbury, 1978.

4 Owen RA, Melton LJ III, Ilstrup DM, Johnson KA, Riggs BL. Colles' fracture and subsequent hip fracture risk. *Clin Orthop* 1982;171:37-43.

(Accepted 3 March 1994)

New method for measuring compliance with long term oxygen treatment

G D Phillips, N K Harrison, A R C Cummin, Jessica Ward, V S Shenoy, V Newey, D Ritchie, Isobel P Williams, F J C Millard

Chest Clinic, St George's Hospital, London SW17 0QT

G D Phillips, senior registrar
N K Harrison, senior registrar
A R C Cummin, lecturer
V S Shenoy, medical technical officer
F J C Millard, consultant physician

Department of Medical Physics
V Newey, medical technical officer
D Ritchie, medical technical officer

Department of Chest Medicine, St Albans City Hospital, St Albans AL3 5PN
Jessica Ward, health visitor
Isobel P Williams, consultant physician

Correspondence to: Dr Phillips.

BMJ 1994;308:1544-5

Long term oxygen treatment used for at least 15 hours a day improves survival in patients with chronic obstructive lung disease and respiratory failure.^{1,2} In 1989 prescriptions for oxygen for use at home, particularly oxygen concentrators, cost the NHS £18 million, and the number of such prescriptions has been increasing.³ There has never, however, been a completely reliable method for measuring compliance with domiciliary oxygen treatment. The most common method uses a clock incorporated into the oxygen concentrator, but patients may take off their nasal cannulas, and leaving the concentrator and clock running, or they may run the machine without wearing the delivery system.

We developed a method to assess compliance that uses plastic electrodes connected to a small (14.3×9.0×3.3 cm), battery operated timing device. This "electrode timer" records the length of time that the nasal cannulas are in contact with the skin (figure); it contains a non-volatile memory that enables recording for up to three weeks. The aims of the present study

Electrode timer worn by patient receiving oxygen treatment. Plastic electrodes make contact with the patient's skin and connect to a portable recording device, which stores the data and cannot be read by the patient (published with permission)

were to validate the electrode timer and to use it to assess compliance in patients with respiratory failure who were receiving long term domiciliary oxygen treatment with an oxygen concentrator and nasal cannulas.

Patients, methods, and results

In the validation study the electrode timer was used in 15 inpatients receiving oxygen treatment with nasal cannulas for acute infective exacerbations of chronic bronchitis or emphysema. The patients were monitored by nurses, who recorded how long the cannulas were worn but did not influence the positioning of the cannulas. A close correlation existed between the length of time recorded by the electrode timer and the length of time a patient received oxygen as observed by the nurses ($r=0.996$, $P<0.001$).

We then studied 56 patients who were receiving long term domiciliary oxygen treatment. All were visited in their homes, where plastic electrodes were cut to size, attached to the patients' nasal cannulas, and connected to the electrode timer. The patients were told to use their oxygen in the usual manner and that the device was to check the oxygen delivery system. One week later the timer was collected, and the times that were recorded by the concentrator's clock and by the electrode timer were noted. The study was approved by the ethics committees of the two hospitals participating in the study.

The mean (range) time for which patients were monitored was 162 (122-193) hours. According to the concentrator's clock, the mean total time for which the concentrator was switched on was 124.7 (38.1-171.5) hours which suggested that oxygen was used for a mean of 18.2 hours a day. The mean total time, however,

that patients received oxygen as determined by the electrode timer was 89.6 (27.3-162.2) hours, which suggested that oxygen was used for a mean of only 12.7 hours a day. This was a significantly shorter length of time than that recorded by the concentrator's clock ($P<0.001$), and no correlation existed between the time measurements of the clock and those of the electrode timer ($r=0.36$).

Comment

Our results suggest that the electrode timer is more accurate than a concentrator's clock for assessing patients' compliance with domiciliary oxygen treatment and that many patients seem to use oxygen for less than the recommended 15 hours a day.⁴ Previous studies that relied on the concentrator's clock to measure compliance may therefore have overestimated the daily oxygen requirement to reduce mortality in patients with respiratory failure. Our findings have potential economic implications and indicate that the daily oxygen requirement recommended to patients may need to be revised.

We thank the audit committees of St George's and St Albans Hospitals for financial support.

- 1 Medical Research Council. Long-term domiciliary oxygen therapy in chronic hypoxic cor pulmonale complicating chronic bronchitis and emphysema. Report of a working party. *Lancet* 1981;i:681-6.
- 2 Nocturnal Oxygen Therapy Trial Group. Continuous or nocturnal oxygen therapy in hypoxic chronic obstructive lung disease. A clinical trial. *Ann Intern Med* 1980;93:391-8.
- 3 Williams B, Nicholl J. Recent trends in the use of domiciliary oxygen in England and Wales. *Health Trends* 1991;23:166-7.
- 4 Department of Health and Social Security. *Introduction of oxygen concentrators to the domiciliary oxygen therapy service*. London: DHSS, 1986. (Publication No FPN 398.)

(Accepted 7 February 1994)

A BOOK THAT CHANGED MY PRACTICE

"The Evolution of Co-operation"

When I became a consultant the most striking difference was the managerial component. I suddenly found myself responsible for a service which was adjusting to the new world of contracting, clinical directorships, and trust status. All sorts of people came to see me—managers, senior nurses, purchasers, and people from social services departments—and I was uncertain of their motives. Furthermore, what commitments might I make which I would subsequently regret? Were the general psychiatrists coercing me into taking patients from them under disadvantageous arrangements? Were my nursing and other managers scheming covertly? Should I react towards general managers with mistrust or just hostility at their efforts to hack my budget? What was the professor's secret agenda?

Around this time I came across Axelrod's book, first published in 1984.¹ It analyses a situation called the iterated prisoner's dilemma. Each of two players must choose whether to cooperate with the other or not. There are therefore four possible outcomes of each move for each player: mutual cooperation, mutual defection, taking advantage of the opponent, or being exploited. The outcomes are assigned different scores. The highest score for an individual—five—is gained by defecting when the other cooperates, but mutual cooperation has a higher total score as each player gains three points. Mutual defection scores one point each, and being ripped off scores zero. The core of the book is a computer tournament of different strategies for playing the game. In two successive tournaments, a strategy known as Tit for Tat (TFT) emerged victorious. Using TFT, a player

cooperates on the first move, but thereafter copies the previous move of the other player—for example, if the opponent defaults on the first move TFT's second move is to default. What seems to make TFT successful is that it is a nice strategy—that is, it is prepared to cooperate and will always cooperate if the opponent is using a nice strategy, but that it is not easily exploited.

What has this to do with my role in NHS management? Firstly, it didn't really matter what motivated other people. If they showed they would cooperate with me it was okay to be cooperative. Secondly, there was evidence to support my intuition that being cooperative does better in the long run. Thirdly, I could see why individuals with a paranoid view of the world, while often correct, did not get things their way. Fourthly, you should not be a doormat either. All this was liberating as it swept away many of my concerns, enabling me to concentrate on the issues.

There is always a footnote. One problem with TFT was that it seemed that you could get locked into consistent defaulting if you encountered someone who did not initially cooperate. Recently, it seems that another strategy, called Pavlov or "win-stay, lose-shift," outperforms TFT.² Pavlov may be more representative of biological situations. None the less, my general conclusions are the same: be positive but respond firmly to attempts at exploitation.—TOM DENING is a psychiatrist for the elderly in Cambridge

- 1 Axelrod R. *The evolution of co-operation*. Harmondsworth: Penguin, 1990.
- 2 Nowak M, Sigmund K. A strategy of win-stay, lose-shift that outperforms tit-for-tat in the prisoner's dilemma game. *Nature* 1993;364:56-8.