committee and from A G Elder, in which in clear terms are laid out the case for screening after admission¹ and that against screening before admission.⁴ My only slight disagreement with Elder is that if he does not believe hepatitis B is the thin end of the wedge, he should look again at the lack of compelling arguments on which these current guidelines have been introduced and at Richards and Harries comment that hepatitis B may only be the "tip of the iceberg"—presumably as a criterion for exclusion from medical school.

Richards and Harries are at pains to underline the concept of responsibility. The point here is that the Committee of Vice-Chancellors and Principals has a massive responsibility. It has a responsibility to the medical student who is now halfway through his or her course, having been accepted and been held in limbo for the two years which the committee has taken to debate the matter. This student's future has now been significantly affected by the guidelines, and whereas a modified clinical course was in prospect, he or she is now being pressured to leave or change course.

The responsibility of the committee pertains particularly to medical schools, some of which are continuing their policy of screening after admission. The fear is that, in view of the justifiably high regard in which the committee is held, some schools will, against their own better judgment, exclude or dismiss outstanding students on the sole grounds that to admit them would be to go against one of the committee's pronouncements.

Admission to medical schools should be on the basis of merit and suitability to become a doctor, not on grounds of infinitesimal risk factors which are preventable. The concept of accepting an applicant who is less suited for medicine on grounds of intellect, attitude, and personality rather than someone who is more suited but is positive for hepatitis B antigen—possibly temporarily—is not in the best interests of the profession or the public, whereas constructive use of able individuals and creation of an appropriate training programme for them is. I believe the debate should continue and the Committee of Vice-Chancellors and Principals should re-evaluate its position in the light of responsible opinion, such as that provided by the BMA medical students committee.

A M L LEVER
Honorary consultant physician

Department of Medicine,
University of Cambridge School of Medicine,
Addenbrooke's Hospital,
Cambridge CB2 2QQ

- 1 Lever AML. Hepatitis B and medical student admission. *BMJ* 1994;308:870-1. (2 April.)
- 2 Richards P, Harries F. Hepatitis B and admission to medical school. *BMJ* 1994;308:1161. (30 April.)
- 3 Gauntlett R, Bailey M. Hepatitis B and admission to medical school. *BMJ* 1994;308:1161. (30 April.)
- 4 Elder AG. Hepatitis B and admission to medical school. *BMJ* 1994;308:1161. (30 April.)

Guidelines apply to dental students too

EDITOR,—A M Lever highlights the problems of implementation of the recently issued guidelines for universities regarding medical students.¹ These guidelines also apply to dental student admission. The aim of courses for dental students is to produce a graduate with academic and clinical knowledge and skills, ready for independent practice as a dental physician and surgeon. Therefore as part of these courses all students are required to carry out a wide range of treatment procedures. By the Department of Health's guidelines virtually all dental procedures are "exposure prone" in that the operation takes place within a body cavity. On qualification virtually all graduates will continue to perform clinical work as there are few jobs in dentistry which do not include clinical work.

It is appropriate to screen dental students and

start hepatitis B vaccination during the first term of training as they are unlikely to perform exposure prone procedures then. Dental schools can avoid pre-admission screening. Lever suggested that one in 5000 medical school applicants could be infectious, which would suggest that one carrier of e antigen might be accepted to dental studentship every six years—surely a surmountable problem. The cost of the vaccination programme to dental schools or students is considerable—vaccination by the student's own general practitioner necessitates travel and time costs. This is accepted as necessary to protect the patients and the dentist.

We considered the problem of the HIV positive dental student at a European Community Working Party² since employment opportunities in dentistry would be very limited for such students. On admission to dental school, students should be informed of the occupational risk and consequences for employment of becoming HIV positive. There is one reported case of a dental undergraduate who was found to be HIV positive.³ Testing should remain on a voluntary basis but once a student is identified as positive a sympathetic approach by the dental school, university, and General Dental Council should allow a clinical student to complete the course and be examined while avoiding exposure prone procedures. The graduate could then seek research or administrative work or obtain graduate entry to a non-dental course.

A recent report indicated that "over 70% of hospital ward doctors had been accidentally jabbed by 'sharps' during a two year period."⁴ The emphasis in dentistry has, for some years, been to prevent injury and possible contraction of HIV. Risk assessment⁵ has led to modification of techniques and so reduced the danger to the operator and assistant. Anecdotal evidence points to a reduction in injuries, but only time will tell how effective these changes in techniques have been in preventing dentists from contracting HIV infection.

PETER L ERRIDGE
Director

Special Needs Unit,
Guy's Dental Hospital,
London SE1 9RT

- 1 Lever AML. Hepatitis B and medical student admission. *BMJ* 1994;308:870-1. (2 April.)
- 2 Smith CJ. HIV infection and AIDS: educational and ethical aspects in relation to dentistry. *Br Dent J* 1993;175:75-7.
- 3 Comer RW, Myers DR, Steadman CD. Management considerations for an HIV dental student. *J Dent Educ* 1991;55:187-91.
- 4 Zinn C. Australian doctors fight over HIV testing. *BMJ* 1994;308:1058. (23 April.)
- 5 Health and Safety Commission. *Management of Health and Safety at Work Regulations* 1992. London: HMSO, 1992.

General practitioners and necropsies

General practitioners are welcome at necropsies

EDITOR,—Recent articles on necropsies written by the non-forensic community are muddled and consistently omit to mention crucial differences between the coroner's necropsy (under the Coroners Act 1988 and the Coroners Rules 1984) and the hospital necropsy (under the Human Tissue Act 1961, as amended by the Anatomy Act 1984).^{1,2} The relevant cofactors that merit consideration by those indulging in medical jurisprudence are stated elsewhere.^{4,5}

The comment that a deterrent to general practitioners requesting necropsies is their belief that a necropsy is needed only if the cause of death is not known¹ is wrong. When the cause of death is unknown the case is referred to the coroner and a request from the general practitioner for a necropsy is redundant.

Although the coroner's investigation of a death primarily serves statutory requirements and medicolegal interest, modern forensic pathologists endeavour to find a medical reason for the death for the benefit of clinical colleagues and the bereaved family. All too commonly, and to the detriment of clinicopathological answers that may be gleaned from a necropsy, no clinical history is available.

The general practitioner is welcome at the necropsy (under rule 7 of the Coroners Rules 1984) and entitled to be notified when practicable of the time and place. It is constructive and of mutual benefit for doctors to liaise with the coroner's pathologist before the necropsy. More communication and greater rapport should be encouraged.

The financial burden of a medicolegal necropsy falls on the coroner or the police authority and is relieved by public funds. A prescribed fee (if any) for the necropsy report (under rule 57 of the Coroners Rules 1984) is a reasonable token payment. It may selectively control properly interested applicants. The report and other evidence before the coroner may be inspected without charge.

A proposal that a copy of every necropsy report should be readily available to the general practitioner is inappropriate under current legislation. In practice, the information in many medicolegal cases can be given in a telephone call, and in certain cases premature disclosure is inadvisable, particularly if legal proceedings or criminal charges are likely and the case material is sub judice. The disclosure of information without the coroner's prior authorisation is forbidden (under rules 10 and 13 of the Coroners Rules 1984) in coroner's cases.

A necropsy should not represent failure for a doctor but should be regarded as vital to the process of continuing postgraduate education and valuable to medicine.

F PATEL
Home Office pathologist

Department of Forensic Medicine,
UMDS Guy's Hospital,
London SE1 9RT

- 1 Thomas KB, Weller RO. General practitioners and necropsies. *BMJ* 1994;308:1054. (23 April.)
- 2 Berlin A, Wagstaff R, Bhopal R, Spencer J. Postmortem examinations: general practitioners knowledge, behaviour, and attitudes. *BMJ* 1994;308:1080-1. (23 April.)
- 3 Slavin G. *The autopsy and audit. Report of the joint working party of the Royal Colleges of Pathologists, Physicians (London) and Surgeons (England)*. London: Royal College of Pathologists, 1991.
- 4 Patel F. Over my dead body: coroner's case. *Br J Hosp Med* 1991;46:11.
- 5 Patel F. The autopsy and audit: the forensic co-factor. *Bulletin of the Royal College of Pathologists* 1992;78:15-6.

May harm relationships with relatives

EDITOR,—K B Thomas and R O Weller suggest that general practitioners hesitate to request necropsies because the death of a patient starkly highlights their failure as doctors¹; the general practitioner can be seen as attempting to bolster his or her flagging self confidence by denying that the death happened. I challenge this view and suggest that the reluctance of general practitioners to request necropsies stems from two main factors.

Whereas hospital pathology departments have little to do with dead patients' relatives, these relatives are often the patients of the general practitioner concerned, who may have developed a close relationship with them. A desire not to hurt a relative's feelings may seem a trivial reason not to press for a necropsy when viewed from the point of view of a hospital (or even a university department of general practice). But to a general practitioner, pressing unwilling relatives for permission to perform a necropsy may mean knowingly causing further stress and hurt to

someone he or she has cared for for many years and may damage subsequent doctor-patient relationships with any member of the family or their close friends.

The second reason for general practitioners' reluctance to request necropsies is their wish to avoid the ever increasing threat of litigation and complaints. Should the patient be shown to have died of a condition that was not suspected before death the doctor is wide open to complaints and litigation by the relatives. The quoted figures that 15% of main diagnoses and 42% of causes of death were not confirmed at necropsy hardly make comforting reading when viewed in this context.² A doctor could find his or her career in ruins simply by requesting a necropsy that failed to confirm the antemortem diagnosis.

JOHN LEIGH
General practitioner

Washington,
Tyne and Wear NE38 9BT

- 1 Thomas KB, Weller RO. General practitioners and necropsies. *BMJ* 1994;308:1054. (23 April.)
- 2 Cameron HM, McGoogan E, Watson H. Necropsy: a yardstick for clinical diagnosis. *BMJ* 1980;281:985-8.

Transport costs unmet

EDITOR,—The editorial¹ and paper² on general practitioners and necropsies do not mention the main stumbling block to obtaining a necropsy on a patient who died in the community—namely, transport of the patient to the postmortem room. About three years ago I tried to obtain a necropsy on a patient with clinical sarcoid who died at home. There was no tissue diagnosis, and tuberculosis had not been wholly excluded. The pathology department (which was headed by the author of the report *Autopsy and Audit*³) was willing to perform the necropsy but insisted that the transport costs be paid either by me or by the patient's relatives. I am not in a fundholding practice and did not think that either source of funding for the transport was appropriate, so the necropsy was not done.

My second point relates to reports on coroner's necropsies. I have repeatedly requested written reports without success. In this area I am able to obtain only a verbal report, by telephoning the coroner's officer after the necropsy has been completed. Therefore I strongly support the proposal that necropsy reports should be sent to general practitioners routinely and preferably within a few weeks, before the patient's medical record is returned to the family health services authority.

PETER MCCARTNEY
General practitioner

St Peter's Street Medical Practice,
London N1 8JG

- 1 Thomas KB, Weller RO. General practitioners and necropsies. *BMJ* 1994;308:1054. (23 April.)
- 2 Berlin A, Wagstaff R, Bhopal R, Spencer J. Postmortem examinations: general practitioners' knowledge, behaviour, and attitudes. *BMJ* 1994;308:1080-1. (23 April.)
- 3 Slavin G. *The autopsy and audit. Report of the joint working party of the Royal Colleges of Pathologists, Physicians (London) and Surgeons (England)*. London: Royal College of Pathologists, 1991.

Service may not be available

EDITOR,—I agree that general practitioners should receive copies of coroners' necropsy reports and that more non-coroners' necropsies are necessary for medical audit and education,^{1,2} but these may be difficult to implement in practice. With the advent of purchaser-provider contracts it is wrong to assume that a service for non-coroners' necropsies is automatically available. Personal conversation with colleagues indicates that the provision of non-coroners' necropsies has largely been omitted from contracts with district health authorities with reference to non-fundholding and fundholding general practitioners. Indeed, its inclusion in such

contracts in future seems unlikely in view of increasing workloads in histology and minimal expansion of the number of pathologists. In addition, the cost of transporting a body from the place of death to the mortuary must not be overlooked; in some areas this cost has been borne by relatives.

Currently, copies of coroners' necropsy reports can be released only with expressed permission. This practice varies widely throughout Britain and warrants central legislation to allow the release of reports to general practitioners. In some areas the non-availability of coroners' necropsy reports is to a degree circumvented by the coroners' rules, which require coroners to notify the dead person's regular medical attendant of the date, time, and place of necropsy.³ Implicit in this is the desirability for general practitioners to attend the necropsy and personally observe the findings, although this is often not feasible. In my experience, however, the prime requirement of many general practitioners is rapid notification of the cause of death, so that relatives can be counselled appropriately.

As practised locally, after the necropsy has been undertaken general practitioners should be encouraged to telephone the pathologist and discuss the necropsy findings. Such communication invariably proves invaluable for both the general practitioner and the pathologist,

DAVID SLATER
Consultant histopathologist

Rotherham Hospitals NHS Trust,
Rotherham S60 2UD

- 1 Thomas KB, Weller RO. General practitioners and necropsies. *BMJ* 1994;308:1054. (23 April.)
- 2 Berlin A, Wagstaff R, Bhopal R, Spencer J. Postmortem examinations: general practitioners' knowledge, behaviour, and attitudes. *BMJ* 1994;308:1080-1. (23 April.)
- 3 Burton JD, Chambers DR, Gill PS. *Coroners' inquiries: a guide to law and practice*. Middlesex: Kluwer Law, 1985:34.

More sophisticated studies needed

EDITOR,—Anita Berlin and colleagues' questionnaire survey of general practitioners' knowledge of and behaviour and attitudes towards necropsies addresses some of the straightforward issues but few of the difficult ones.¹ Our experience supports their observation that many general practitioners know that they can request clinical (non-coroner's) necropsies, but few do so in practice. The authors' only explanation for this relates to the complexities of arranging such an unusual procedure, yet the authors claim to have assessed the attitudes of general practitioners towards necropsies. Opposition from relatives, confidence in clinical diagnoses, and the level of general interest in necropsies among doctors, including pathologists, are the most influential factors in any decision to request a necropsy,² and these were not discussed in the study.

The paradoxical finding that many general practitioners believe that more necropsies should be requested to improve medical audit but that rates of request remain low is similar to the situation that exists in some hospitals and requires further attention. Studies would be enhanced by collaboration with pathologists (none of the authors of the paper seem to be pathologists) and psychologists. Input from these informed sources would enable a more detailed analysis of behaviour, intentions, and attitudes with regard to necropsy.^{3,4} More necropsies will be requested only as a result of the requesting clinicians' belief in the practical utility of information obtained and the re-education of doctors from all disciplines.

Access to necropsy reports should be encouraged, but it is naive to suggest that this would have an appreciable influence on behaviour with regard to requests for necropsies. Hospital doctors have had constant access to necropsy reports, and rates of clinical necropsy continue to decline.⁵ The problem is far more complex than the authors

suggest and requires more sophisticated investigation.

ROGER D START
Lecturer
DENNIS W K COTTON
Reader

Department of Pathology,
University of Sheffield Medical School,
PO Box 596,
Sheffield S10 2UL

- 1 Berlin A, Wagstaff R, Bhopal R, Spencer J. Postmortem examinations: general practitioners' knowledge, behaviour and attitudes. *BMJ* 1994;308:1080-1. (23 April.)
- 2 Start RD, Hector-Taylor MJ, Cotton DWK, Startup M, Parsons MA, Kennedy A. Factors which influence necropsy requests: a psychological approach. *J Clin Pathol* 1992;45:254-7.
- 3 Birdi KS, Start RD, Cotton DWK, Bunce D. A psychological evaluation of factors influential in clinician necropsy requests. *J Pathol* 1993;170S:374A.
- 4 Sherwood SJ, Start RD, Birdi KS, Cotton DWK, Bunce D. Why do patients' relatives refuse permission for autopsies? *J Pathol* 1994;172S:149A.
- 5 Start RD, McCulloch TA, Benbow EW, Lauder I, Underwood JCE. Clinical necropsy rates during the 1980s: the continued decline. *J Pathol* 1993;173:63-6.

Women consultants and merit awards

EDITOR,—Women consultants are poorly represented among holders of merit awards (20% hold awards, compared with 37% of men). Two possible partial explanations have recently been suggested.¹ Firstly, the age distribution of women consultants, which is skewed towards younger age, means that they are less likely to have awards than their older male counterparts because the chances of having an award increase with age. Secondly, the specialities that women choose to go into particularly radiology, anaesthetics, paediatrics, public health, and psychiatry—are relatively poorly represented with regard to awards.

Age could be taken account of if the available data on the proportion of consultants with merit awards were presented by age and by sex to determine the extent to which this suggestion applies. The association of women with specialities poorly represented among merit award holders means that this is the cause of the difference between the sexes. Possibly these specialities are less well rewarded because many of the specialists in them are women, who as a group are less likely to be put forward for an award. In addition, the specialities listed are not as poorly underrepresented as might have been expected from the comment (all specialities 34.4%, paediatrics 35.4%, anaesthetics 28.7%, public health 28.5%, mental illness 27.8%, radiology 25.7%).² Probably, however, within each of these specialities a lower proportion of women than men hold awards.

Publication of more disaggregated information on holders of merit awards would help to clarify the situation and ensure that the discussion was based whenever possible on fact rather than opinion.

ALLISON STREETLY
Locum consultant in public health medicine

South East London Health Authority,
London SE1 7RJ

- 1 Beecham L. Women consultants lag behind in merit awards. *BMJ* 1994;308:1106. (23 April.)
- 2 Secretariat of Advisory Committee on Distinction Awards. Distinction awards: analysis by type of ward, specialty and percentage distribution at 31 December 1992—England and Wales. *Health Trends* 1994;20:152.

Merit awards

EDITOR,—I am a member of a regional higher awards committee. Though I approve of the merit awards system, I have several reservations. I agree with Jeffrey Tobias that all consultants are not equal and that some system is required to acknow-