follow the pilot study with a randomised trial comparing intraluminal treatment with or without external beam radiotherapy (OE04 trial). The aim is to improve the quality and duration of palliation with minimal disruption to patients' lives. The OE03 and OE04 trials have already been agreed to in principle by the MRC's Cancer Therapy Committee, pending the results of the respective pilot studies.

We urge clinicians who treat oesophageal cancer to collaborate in large randomised trials. Such trials offer the most reliable method of assessing and comparing treatment policies and so of improving the results of treatment. Copies of the protocols and further information can be obtained from the MRC's Cancer Trials Office. We are now particularly eager to recruit more centres into the OE02 trial.

R J DONNELLY

Chairman, MRC Oesophageal Cancer Working Party
Cardiothoracic Centre,
Liverpool L14 3PE

D J GIRLING

Secretary, MRC Oesophageal Cancer Working Party

P M FAYERS

Statistician, MRC Oesophageal Cancer Working Party,
MRC Cancer Trials Office,
5 Shaftesbury Road,
Cambridge CB2 2BW

1 Weil JG, Munday SJ. Management of oesophageal cancer. *BMJ* 1994;308:1103. (23 April.)

2 O'Reilly S, Forastiere A. New approaches to treating oesophageal cancer. *BMJ* 1994;308:1249-50. (14 May.)

3 Hatlevoll R, Hagen S, Hansen HS, Hultborn R, Jakobsen A, Mantyla M, et al. Bleomycin/cisplatin as neoadjuvant chemotherapy before radical radiotherapy in localised, inoperable carcinoma of the esophagus. *Radiother Oncol* 1992;24:114-6.

Equity in the NHS

EDITOR,—Margaret Whitehead's picture of health services before the establishment of the NHS reminds us of the dangers of market forces operating in health care.¹ She indicates clearly that we are now well along the road to requiring certain sections of the population to pay directly for the consequences of ill health. What causes the greatest concern, though, is that the main financial burden is falling on the most disadvantaged in society.

The continuing care of older people has in effect been taken out of the public sector and privatised. As this has been accomplished without any public debate most of the population are unaware of the profound effect it will have on financial arrangements for old age. It has been allowed to occur despite repeated statements from the Department of Health and ministerial pronouncements that long term care in the NHS, free to patients, still has a place.^{2,3} Despite the protest by the Social Security Committee in 1991,⁴ the view held by the National Association of Health Authorities and Trusts—that the provision of long term nursing care of older people is no longer a function of the health service—seems to have prevailed. More recently, Sir Duncan Nichol, when chief executive of the NHS Management Executive, confirmed that when medical assessment deems that long stay care is required on the grounds of ill health it should be provided free by the NHS.⁵ Thus the onus is on consultants who must make such assessments, but the financial consequences are likely to influence the outcome in favour of social care and discharge from hospital.

In a survey that I carried out last year on behalf of the British Geriatrics Society members were asked their opinions on issues relating to long term care. Of the 489 respondents, 196 agreed with the statement that all forms of such care should be means tested while 377 thought that some long stay beds should be retained in the NHS. This suggests that a substantial proportion of doctors who care for older people continue to favour the traditional

role of the NHS in the provision of long stay care at no financial cost.

Inequity in the NHS has now reached the point that adequate funding of acute care is possible only because of the transfer of resources from the long term care of older people. This amounts to blatant discrimination against those who are both old and chronically ill or disabled.

ALISTAIR RITCH
Consultant geriatrician

City Hospital,
Birmingham B18 7QH

1 Whitehead M. Who cares about equity in the NHS? *BMJ* 1994;308:1284-7. (14 May.)

2 Department of Social Security. *Financing of private and voluntary residential care*. London: HMSO, 1992.

3 Mawhinney B. *Caring for people: speech to IHSM/ADSS conference*. London: HMSO, 1992.

4 Social Security Committee. *Fourth report. The financing of private residential and nursing home fees*. London: HMSO, 1991.

5 Warden J. The lottery of long term care. *BMJ* 1994;308:742. (19 March.)

World Medical Association

EDITOR,—George Giri asks which organisation represents Spanish doctors in the World Medical Association (WMA), because as far as he can ascertain there is no equivalent of the British Medical Association (BMA) in Spain. He asks also whether the Spanish doctors' dues are paid by the government.

Spanish doctors are represented in the WMA by the Consejo General de Colegios Médicos de España, which like the BMA is both a founder of the WMA and a current member of the Standing Committee of Doctors of the European Union. Its membership, and its funding, is drawn from physicians of all specialties throughout Spain and it is not subsidised by the government.

Exact equivalence to the BMA is not a prerequisite for membership of the WMA, because many of the representative organisations from other countries differ from each other in certain respects—even those which may seem at first glance to be identical. To belong to the WMA an organisation must have a "membership broadly representative of physicians in that country" and it "must not be subject to, or controlled by, any office or agency of the government."

IAN T FIELD
Secretary general

World Medical Association,
PO Box 63,
01212 Ferney-Voltaire,
France

1 Giri G. World Medical Association. *BMJ* 1994;308:921. (2 April.)

Waiting times and the patient's charter

EDITOR,—Surely no one feels that seeing 90% of outpatients within 30 minutes of their appointment time is an unreasonable objective. However, to impose this particular standard in the patient's charter now will almost certainly lead to consequences which are against patients' interests in the short and intermediate term at this hospital.

In 1989, when norms were set, we were already trying to cope with a patient population close to double that recommended for the number of consultants in post. To increase our throughput would have been unsustainable. We have already had to accede to the divisive requirements of a two tier system engendered by fundholding and non-fundholding practices, which is inevitable (and undeniable) in a situation where demand outstrips the provision of services.

Now we face a challenge which is likely to prove insurmountable and has a financial penalty that

will exacerbate the problem. We will respond seriously to the challenge, but to have any hope of seeing patients within 30 minutes of their appointment time it is more than likely that we will see considerably fewer routine patients (except for those of fundholders); be refractory to requests for seeing urgent and expedited patients whom we traditionally have squeezed in; decline requests from medical students for special attachments; discourage postgraduate activities such as teaching and accommodation of medical and paramedical visitors; and be inclined to send away and rebook those patients who will not be seen "in time" because of unexpected absences, such as staff sickness.

Many of our patients, new and old, require several detailed assessments by professional support staff before seeing a member of the medical staff. Many depend on ambulances, making appointment planning optimistic. We urge the rulemakers to reconsider the simplistic requirement that patients be seen within 30 minutes of their appointment to see a doctor. If the requirement were to relate to appointments with any staff member this would help us to develop and improve the running of our paramedical support services.

If money did follow the patient, as was promised, an already highly cost effective hospital like ours could afford to deliver the exemplary, innovative, and often complex management that is properly expected of us, while fulfilling our training and research commitments.

A M ANSONS	J L NOBLE
Consultant ophthalmic surgeon	Consultant ophthalmic surgeon
N P JONES	E P O'DONOGHUE
Consultant ophthalmic surgeon	Consultant ophthalmic surgeon
B LEATHERBARROW	A E A RIDGWAY
Consultant ophthalmic surgeon	Consultant ophthalmic surgeon
K B MILLS	A B TULLO
Consultant ophthalmic surgeon	Consultant ophthalmic surgeon
D McLEOD	G S TURNER
Professor of ophthalmology	Consultant ophthalmic surgeon

Royal Eye Hospital,
Manchester M13 9WH

Elderly people who get stuck in the bath

EDITOR,—As a medical nonagenarian with bilateral hip replacements, I agree that it is often difficult and sometimes impossible to get out of a bath unaided.¹ If I lie in a normal sized bath I am unable to get out unaided. Fortunately, however, my own bath, which is old, is slightly broader than usual baths, and this allows me to roll over on to my front, from which position I can rise to my knees and then stand erect and step out of the bath.

What I cannot understand is why anyone should get seriously burnt or scalded in the bath. Surely it is dangerous to leave the hot tap running when you are in the bath. It is common sense to test the heat of the water before getting into it; a thermometer is not required.

A M CAMPBELL

Skipton,
North Yorkshire BD23 1LN

1 Cadier MA, Shakespeare PJ. Elderly people who get stuck in the bath. *BMJ* 1994;308:1240. (7 May.)

Correction

New public health

A typesetting error occurred in this letter by Nigel Bruce and others (11 June, p 1568). The sentence in the middle of the penultimate paragraph should read, "These processes—central to Health for All theory and increasingly important in national health policy"—require commitment, resources, and experience."