ledge is often one of the basic requirements when assessing quality of care.

It has been claimed that the purchaser-provider split in the health system provides more choice for patients and doctors.² For this new found freedom to be translated into improved care, those bodies responsible for buying services on behalf of patients must be large enough to have at their disposal sufficient information to make the best choice among competing provider units. The resources required for this would normally be available only to relatively large scale commissioning bodies such as health authorities or commissioning consortiums. Unfortunately, Potamitis and colleagues do not say how many of the general practitioners who responded in their study were fundholders or had been involved in commissioning ophthalmic services.

Modern techniques do not necessarily equate with improved quality of care. C M S Royston and colleagues reported that in laparoscopic surgery morbidity figures improve only when training and assessment are adequate.³ Is there not a danger that units will, in an attempt to attract potential purchasers, adopt new practices before they have been adequately evaluated? We may be left with the situation where internal market forces have allowed surgical innovation to be led from a position of relative ignorance.

MARK DAVIES
General practitioner

JANE LOGAN
General practitioner

JENNIFER LAW
General practitioner

Mawbey Brough Health Centre,
London SW8 2UD

1 Potamitis T, Fouladi M, Aggarwal RK, Jones HS, Freider AR. General practitioners' awareness of different techniques of cataract surgery: implications for quality of care. *BMJ* 1994; 308:1334-5. (21 May.)

2 Secretary of State for Health. *The health of the nation*. London: HMSO, 1991.

3 Royston CMS, Landsdown MRJ, Brough WA. Teaching laparoscopic surgery: the need for guidelines. *BMJ* 1994;308:1023-5. (16 April.)

Integrated care for diabetes

EDITOR,—The Diabetes Integrated Care Evaluation Team reports a comparison of integrated care versus conventional care for diabetes.¹ In our opinion the results of this study cannot be generalised to all diabetic patients, even if the exclusion criteria are taken into account. The mean glycated haemoglobin concentration when the patients were recruited is impressive, being nearly normal (5.3% (SD 1.4%); normal < 5.3%); it is evidence of good metabolic control before the trial. In the diabetes control and complications trial the mean glycated haemoglobin concentration at recruitment varied from 8.8% (1.6%) to 9.0% (1.5%) (normal value 5.05% (0.5%)).² After two years the value in the intensive therapy group had fallen to about 7%, thus remaining clearly above normal.

The degree of glycaemic control among non-insulin dependent patients who receive conventional care is not well known. The United Kingdom prospective diabetes study should answer this question next year. Nevertheless, in that study the median glycated haemoglobin concentration over three years varied from 6.2% to 7.2%,³ with exclusion criteria at recruitment similar to those of the diabetes integrated care evaluation study. In France, Guillausseau *et al* reported a mean glycated haemoglobin concentration of 8.4% (1.9%) in 246 non-insulin dependent patients and of 9.3% (1.9%) in 125 insulin dependent patients attending a diabetic clinic (normal value 6.2% (0.7%)).⁴ We would therefore be interested to know the training programme of the traditional hospital clinic model in the Aberdeen Health Centre. We think that the statement that "the model of integrated care for diabetes is at

least as effective as conventional hospital clinic care" applies only to diabetic patients with good metabolic control.

P CHEDIN
Medical practitioner
J CAHEN-VARSAX
Chief doctor

Department of Diabetology,
Hopital Victor Dupouy,
95107 Argenteuil Cedex,
France

1 Diabetes Integrated Care Evaluation Team. Integrated care for diabetes: clinical, psychosocial, and economic evaluation. *BMJ* 1994;308:1208-12. (7 May.)

2 Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977-86.

3 United Kingdom Prospective Diabetes Study. VIII. Study design, progress and performance. *Diabetologia* 1991;34:877-90.

4 Guillausseau PJ, Charles MA, Godard V, Timsit J, Chanson P, Paolaggi F, *et al*. Comparison of fructosamine with glycated hemoglobin as an index of glycemic control in diabetic patients. *Diabetes Res* 1990;13:127-31.

Chronic fatigue syndrome and myalgic encephalomyelitis

EDITOR,—Our recent editorial on the chronic fatigue syndrome and myalgic encephalomyelitis prompted considerable correspondence,¹ which raised issues of case definition, clinical management, and attitudes towards people with psychiatric illnesses. Sadly, many of our critics show that the editor of the *BMJ* is wrong to state in the "editor's choice" in the issue of 14 May that "only the naivest medical students think that diseases have some independent, objective reality." Medical students show greater intellectual sophistication in tackling the classification of ill defined illnesses than many patients and doctors—and particularly medical practitioners with self diagnosed myalgic encephalomyelitis.¹

Case definition—Ellen M Goudsmit¹ and Nick Anderson¹ assert that research criteria for the chronic fatigue syndrome fail to distinguish myalgic encephalomyelitis and exaggerate psychiatric associations. The best replicated research finding, however, is that patients suffer substantial emotional morbidity, whether the chronic fatigue syndrome is defined by British or, as patient groups prefer, Australian or American criteria.² All three sets of criteria can be used to identify cases on a continuum of fatigue,³ which includes myalgic encephalomyelitis. We did not cite D O Ho-Yen's prevalence study as it used an idiosyncratic definition of cases of the chronic fatigue syndrome and surveyed doctors' diagnoses rather than patients.

Management—Patients' self help groups can have valuable informative and supportive functions. In stating that we implied that this was not so, Ho-Yen and Grant fail to differentiate between self help and self help groups and between treatment and support.¹ We referred to attributional style and its prognostic implications for individual patients; overemphasis of a viral aetiology may promote demoralisation and avoidance of activity and thus contribute to a poor outcome. We are encouraged that the patients' organisations now advocate a gradual increase in activity and provide more optimistic outlooks with regard to outcome than they once did.¹ The prognosis seems to be better still for treated patients with the chronic fatigue syndrome or myalgic encephalomyelitis: 70% of longstanding cases improved with cognitive-behavioural treatment and with antidepressants as required, compared with 20% of patients who refused such treatment.¹

Attitudes—Clare Fleming claims that we concentrated on psychological slants rather than taking a holistic view.¹ All specialists need to beware of conceptualising illness in models most

familiar to themselves; but the association between the chronic fatigue syndrome or myalgic encephalomyelitis and psychiatric or psychological problems is indisputable,² even though the conditions are not entirely explicable in these terms. Karen Prince states that "what is really at issue is whether this illness is real or imaginary."³ Such dualistic attitudes to physical and mental illness from members of the medical profession, and particularly from members of the profession who suffer from the chronic fatigue syndrome or myalgic encephalomyelitis, make it all the more understandable that patients balk at suggestions that psychiatric and psychological factors are relevant. The repeated criticism of our editorial and the scientific medical journals—that they are biased towards psychiatric and away from "organic" aetiological theories—is based on isolated reports of results of brain imaging and neuroendocrine findings, which have not been replicated and have not controlled for depression and inactivity. These comments typify stereotyped views of psychiatric illnesses as not real, unrelated to cerebral disorder, and therefore the sufferers' fault—none of which is true.

Accusations of bias arise from a misunderstanding of the nature of progress in medical research. Replicated, peer reviewed findings ensure that doctors and well informed patients are not misled by preliminary reports of research or exaggerated claims in the media, examples of which are numerous concerning the chronic fatigue syndrome and myalgic encephalomyelitis. Constructive debate with the patients' organisations, based on scientific evidence, is part of the process of improving the treatment and outlook for sufferers. Doctors must learn from their patients' experiences, but the myalgic encephalomyelitis societies should not try to set the research agenda or shout down views with which they disagree. The editors of medical journals could equally be accused of bias towards the self appointed patients' representatives in printing their simplistic criticism of evolving knowledge in these complex conditions.

S M LAWRIE
Lecturer

Department of Psychiatry,
Royal Edinburgh Hospital,
Edinburgh EH10 5HF

Hairmyres Hospital,
East Kilbride, Glasgow G75 8RG

A J PELOSI
Consultant psychiatrist

1 Correspondence. *BMJ* 1994;308:1297-301. (14 May.)

2 David AS. Postviral fatigue syndrome and psychiatry. *Br Med Bull* 1991;46:966-88.

3 Bates DW, Schmitt W, Buchwald D, Ware NC, Lee J, Thoyer E, *et al*. Prevalence of fatigue and chronic fatigue in a primary care practice. *Arch Intern Med* 1993;153:2759-65.

4 Bonner D, Ron M, Chalder T, Butler S, Wessely S. Chronic fatigue syndrome: a follow-up study. *J Neurol Neurosurg Psychiatry* 1994;57:617-21.

Alcohol and blood pressure

EDITOR,—The INTERSALT group addressed the relation between alcohol intake and blood pressure and advocates a reduction in population mean alcohol intake. It concludes, "Overall, the data indicate the usefulness of an approach targeting those at high risk as well as the general population to reduce the adverse effects of alcohol on blood pressure."¹ The group also states that a reduction in systolic blood pressure of 5 mm Hg achieved by "favourable lifestyle changes" would reduce coronary mortality by 9%. Although agreeing with this statement in general, I doubt whether it applies to reductions in pressure achieved by limiting alcohol consumption.

Men who are light to moderate drinkers (consuming 1-299 ml/week) may reduce their blood pressure by 1 mm Hg by stopping drinking, but epidemiological evidence suggests that their coronary mortality may increase by up to 10%.²

Male drinkers consuming ≥ 300 ml/week may experience a 4 mm Hg fall in systolic pressure and possibly a 27% fall in coronary mortality. But heavy drinkers constituted only about 20% of the men studied, light to moderate drinkers about 50%, and non-drinkers 30%. Thus, for men it is theoretically possible to increase cardiac mortality by 10% in half the population and to reduce it by 27% in a fifth. With a population approach in women 45% may experience an increase in coronary mortality and 5% a decrease. The net result is that lowering blood pressure by reducing alcohol intake is unlikely to reduce coronary mortality.

All drinkers should be advised of the increased risks of non-cardiovascular death, but only heavy drinkers should be advised of the increased risk of coronary death.

C J BULPITT
Professor

Division of Geriatric Medicine,
Royal Postgraduate Medical School,
Hammersmith Hospital,
London W12 0NN

1 Marmot MG, Elliott P, Shipley MJ, Dyer AR, Ueshima H, Bevers DG, *et al.* Alcohol and blood pressure: the INTER-SALT study. *BMJ* 1994;308:1263-7. (14 May.)

2 Klatsky AL, Armstrong MA, Friedman GD. Alcohol and mortality. *Ann Intern Med* 1992;117:646-54.

Managing clinical risk

EDITOR,—Jonathan Mant and Alex Gatherer aver that "risk management is concerned with the quality of care only as much as better quality care might reduce negligence claims." Sadly, few involved with risk management programmes see them in any other way. That is why Action for Victims of Medical Accidents insists that risk management paradoxically carries risk for patients. If the main objective is to reduce negligence claims rather than to improve patients' care a conflict of interest between hospitals and patients cannot be avoided. For that reason it is essential that risk management programmes are not left to commercially minded consultants and managers but involve both doctors' and patients' organisations.

ARNOLD SIMANOWITZ
Executive director

Action for Victims of Medical Accidents,
London SE23 3TP

1 Mant J, Gatherer A. Managing clinical risk. *BMJ* 1994;308:1522-3. (11 Jan.)

Log books for doctors

EDITOR,—In commenting on the case of Dr Behrooz Irani, D J Dye suggests that locum staff of any grade should be under the supervision of a senior member of consultant staff and should carry a log book.¹ No one disputes that all doctors who are negligent in their professional duties should be dealt with appropriately. But Dye seems to be selective and ignores negligence among doctors in substantive posts. The reference system used for locum doctors is the same as that used for doctors in substantive posts. So why doesn't Dye suggest that the same measures should be used for doctors in substantive posts to eliminate such problems? I wonder whether Dye would be happy to be supervised by a senior member of staff and would like to carry a log book because of negligence by some doctors holding substantive posts. When a time sheet was proposed for all the consultants working in Medway Health Authority a couple of years ago consultants' leaders circulated letters to all the consultants to oppose this.

Locum doctors practise the same medicine and are trained in the same NHS hospitals as doctors in substantive posts. But the most important

differences are that locums lack recognition of their work and have no career prospects and are being exploited in this respect. Any measures that do not address these fundamental defects are bound to fail. I suggest that Dye and any other people interested in improving locum services should refer to my previous letter on this subject.²

R MACHERLA
Consultant physician

Medicine for the Elderly,
St George's Hospital,
London SW17

1 Dye DJ. Locum doctors should have log books. *BMJ* 1994;308:1572. (11 June.)

2 Macherla R. Locum doctors are undervalued. *BMJ* 1993;307:1212.

Network of cancer services proposed

EDITOR,—It is bizarre that the chief medical officer of the NHS, Dr Kenneth Calman, should propose "a national network of designated cancer units in district hospitals linked to specialised cancer centres in larger general hospitals."¹ As the experienced chairman of a sensible committee Dr Calman may propose, but as a member of the NHS Executive Dr Calman cannot dispose. The internal market that he and his colleagues on the executive have created make the proposal for a national plan for cancer services useless.

The events at St Bartholomew's Hospital have shown that Dr Calman's well planned cancer center in any major hospital could be closed down weeks after opening if the hospital in which it was based did not attract enough business within the market. Within the two tier system created by general practice fundholding "general practitioners may be expected to establish local referral and follow up patterns with providers of specialised cancer care, using national agreed guidelines,"² but only patients of non-fundholding general practitioners (a dwindling number) can be guaranteed to be referred to the local centre. Fundholding general practitioners, driven by Dr Calman's market, are bound to look over the border of their district to see if neighbouring or even more distant district cancer services can provide cheaper care.

Professor Selby states, "It was fairly obvious to the group that clinical freedom was not producing the best cancer services"; at least Dr Calman's report could lead to a "uniformly high standard of care."³ market freedoms, however, can create only a framework for chaos.

"The reality of change" reported by Jane Smith⁴ is a reality that Dr Calman seems to lecture others to accept but to be incapable of accepting himself.

RICHARD STONE
Former general practitioner

London NW8 0LU

1 Dillner L. Cancer services to be radically developed in Britain. *BMJ* 1994;308:1317. (21 May.)

2 Smith J. A rebuke from the chief medical officer. *BMJ* 1994;308:1322. (21 May.)

Nestlé's donation

EDITOR,—The donation from Nestlé (UK) Ltd of £140 000 over four years¹ is the largest single donation ever received by the British Paediatric Association and should have been debated among the membership. The infant mortality and morbidity worldwide which results from bottle feeding is well known,² as is the WHO code of practice,³ which is supported by a European directive and, in principle though not yet in law, by the British government. The British Paediatric Association stated its support for this code to be

enacted in law in 1990 and 1994. The code prohibits promotion of formula milk and donation of free samples and supplies and requires any financial contributions from formula milk companies to be declared. Nestlé has the largest share of formula milk sales in developing countries and, according to International Baby Food Action, continues to breach the code and to contravene government policies in less developed countries and in Europe: hence the international boycott of Nestlé products supported by many organisations.¹

It is true that a number of paediatric research projects are funded, directly or indirectly, by Nestlé (UK) Ltd. However, individual paediatricians can choose whether to work with such projects. In contrast, the British Paediatric Association is our only professional body; we have no choice but to back it.

At the time of their donation in July 1993, Nestlé (UK) Ltd requested that acceptance be confirmed by the British Paediatric Association's council as soon as possible. Despite this request, the source of the donation was not revealed to council members at meetings in October 1993 and February 1994 and was not raised at council until June 1994, following a letter to the president from Dr Anthony Costello. The accounts published in the association's annual report in 1994 cited an anonymous donation of £35 000 for the year 1993-4.

These successive lapses have been explained as a series of oversights by the association's administration. As members of the British Paediatric Association we ask the president, firstly, to explain fully to the membership why the donation was accepted covertly, and, secondly, to give an assurance that in future, the association's administration will be properly accountable to the membership. Nestlé funding compromises the British Paediatric Association's standing as a national and international professional body able to comment on all matters related to child health. The Nestlé donation should be returned and alternative sources of funding sought.

NELLIE ADJAYE, J R BEESLEY, NIGEL BREWSTER,
A BUSH, P F B CARTER, ELAINE CARTER,
MICHAEL CHAN, NNENNA COOKEY,
MALCOLM G COULTHARD, SIMON COURT,
DAVID B CUNDALL, GEOFF DEBELLE,
CAROL DEZATEUX, G J EBRAHIM, DAVID ELLIMAN,
D GIBB, RUTH GILBERT, EDMUND HEY, W T HOULSBY,
HELEN ISSLER, CHRISTINE JESSEN, P M JONES,
W H LAMB, H J LAMBERT, CAMILLE LAZARO,
SIMON LENTON, MICHAEL LEVIN, M F LOWRY,
STUART LOGAN, AIDAN MACFARLANE,
ROSEMARY MENZIES, D C MORLEY, H M PALMER,
CATHERINE PECKHAM, C A PORTER, EUAN ROSS,
A N P SPEIGHT, NICK SPENCER, J P STANFIELD,
J E THOMAS, ANDREW TOMKINS, TONY WATERSTON,
C M WRIGHT, JANE WYNNE, PAM ZINKIN

Members of British Paediatric Association,
London NW1 4LB

1 Court C. Paediatricians question Nestlé's donation. *BMJ* 1994;308:1661-2. (25 June.)

2 Cunningham AS, Jelliffe DB, Jelliffe EFP. *Breastfeeding, growth and illness: an annotated bibliography*. New York: UNICEF, 1992.

3 World Health Organisation. *International code of marketing of breast milk substitutes*. Geneva: WHO, 1981.

Your letter should be less than 400 words long, contain a maximum of five references including one to the *BMJ* article to which you refer, and be typed with double spacing. All authors need to sign the letter and provide one current appointment and address. Your letter may be shortened. We encourage you to declare any conflict of interest that you may have. Please enclose a stamped addressed envelope if you require an acknowledgment.