"computer illiterate" for some years yet. Computer skills should be integrated into all postgraduate courses. For example, we would add "information management including computer skills" to the content of the master's degree in general practice suggested by Lindsay F P Smith.⁴ Students taking the master of public health at Glasgow University are now required to attain basic computer literacy.

Secondly, the need for an appropriate bridge between JANET (the joint academic network, the British end of the internet) and NHS networks has been discussed in seminars as part of the "Cumberlege process."⁵ Given the rate of growth of the internet, a well planned solution is needed urgently to prevent the growth of unofficial and uncontrolled connections.

RAY JONES
Senior lecturer in health informatics

Department of Public Health,
University of Glasgow,
Glasgow G12 8RZ

SUE KINN
Research fellow

Department of Postgraduate Medical Education,
University of Glasgow

- 1 LaPorte RE, Akazawa S, Hellmonds P, Boostrom E, Gamboa C, Gooch T, et al. Global public health and the information superhighway. *BMJ* 1994;308:1652-3. (25 June.)
- 2 Jones RB, Navin LM, Barrie J, Hillan E, Kinane D. Computer literacy among medical, nursing, dental and veterinary undergraduates. *Med Educ* 1991;25:191-5.
- 3 Kidd MR, Connolly GL, Cesnik B, McPhee W. What do medical students know about computers? *Med J Aust* 1993;158:283-4.
- 4 Smith LFP. Higher professional training in general practice; provision of master's degree courses in the United Kingdom in 1993. *BMJ* 1994;308:1679-82. (25 June.)
- 5 *Managing the knowledge base of healthcare. Report of a seminar held on 22nd October 1993 at the King's Fund Centre, London.* London: British Library Research and Development Department, 1994. (Report 6133.)

Computer linked fax system is efficient

EDITOR,—The chief medical officer recently stated that an information cascade system should be established to transmit messages to doctors.¹ Consequently, district health authorities, family health services authorities, NHS trusts, directly managed units, and other health care organisations have had to consider how to disseminate information within their organisation and to other organisations. In the event of a major incident or communicable disease outbreak—and for routine administration—an efficient communication system is called for.

We wish to highlight the potential use of a computer linked facsimile (fax) system as an efficient means of transmitting the same message to several destinations. This facility, available with many computer software packages, requires each recipient to have access to a fax machine. The sender can import a word file with the message into the fax linked software. This merges the file with the fax numbers of intended destinations selected from a file containing the fax numbers of key agencies. Computer activation of the sender's fax results in the automatic transmission of the message to target fax machines. This obviates the need to address each message separately and to load individual sheets manually into the sender's fax machine.

This means of communication is suitable for use during working hours. Out of hours a recipient may be alerted to receipt of a fax by a tonemaster bleep activated through a computer modem linked to the recipient's fax machine. Acknowledgement of receipt of the message could be faxed back to the sender.

The time saved by use of this facility is of benefit in both urgent and routine situations. The wide availability of fax machines makes this method of communication between agencies (both within and outside the NHS) broadly applicable and independent of more complex interagency information systems, such as EPINET (the Public

Health Laboratory Service Communicable Disease Surveillance Centre communication system) and electronic mail.

JOYSHRI SARANGI
Registrar in public health medicine
GILLIAN EVANS
Senior registrar in public health medicine

Bristol and District Health Authority,
Bristol BS2 8EE

- 1 Calman K. *Communications with the profession*. London: DoH, 1994. (PL/CMO(94)3.)

Global health network university proposed

EDITOR,—Recently we pointed out the potential of the information superhighway to become a critical factor in improving global health in the next century.¹ One important aspect of the telecommunications revolution is to enable distance education. We believe that it is time to establish a global training programme in public health through the capabilities that the internet provides and are enlisting faculty and students to join this effort.

Developing countries are facing the twin burdens of chronic disease and infectious disease. As the result of increased longevity, chronic diseases have emerged, yet infectious diseases have not been eliminated. Recently a World Bank report argued that cost effective public health measures represent a primary means for disease prevention and a reduction of the enormous cost of disease.²

The establishment of public health measures requires trained public health staff. Many developing countries have no choice but to send students abroad for training. This is enormously expensive—tuition, fees, and stipends at an American school of public health for a two year MPH degree cost over \$80 000, but unfortunately the costs are higher as more than 50% of students do not return home after completing their degree. Thus, increasing the number of foreign trained MPH staff in a country requires an investment of over \$160 000. In addition, we are in a global community of health, yet our students in developing and developed countries alike have little international experience.

Until now, a student could acquire the public health tools of access to information, the network of students, the availability of leading scientists, and international experience by few means other than through foreign training. Now, however, potentially equivalent international expertise and public health care education can be provided through the internet, which can reach broad audiences and avoid relocation and other unnecessary expenses.

We are in the process of establishing a telecommunications based public health training program. This program will grant degrees in public health, as well as provide internationally recognised certificates for those students receiving their public health degrees from their home institutions. Students can also take courses to gain international experience. We are seeking expert members of faculty of public health in schools, ministries of health, and other areas who could serve as mentors and instructors for a new "metaschool" of public health that could train students through the internet on the basis of how we learn through networks.³

The first step of this is to identify potential faculty in epidemiology and other disciplines from schools of public health who are facile on the internet who would like to teach in this metaschool. The second step is to network, across country boundaries, young people starting their training in public health so as to start a global training dialogue among students themselves and the faculty. The third step is to begin the degree granting programme.

Students and faculty interested in forming a global health network university (GHNet-U), please contact one of us below.

RONALD E LAPORTE
Director,
Disease Monitoring and Telecommunications

WHO Collaborating Center,
Graduate School of Public Health,
University of Pittsburgh,
Pittsburgh, PA, USA
(rlaporte@vms.cis.pitt.edu)

SHUNICHI AKAZAWA

World Health Organisation,
Geneva, Switzerland
(akazawa@who.ch)

EUGENE BOOSTROM

World Bank,
Washington, DC
(eboostrom@worldbank.org)

MIGUEL CAMPOS

Universidad Peruana,
Cayetano Heredia,
(viper@upch.edu.pe)

CARLOS GAMBOA

Pan American Health Organisation,
Washington, DC
(gamboa@nlm.nih.gov)

TONY GOOCH

Dumfries, VA
(tgooch@usaid.gov)

HONG-KYU LEE

Seoul National University,
Seoul, Korea
(hklee@cd4680.snu.ac.kr)

INGRID LIBMAN

University of Pittsburgh,
Pittsburgh, PA
(iml@vms.cis.pitt.edu)

ERIC MARLER

North Tarrytown, NY
(gmarler@rhcvm15.vnet.ibm.com)

KEN ROKO

US Agency for International Development,
Rosslyn, VA
(kroko@usaid.gov)

FRANCOIS SAUER

Digital Corporation,
Dearfield, FL
(sauer@lacv01.enet.dec.com)

NAOKO TAJIMA

Jikei University,
Tokyo, Japan
(ntajima@jpnst01.bitnet)

WALTER WIEBE

Federal Networking Council,
National Science Foundation
(wwiebe@nsf.gov)

- 1 LaPorte RE, Akazawa S, Hellmonds P, Boostrom E, Gamboa C, Gooch T, et al. Global public health and the information superhighway. *BMJ* 1994;308:1651-2. (25 June.)
- 2 World Bank. *Development report, 1993: investing in health*. Oxford: Oxford University Press, 1993.
- 3 Riel M. Global education through learning circles. In: Harasim L, ed. *Global networks*. London: MIT Press, 1993: 221-6.

Necrotising fasciitis

Use of skin flaps needs experience

EDITOR,—H Cox and colleagues seem to misunderstand the circulation of the skin.¹ The primary problem in necrotising fasciitis relates not just to the underlying fascia but also to the subcutaneous tissues. Any plastic surgeon will have seen cases in which the underlying subcutaneous tissues become necrotic. When this dead tissue is excised the skin above becomes necrotic because the feeding blood supply based on the fasciocutaneous perforated system has been removed. If excision of the necrotic tissue is adequate it will include both the fascia and the subcutaneous tissue, which leaves the skin to survive only on its dermal vascular network. The length and breadth of the pedicle are therefore severely compromised, particularly in areas where fasciitis is most common (the groin, perineum, abdomen, and leg).

It is therefore dangerous to suggest the use of skin flaps by those who are not experienced or trained in the technique. It is essential that patients